

TABLE OF CONTENTS

INTRODUCTION

	Test Objectives and Procedures	1
	1998 Statewide Growing Conditions	2
	Variety Characterization	3
RESU	JLTS: ALFALFA PERFORMANCE TESTS	
	NORTHEASTERN KANSAS Riley County, dryland Table 1	1
	NORTH CENTRAL KANSAS Republic County, dryland Table 2	5
	SOUTHEAST KANSAS Labette County, dryland Table 3	5
	SOUTH CENTRAL KANSAS Reno County, dryland Table 4	7
	SOUTHWESTERN KANSAS Finney County, irrigated Table 5	3
APPE	ENDIX	
	Entrants in the 1998 Kansas Alfalfa Performance Tests with unverified fall dormancy and pest resistance ratings	9
	Electronic Access and University Research Policy	3

1998 KANSAS ALFALFA PERFORMANCE TESTS

INTRODUCTION

TEST OBJECTIVES AND PROCEDURES

The Kansas Agricultural Experiment Station established an official alfalfa performance testing program in 1980 to provide Kansas growers with unbiased performance comparisons on alfalfa varieties marketed in the state. Each year, private companies are asked to enter varieties voluntarily the locations slated establishment that year. Announcements and entry forms are mailed to private companies in June for entry in fall-seeded tests. Companies enter varieties of their choice and pay entry fees to cover part of the costs of conducting the tests. Most tests are planted in mid-August or September; however, the Southeast Kansas test usually is planted in the spring. Individual tests are conducted for a minimum of 3 or 4 years. New tests are established during the final production year of the previous test.

Alfalfa tests are currently in progress at 7 locations around the state. This year, no results are included from the Sandyland Experiment Field near St. John or the Cornbelt Experiment Field near Powhattan because of stand establishment problems or delays. The other testing sites include the Southwest Research-Extension Center at Garden City, the Southeast Agricultural Research Center at Parsons, the South Central Kansas Experiment Field near Hutchinson, the North Central Experiment Field near Belleville, and the Agronomy North Farm at Manhattan.

Descriptive information is presented with the results for each test (Tables 1-5). This information, including soil type, establishment methods, fertilization, pest control, irrigation, harvest dates, and growing conditions unique to that location, can help explain test and/or variety performance.

FORAGE YIELDS were estimated by harvesting four replications of each variety with a plot

harvester. The amount of forage produced from a specific area (35-80 ft²) was weighed, and a subsample was taken to determine moisture content. This information was used to convert the plot weights to tons of dry matter per acre for each cutting, the season total, and the total for each previous season as presented in Tables 1-5. The forage yield over the lifetime of a particular test is presented as the total tons of dry matter produced per acre, as the total tons of 15% moisture hay, and as a percentage of the test average.

At the bottom of each column, the <u>Least Significant Difference</u> (LSD) is listed at the 0.05 and 0.20 levels. These values indicate how large a difference is needed to be confident that one variety is superior to another. Differences between varieties that are equal to or greater than the 0.05 LSD have a 1 in 20 chance of not being real. Differences equal to or greater than the 0.20 LSD have a 1 in 5 chance of not being real.

The Coefficient of Variability (CV) provides an estimate of the consistency of the results of a particular test. In these tests, CV's below 10% generally indicate reliable, uniform data, whereas CV's of 10-15% are not uncommon and generally indicate that the data are acceptable for rough comparisons. Tests with CV's over 15% may still be useful, but variety comparisons lack precision.

The Mean Coefficient of Variability (MCV) is similar to the CV in that it serves as an indicator of test precision. The MCV is calculated by dividing the 0.05 LSD by the test mean (average) and multiplying by 100. The MCV reveals the percent difference required to detect differences between varieties with 95% confidence. Many alfalfa breeders and testers agree that tests with MCV values greater than 10% are of no benefit.




Figure 1. Statewide alfalfa harvest progress.

1998 STATEWIDE GROWING CONDITIONS

Alfalfa harvests were generally ahead of normal during the entire season (Figure 1). Warm temperatures early in May enabled producers to start the first harvest ahead of schedule, and low rainfall in the second half of the month facilitated a rapid finish (Figures 2 and 3). Continued warm temperatures caused the second and third cuttings to be slightly ahead of average as well. The fourth cutting started slightly behind last year's but progressed more rapidly because of very dry conditions in late August and all of




Figure 2. 1998 Kansas weekly maximum and minimum temperatures.




Figure 3. Statewide topsoil moisture status.

September. (From Crop-Weather reports, Kansas Agricultural Statistics, Topeka).

Entomologists found several insect pests in alfalfa fields in 1998. Weevils caused extensive damage in the eastern two thirds of the state before the first harvest. Several producers applied insecticides to control weevils in late April and early May. Pea aphids prompted insecticide applications in some south-central fields during the same period. Grasshoppers caused some concern in June and July, especially in south central Kansas, much of which suffered from drought conditions. Gray and black blister

beetles appeared sporadically beginning in mid-June through September. Several species of insect larvae damaged alfalfa late in the season (beet armyworm, fall armyworm, yellow striped armyworm, green cloverworm, corn earworm, various webworms). These worms completely destroyed some new plantings. (From Cooperative Economic Insect Survey reports, Kansas Department of Agriculture).

No significant disease problems were reported in 1998. The lack of disease pressure was most likely due to the relatively hot, dry weather in May when most of the foliar diseases develop. (From Plant Disease Survey Reports, Kansas Department of Agriculture).

The November 10 Kansas Agricultural Statistics report predicted total 1998 alfalfa hay production of 4.18 million tons from 950,000 acres. This is up from 3.6 million tons produced from 900,000 acres in 1997. The predicted average yield of 4.4 tons per acre exceeded the final 1997 average by 0.4 tons/acre.

Contact Kraig Roozeboom for alfalfa test information on disk or via e-mail. Text and tables can be sent in a variety of formats (txt, Excel, dBase). See note on page 13 about electronic access.

VARIETY CHARACTERIZATION

For variety selection, producers should consider the performance of a variety in each of the current tests where it appears, its performance over time and locations relative to familiar or check varieties, and the disease and insect resistance characteristics that are potentially important in their situation. Tables 1-6 contain updated yield data from individual tests currently in progress. The appendix contains additional descriptive information and marketing contacts for all varieties included in the 1998 Kansas Alfalfa Performance Tests. Fall dormancy, disease resistance, and insect resistance ratings were provided by developers of each variety and were reviewed by the Association of Official Seed Certifying Agencies (AOSCA) National Alfalfa Variety Review Board (NAVRB). The Alfalfa Council uses that information to publish its annual Fall Dormancy & Pest Resistance Ratings for Alfalfa Varieties, which was used as the source of the information in the appendix. Entrants provided ratings for those varieties not included in the Alfalfa Council publication.

Fall dormancy values are based on the fall canopy height measured in Minnesota. Dormancy values often are related to the speed of regrowth. The rapid regrowth types have higher values, and the slower regrowth types have lower values.

TABLE 1. RILEY CO. ALFALFA PERFORMANCE TEST RESULTS, 1995-1998.

	Forage Yield												
		Plant	t				to	ns/acre					95-98
	H	Heigh	ıt				Dry M	atter				Total,	Total,
	i	inche	s		19	998		1997	1996	1995	95-98	15%	% of
BRAND	NAME	8-18	5-11	6-17	7-13	8-18	10-21	Total Total	l Total	Total	Total	Moist	Mean
Released Cultivars													
Dairyland Seeds	Magnum IV	21	2.19	1.28	1.21	2.30	1.14	8.12 11.3	7 7.61	7.05	34.15	40.18	107
Garst	630	21	2.20	1.39	1.08	1.81	0.91	7.39 11.2	9 7.68	7.47	33.83	39.80	106
Hobart Seed	SuperCuts	21	2.24	1.25	1.08	1.80	0.97	7.35 11.2	6 7.05	7.34	33.00	38.82	103
NK	Ciba 2444	20	2.21	1.17	1.04	1.59	0.96	6.96 11.2	7.66	7.00	32.82	2 38.61	103
Cal/West	OK49	23	2.05	1.29	1.02	1.69	1.02	7.08 10.8	4 7.53	7.26	32.71	38.48	102
DSS	Reward	22	2.12	1.36	1.11	1.85	1.11	7.54 10.4	6 7.52	7.10	32.62	38.38	102
W-L Research	WL 323	21	1.99	1.34	1.01	2.03	1.00	7.37 10.9				2 38.38	
America's Alfalfa	Archer	22	2.03	1.29	1.10	1.64	1.10	7.16 10.7				38.08	
KS AES & USDA	Riley	21	2.07	1.15	1.10	2.03	1.04	7.39 10.9				38.07	
DeKalb	DK 133	20	1.77	1.31	1.03	2.42	1.40	7.93 10.1				37.98	
Star	Asset	19	2.01	1.09	0.89	1.45	1.14	6.59 11.2	-			37.81	-
America's Alfalfa	Aggressor	22	2.06	1.27	1.03	1.73	1.04	7.13 10.6				37.53	
Garst	645	21	2.32	1.27	0.97	1.71	0.96	7.23 10.6	-			37.45	
Cargill	Crown II	22	1.95	1.14	1.01	1.82	1.04	6.97 10.4				37.44	
Mycogen	TMF Generation		2.17	1.19	0.91	1.95	0.88	7.09 10.7			31.58		
NE AES & USDA	Perry	21	1.80	1.06	0.93	1.71	1.02	6.53 10.7				36.66	
America's Alfalfa	Apollo Supreme	22	2.36	1.09	0.91	1.83	0.87	7.05 10.5				36.65	
Star	A-100	21	1.97	1.23	0.84	1.64	1.08	6.76 10.7				36.59	-
W-L Research	WL 322 HQ	22	2.13 2.03	1.17	1.03	1.90	0.91	7.14 10.6				36.56	
KS AES & USDA NK	Kanza Fortress	22 22	1.93	0.87 1.06	0.91 0.83	1.75 1.56	0.82 0.99	6.38 10.4 6.38 9.88				36.27 34.12	
Experimental Strain			1.93	1.06	0.63	1.56	0.99	0.30 9.00	0.29	0.40	29.00	34.12	91
Pioneer	90W3PR1 Exp	23	2.34	1.16	0.90	1.72	1.04	7.17 11.2	1 8.06	8 12	34.56	3 40 66	108
Pioneer	91112PJ1 Exp	22	2.08	1.46	1.05	1.83	1.05	7.48 11.4				38.88	
ABI	ABI 9142	20	1.98	1.15	0.90	1.71	1.08	6.82 11.0	-			38.60	
ABI	ABI 9141 Exp	20	2.15	1.20	1.16	1.85	1.12	7.49 11.2			32.65		102
MBS	PGI3212 Exp	20	2.06	1.31	1.06	1.80	0.96	7.19 11.2	_			38.34	
ABI	ABI 923DD Exp	21	2.21	1.19	1.13	1.93	0.99	7.46 11.0		-		37.65	
Pioneer	91CO2PR1 Exp	21	2.21	1.19	0.98	1.69	0.87	6.94 11.0				37.47	
Pioneer	88C2PI2 Exp	22	1.87	1.36	1.17	2.02	0.99	7.41 10.1				36.52	
Cal/West	1346 Exp	20	2.02	1.28	1.02	1.72	0.96	6.99 10.1	3 7.33	6.50	31.00	36.47	
Pioneer	91CO1PR1 Exp	22	1.93	1.10	1.06	1.71	0.94	6.75 10.1	7.60	6.30	30.75	36.18	96
Cal/West	1344 Exp	20	2.30	1.07	0.82	1.68	0.92	6.80 10.2	0 6.68	6.96	30.64	36.05	96
Cal/West	1469 Exp	21	1.93	1.03	1.09	1.93	0.94	6.92 9.99	7.17	6.44	30.52	2 35.91	96
MBS	PGI3392 Exp	21	1.79	1.18	1.00	1.78	0.90	6.65 10.0	3 6.80	6.78	30.26	35.60	95
Summary Statistics	5												
Average	Average	21	2.07	1.21	1.01	1.81	1.01	7.10 10.73	3 7.10	7.00	31.93	37.56	100
LSD(0.05)	LSD(0.05)	1	0.17	0.18	0.16	0.24	NS	0.56 0.65	0.34	0.60	1.25	1.47	4
LSD(0.20)	LSD(0.20)	1	0.13	0.14	0.12	0.19	0.15	0.44 0.43	0.22	0.46	0.80	0.94	3
CV(%)	CV(%)	6	6.95	12.64	13.44	11.53	15.96	6.68 4.31	3.45	7.27	2.73		
MCV(%)	MCV(%)	7	8.17	14.78	15.72	13.52	18.68	7.85 6.10	4.79	8.57	3.91	3.91	4
LOCATION: Northea	ast Kansas		1998 F	ERTIL	IZATIO	N:		1998 C	ONDIT	ONS:			
Site: Agronomy	North Farm		None;	soil te	sted hi	gh in P	and K	The fire	st harve	est was	made	early to	,
County: Riley						<u>.</u>		preven	exces	sive we	eevil da	mage.	
Town: Manhattan				PEST C				Second					sed
Soil: Smolan sil				icide a				by hot,					
ESTABLISHMENT:			contro	l weev	iis and	ieat no	ppers.						
								August					
3/17/94 ; RCBD, 4	•							the 4th and Se					
Plots 3'x12'; 3'x12'	narvested							reduce				owiii a	iiu
15 lb seed/acre								reduce	a Juli C	atting)	ri c ius.		

TABLE 2. REPUBLIC CO. ALFALFA PERFORMANCE TEST RESULTS, 1998.

		Forage Yield								
		tons/acre								
						Total,	Total,			
BRAND	NAME	5-28	1998 D 7-13	ry Matter 8-15	Total	15% Moist.	% of Mean			
Released Cultivars		0 20	0	0.0						
AgriPro	Dominator	2.98	1.51	1.95	6.44	7.58	109			
DeKalb	DK 127	2.74	1.53	2.02	6.29	7.40	106			
DeKalb	DK 142	2.81	1.64	1.83	6.28	7.39	106			
Germains	WL 324	2.81	1.62	1.78	6.21	7.31	105			
Pioneer	5454	2.76	1.44	1.87	6.07	7.14	103			
AgriPro	Depend+EV	2.85	1.43	1.78	6.06	7.13	102			
Star	Asset	2.75	1.62	1.66	6.04	7.11	102			
Star	Spur	2.76	1.49	1.77	6.03	7.09	102			
Germains	WL 325 HQ	2.69	1.54	1.77	6.01	7.07	102			
NE AES & USDA	Perry	2.56	1.45	1.71	5.72	6.73	97			
KS AES & USDA	Kanza	2.04	1.46	2.00	5.50	6.47	93			
Experimental Strai										
ABI	ZN9646 Exp	2.92	1.37	1.45	5.74	6.75	97			
ABI	ZN9541 Exp	2.80	1.39	1.46	5.65	6.65	95			
ABI	ZN9540 Exp	2.68	1.42	1.54	5.64	6.64	95			
ABI	ZC9641 Exp	2.25	1.19	1.65	5.09	5.99	86			
Summary Statistic	s	2.00	4 47	4.75	F 00	0.00	100			
Average		2.69	1.47	1.75	5.92	6.96	100			
LSD(0.05)		0.19	0.14	0.21	0.33	0.39	6			
LSD(0.20)		0.15	0.11	0.16	0.26	0.31	4			
CV(%)		6.03	7.81	10.11	4.71					
MCV(%)		7.18	9.31	12.03	5.60	5.60	6			

LOCATION: North Central Kansas

Site: North Central Kansas Exp. Field

County: Republic
Town: Belleville
Soil: Crete silt loam

ESTABLISHMENT:

9/6/97 ; RCBD, 4 reps Plots 5'x15'; 3'x15' harvested

18 lb seed/acre

1998 FERTILIZATION:

September, 1997; 0-100-0

1998 PEST CONTROL:

Herbicide applied preplant to control grasses. Insecticide applied on May 1 for armyworm control.

1998 CONDITIONS:

April was wetter than normal, but May and the first part of June were very dry. A high temperature of 100 F was recorded in the first week of May. High temperatures in the first week of June were only in the 50's. Temperatures in July-September were very warm. Rainfall in July was ideal, but little rain fell in August and September.

TABLE 3. LABETTE CO. ALFALFA PERFORMANCE TEST RESULTS, 1995-1998.

							tons/a						
						Dry	/ Matte		4000	4005	05.00	Total, 15%	
			6-4	7-8	8-6	10-22	Total	1997 Total	1996 Total	1995 Total	95-98 Total	Moist.	
Released Cultivars	3												
Hobart Seed	SuperCuts	2.72	1.44	0.70	0.54	1.12	6.52	9.13	4.98	3.48	24.11	28.36	10
AgriPro	Depend+EV	2.68	1.49	0.68	0.44	1.19	6.48	9.30	4.79	3.23	23.80	28.00	10
Mycogen	TMF Generation	2.77	1.43	0.58	0.44	1.29	6.51	9.18	4.94	3.13	23.76	27.95	10
America's Alfalfa	Total+Z	2.79	1.41	0.58	0.37	1.26	6.41	8.84	5.28	3.22	23.75	27.94	10
America's Alfalfa	Affinity+Z	2.77	1.40	0.64	0.45	1.20	6.46	9.14	4.83	3.16	23.59	27.75	10
Dairyland Seeds	Magnum IV	2.57	1.48	0.74	0.54	1.31	6.64	8.93	5.19	2.74	23.50	27.65	10
DeKalb	DK 133	2.67	1.34	0.60	0.47	1.12	6.20	8.91	5.10	3.27	23.48	27.62	10
Great Plains	Haygrazer	2.83	1.30	0.57	0.42	1.31	6.43	9.22	4.82	2.86	23.33	27.45	10
W-L Research	WL 323	2.72	1.37	0.59	0.39	1.27	6.34	8.77	4.92	3.24	23.27	27.38	10
W-L Research	WL 252 HQ	2.65	1.29	0.67	0.50	1.30	6.41	8.87	4.82	3.14	23.24	27.34	10
America's Alfalfa	Innovator+Z	2.66	1.40	0.58	0.43	1.31	6.38	8.68	4.62	3.42	23.10	27.18	10
DeKalb	DK 127	2.71	1.34	0.59	0.33	1.26	6.23	8.96	4.53	3.09	22.81	26.84	98
NK	Rushmore	2.57	1.33	0.59	0.49	1.16	6.14	8.67	4.78	3.06	22.65	26.65	98
NE AES & USDA	Perry	2.84	1.30	0.47	0.25	1.32	6.18	8.58	4.75	2.75	22.26	26.19	96
KS AES & USDA	Riley	2.59	1.35	0.58	0.36	1.26	6.14	8.69	4.72	2.57	22.12	26.02	95
KS AES & USDA	Kanza	2.47	1.35	0.59	0.36	1.10	5.87	8.29	4.89	2.54	21.59	25.40	93
Experimental Strai	ns												
ABI	ABI 9141 Exp	2.68	1.45	0.72	0.46	1.35	6.66	9.22	5.01	3.29	24.18	28.45	10
Forage Genetics	3T26 Exp	2.65	1.45	0.62	0.40	1.09	6.21	8.78	4.61	3.22	22.82	26.85	98
Summary Statistic	s												
Average		2.69	1.38	0.61	0.42	1.23	6.33	8.90	4.88	3.07	23.18	27.27	10
LSD(0.05)		0.16	0.10	NS	0.10	0.11	0.32	0.41	NS	0.35	0.75	0.88	3
LSD(0.20)		0.12	0.08	0.09	0.08	0.09	0.21	0.27	0.28	0.23	0.49	0.58	2
CV(%)		4.93	5.97	16.31	19.12	7.64	3.53	3.25	6.21	9.20	2.29		
MCV(%)		5.83	7.09	19.48	22.90	9.09	5.02	4.60	NS	11.40	3.24	3.24	3
Site: Southeast County: Labette	t Ag. Research Cen	ter	1998 FI May; 2 1998 P I	1-54-1	82			1998 CONDITIONS: Growth after the August 6 cutting was suppressed by dry conditions that were not alleviated until the rains returned or					

Town: Mound Valley

Soil: Parsons silty clay loam

ESTABLISHMENT:

4/6/95; RCBD, 4 reps Plots 5'x30'; 3'x20' harvested

15 lb seed/acre

No pesticides needed; a transient swarm of blister beetles appeared just prior to the first cutting but not alleviated until the rains returned on September 13.

TABLE 4. RENO CO. ALFALFA PERFORMANCE TEST RESULTS, 1998.

					Forage Yield								
		DI.								97-98			
			nt Hei inche:	_				ry	Matter				Total,
BRAND	NAME		6-26		5-21	6-26	998 7-2	2	Total	1997 Total	97-98 Tatal	15% Moist	% of Mean
Released Cultivars		0 2 1	0 20		0 2.	0 20		_	Total	Total	Total	WIOISL	Wican
Mycogen	TMF Generation	24	12	18	2.58	1.15	1.1	Λ	4.83	5.13	9.96	11.72	105
Star	Spur	24	13	17	2.44	1.29	1.1		4.90	5.12	10.02	11.79	106
Germains	WL 324	24	13	18	2.40	1.24	1.3		4.95	5.04	9.99	11.75	105
Casterline	ProGro 424	24	12	19	2.27	1.26	1.3		4.83	4.98	9.81	11.54	103
Garst	645	24	12	16	2.45	1.22	1.1		4.81	4.97	9.78	11.51	103
Great Plains	Key	25	14	18	2.56	1.30	1.1		5.04	4.97	10.01	11.78	105
Star	Asset	25	13	17	2.54	1.24	1.3		5.10	4.87	9.97	11.73	105
Star	A-100	25	12	17	2.48	1.06	1.0		4.62	4.87	9.49	11.16	100
Germains	WL 325 HQ	23	14	19	2.22	1.32	1.3		4.86	4.83	9.69	11.40	102
Mycogen	TMF Multiplier II	25	14	19	2.46	1.21	1.2		4.92	4.81	9.73	11.45	103
America's Alfalfa	Affinity+Z	25	12	18	2.49	1.13	1.2		4.82	4.80	9.62	11.32	101
Dairyland Seeds	Magnum IV	22	12	17	2.56	1.11	1.1		4.79	4.80	9.59	11.28	101
Great Plains	Haygrazer	24	12	18	2.56	1.13	1.1		4.83	4.80	9.63	11.33	101
DeKalb	DK 127	24	13	17	2.51	1.12	1.2	21	4.84	4.78	9.62	11.32	101
W-L Research	WL 414	23	12	18	1.93	1.15	1.2		4.31	4.73	9.04	10.64	95
AgriPro	Depend+EV	22	11	18	2.39	1.18	1.2	20	4.77	4.72	9.49	11.16	100
America's Alfalfa	Archer	24	13	18	2.23	1.25	1.2		4.72	4.69	9.41	11.07	99
KS AES & USDA	Riley	25	12	17	2.40	1.07	1.0)4	4.51	4.68	9.19	10.81	97
Star	Excalibur II	24	13	19	2.29	1.20	1.2	25	4.74	4.67	9.41	11.07	99
W-L Research	WL 252 HQ	23	12	19	2.43	1.04	1.1	5	4.62	4.65	9.27	10.91	98
Sharp	AlfaLeaf II	25	11	18	2.40	1.16	1.0	8	4.64	4.60	9.24	10.87	97
NE AES & USDA	Perry	24	12	16	2.42	1.04	1.1	2	4.58	4.58	9.16	10.78	97
Sharp	Shamrock	25	13	18	2.40	1.06	1.2	23	4.69	4.57	9.26	10.89	98
Star	Stamina	25	12	16	2.35	1.14	1.1	7	4.66	4.48	9.14	10.75	96
KS AES & USDA	Kanza	26	13	18	2.06	1.17	1.1	1	4.34	4.35	8.69	10.22	92
W-L Research	Ace	22	12	18	2.03	1.24	1.3	37	4.64	4.26	8.90	10.47	94
Experimental Stra	ins												
Cal/West	CW 5440 Exp	23	13	19	2.16	1.35	1.3	88	4.89	4.71	9.60	11.29	101
Cal/West	CW 5406 Exp	23	12	18	2.25	1.21	1.3		4.78	4.63	9.41	11.07	99
Cal/West	CW 4429 Exp	23	14	18	2.25	1.21	1.2	28	4.74	4.62	9.36	11.01	99
Summary Statistic	s												
Average		24	12	18	2.36	1.18	1.2	21	4.75	4.74	9.49	11.16	100
LSD(0.05)		NS	NS	NS	0.22	0.12	0.1		0.34	0.35	0.49	0.58	5
LSD(0.20)		2	NS	NS	0.17	0.09	0.1		0.22	0.23	0.32	0.38	3
CV(%)		7	14	9	7.89	8.38	9.3		5.03	5.25	3.64		
MCV(%)		NS	NS	NS	9.28	9.87	10.9	94	7.16	7.38	5.16	5.16	5
LOCATION: South Site: South Ce County: Reno Town: Hutchinso Soil: Ost silt lo ESTABLISHMENT	ntral Experiment Field on am	Nor 1998 Her to c	ne; soi B PES [*] bicide ontrol	T CON applied grasse	IZATION: Issted high in P and K ISONTROL: Isplied on January 27 Isasses. Insecticide oril to control alfalfa						erely ting		
9/1/96 ; RCBD, 4 Plots 5'x20, 3x20' 18 lb seed/acre	4 reps		evils.	•									

TABLE 5. FINNEY CO. IRRIGATED ALFALFA PERFORMANCE TEST RESULTS, 1998.

						orage Yie	eld		,	
					tons	s/acre				97 -98
				I	Dry Matt	er			Total,	Total,
				1998			1997	97-98	15%	% of
BRAND	NAME	6-5	7-14	8-19	9-25	Total	Total	Total	Moist.	Mean
Released Cultivars	3									
W-L Research	WL 414	3.72	2.15	2.74	2.37	10.98	9.62	20.60	24.24	103
Germains	WL 324	4.04	2.19	2.52	2.29	11.02	9.46	20.48	24.09	103
Mycogen	TMF Multiplier II	4.03	2.18	2.54	2.31	11.05	9.29	20.34	23.93	102
Germains	WL 325 HQ	4.13	2.18	2.57	2.26	11.12	9.21	20.33	23.92	102
Star	Stamina	3.94	2.22	2.52	2.32	10.99	9.34	20.33	23.92	102
Casterline	ProGro 424	3.99	2.29	2.55	2.39	11.20	9.08	20.28	23.86	102
Golden Harvest	GH-755	3.99	2.25	2.53	2.31	11.07	8.99	20.06	23.60	101
DSS	Enhancer	3.82	2.26	2.55	2.40	11.02	9.02	20.04	23.58	101
KS AES & USDA	Riley	3.75	2.22	2.57	2.30	10.83	9.21	20.04	23.58	101
Garst	630	3.87	2.32	2.69	2.28	11.16	8.81	19.97	23.49	100
Cargill	Big Horn	3.94	2.14	2.48	2.24	10.79	9.15	19.94	23.46	100
Star	Spur	3.99	2.11	2.46	2.24	10.78	9.14	19.92	23.44	100
Sharp	AlfaLeaf II	3.97	2.08	2.37	2.23	10.64	9.22	19.86	23.36	100
W-L Research	WL 323	3.93	2.11	2.42	2.26	10.72	9.14	19.86	23.36	100
DeKalb	DK 127	3.82	2.17	2.54	2.21	10.73	9.12	19.85	23.35	100
Garst	645	3.73	2.19	2.45	2.17	10.54	9.22	19.76	23.25	99
Star	A-100	3.90	2.16	2.34	2.19	10.58	9.17	19.75	23.24	99
Golden Harvest	GH-766	3.96	2.18	2.46	2.22	10.82	8.78	19.60	23.06	98
W-L Research	Ace	3.73	2.23	2.49	2.20	10.64	8.93	19.57	23.02	98
Dairyland Seeds	Magnum III	3.84	2.13	2.51	2.30	10.78	8.71	19.49	22.93	98
NE AES & USDA	Perry	3.99	2.09	2.43	2.19	10.76	8.80	19.49	22.93	98
Star	Asset	3.80	2.13	2.46	2.22	10.59	8.69	19.28	22.68	97
Sharp	Sure	3.80	2.12	2.42	2.22	10.54	8.68	19.22	22.61	96
KS AES & USDA	Kanza	3.87	2.27	2.53	2.31	10.96	8.13	19.09	22.46	96
Sharp	Shamrock	3.81	2.00	2.12	2.11	10.90	9.02	19.05	22.40	96
Star	Excalibur II	3.66	2.07	2.39	2.24	10.03	8.59	18.95	22.41	95
DeKalb	DK 133	3.75	1.93	2.38	2.24	10.36	8.68	18.93	22.29	95 95
Experimental Strai		3.13	1.55	2.50	۷.۷۱	10.20	0.00	10.30	44.41	30
DSS	DSS 5211X Exp	4.35	2.35	2.72	2.41	11.81	9.71	21.52	25.32	108
Cal/West	CW 5406 Exp	4.35 3.95	2.33	2.72	2.41	11.01	9.71	21.00	23.32	105
	•		2.21							
Cal/West	CW 5440 Exp	3.86	2.32 2.22	2.59 2.61	2.37 2.42	11.12	9.56 9.55	20.68 20.65	24.33 24.29	104 104
Cal/West Cal/West	CW 4420 Exp	3.86				11.10				
	CW 4429 Exp	3.71	2.19	2.46	2.27	10.62	9.37	19.99	23.52	100
DSS Summaru Statiatia	DSS 5106X Exp	3.80	2.15	2.50	2.29	10.73	9.01	19.74	23.22	99
Summary Statistic	S	2.00	2 4 7	2.50	2.00	40.00	2.40	40.00	22.45	400
Average		3.89	2.17	2.50	2.28	10.83	9.10	19.93	23.45	100
LSD(0.05)		NS	0.12	0.16	0.11	0.50	0.27	0.49 0.32	0.58	2
LSD(0.20)		NS	0.10	0.13	0.08	0.39	0.17	0.38	2	
CV(%)		6.44	4.85	5.57	3.95	3.90	2.09 2.95	1.73	2	
MCV(%)		7.57	5.69	6.54	4.63	4.58	2.46	2		
LOCATION: South	west Kansas	1998 FEF	RTILIZAT	ION:		1998 CONDITIONS:				
Site: Southwes	st ResExt. Center	None ne	eded		Extremely hot, dry weather in .				er in Jun	e
County: Finney						appeared to lower 2nd-cutting yie				lds
		1998 PFS	TIMOD TO	$D \cap I$.						

Town: Garden City Soil: Keith silt loam

ESTABLISHMENT:

8/29/96; RCBD, 4 reps Plots 3'x20'; 3'x20' harvested

32 lb seed/acre

1998 PEST CONTROL:

Insecticide applied on April 11 to control aphids.

from typical levels. Flood irrigations were applied five times during the season. Very wet conditions in July reduced the need for irrigations during that part of the summer.

ABI ABI Alfalfa	515-	292-2432	C	Cal/West		608-786-1554
2316 259th St.				R.R. 1, Bo		
Ames, IA 5001				west Sale	em, WI 54669	C 7 0 0 40 44 40 40
ADI 0444 E	1 2 3 4 5 6 7			4044.5		6 7 8 9 10 11 12 13
ABI 9141 Exp	4 H H H H H -			1344 Exp		
ABI 9142	4 H R H H H -			1346 Exp		
•	3			1469 Exp		
ZC9641 Exp				CW 4429 E	-AP	
ZN9540 Exp				CW 4598 E	•	
ZN9541 Exp				CW 5406 E	•	
ZN9646 Exp				CW 5440 E	Exp	
A ari Dro	800-	334-4730		OK49		
AgriPro	800-	334-4730	(Cargill	•	612-742-6743
PO Box 2962			`	_	brid Seeds	3.22 33
	1/0 00004 400	.0		,		
Snawnee Missi	on, KS 66201-130			P.O. Box		
	1 2 3 4 5 6 7			iviinneapo	lis, MN 55440	0 7 0 0 40 44 40 40
•	4 H H H H H -					6 7 8 9 10 11 12 13
Dominator	4 HRHHH -	R - M R		Big Horn		HRRHRH
America's Alfal	fa 913-	384-4940		Crown II	знкнн	H M R
America's Alfal			(Casterline	8	800-444-4137
P.O. Box 2955				Casterline	Seeds, Inc.	
6700 Antioch				Box 1377		
Shawnee Missi	on KS 66201			1st & Map		
Shawhee Missi	1 2 3 4 5 6 7	7 9 0 10 11 12 12			ty, KS 67801	
Affinitus 7	4 H H H H H -			Douge Cit	•	6 7 9 0 10 11 12 12
Affinity+Z	4 H R H H H N			ProGro 424	· · · · · · · · · · · · · · · · · · ·	6 7 8 9 10 11 12 13 H R R M - M
Aggressor				F10G10 424	+ 4 N N N N	TKKW-W
	4 H R H H R -					
Archer	5 M M H R R F					
Innovator+Z	3 H H H H N					
Total+Z	3 H H H H N	IKSWK				
Variety characteriz	zation codes:	Fall dormancy i	ratings:		Pest resistance	ratings:
1 = Fall dormancy	rating	Check variety	Rating	<u>Code</u>	Resistance class	% Resistant plants
2 = Bacterial wilt		Norseman	1	S	Susceptible	0-5%
3 = Verticillium wil	t	Vernal	2	L	Low resistance	6-14%
4 = Fusarium wilt		Ranger	3	M	Moderate resistance	15-30%
5 = Anthracnose ra		Saranac	4	R	Resistance	31-50%
6 = Phytophthora		DuPuits	5	Н	High resistance	>50%
7 = Spotted alfalfa	aphid	Lahontan	6	-	Not adequately tested	d
8 = Pea aphid		Mesilla	7			

4 = Fusarium wilt
Ranger
3 M Moderate resistance
15-30%
5 = Anthracnose race 1 Saranac
4 R Resistance
31-50%
6 = Phytophthora root rot
DuPuits
5 H High resistance
>50%
7 = Spotted alfalfa aphid
Lahontan
6 - Not adequately tested
8 = Pea aphid
9 = Blue alfalfa aphid
Moapa 69
10 = Stem nematode
11 = Aphanomyces root rot race 1
12 = Southern root knot nematode
13 = Northern root knot nematode
13 = Northern root knot nematode

(continued)

DSS	31	6-275-2359		Garst	608-249-8977					
Drussel Seed		2.0 2000	`	Garst See						
2197 W. Para				P.O. Box						
Garden City, h		7 0 0 40 44 40 40		Madison,	WI 53707-7790					
D00 5400V 5		7 8 9 10 11 12 13		000	1 2 3 4 5 6 7 8 9 10 11 12 1					
DSS 5106X Exp	-			630	4 H M R M R M R M M					
DSS 5211X Exp				645	3 H R R H H M R - M M -					
Enhancer		R M		Cormoino	785-674-2062					
Reward	4 H R H R H	RHMMM	,	Germains						
Deimdend Coe	۵۰ ۹۰	0-236-0163			s Seed Co.					
Dairyland See		0-230-0103		P.O. Box	373					
Dairyland See	d Co.			Hill City, I	KS 67642					
P.O. Box 958					1 2 3 4 5 6 7 8 9 10 11 12 1					
West Bend, W	/I 53095			WL 324	3 H R H H H R H - M H -					
	1 2 3 4 5 6	7 8 9 10 11 12 13		WL 325 H	Q 3 H R H H H R R M R R -					
Magnum III	4 R M R M R	MRMML								
Magnum IV	4 H R H R H	M - M R M - M	(Golden Harvest 800-228-990						
				J.C. Robi	nson Seed Co.					
DeKalb	81	5-758-9323		100 J.C. I	Robinson Blvd.					
DeKalb Plant	Genetics Corp.			P.O. Box						
3100 Sycamo	re Rd.				, NE 68069					
DeKalb, IL 60				vvatorioo,	1 2 3 4 5 6 7 8 9 10 11 12 1					
Dortain, 12 oc		7 8 9 10 11 12 13		GH-755	4 H R H H H R R R R R -					
DK 127		H H - R H - R		GH-766	3 H R H H H R R - R R -					
DK 133		RR-MR		GI 1-700	3 II K II II II K K - K K -					
DK 133		H H - R H		Great Plai	ns 919-362-1583					
DR 142	4 11 10 11 10 11	11 11 - 13 11	`		ins Research Co.,Inc.					
Forage Genet	ics 60	8-786-2121			•					
Forage Genet					aire Farm Rd.					
-				Apex, NC						
	Gills Coulee Rd.				1 2 3 4 5 6 7 8 9 10 11 12 1					
West Salem, V				Haygrazer						
	1 2 3 4 5 6	7 8 9 10 11 12 13		Key	4					
3T26 Exp										
Variety character		Fall dormancy r			Pest resistance ratings:					
1 = Fall dormanc	y ratıng	Check variety	Rating		·					
2 = Bacterial wilt	.:112	Norseman	1	S	Susceptible 0-5%					
3 = Verticillium w		Vernal	2	L	Low resistance 6-14%					
4 = Fusarium wilt 5 = Anthracnose		Ranger	3	M	Moderate resistance 15-30%					
		Saranac DuPuits	4 5	R H	Resistance 31-50% High resistance >50%					
6 = Phytophthora		Lahontan	5 6	н -	S .					
7 = Spotted alfalf 8 = Pea aphid	α αμιτιυ	Mesilla	7	-	Not adequately tested					
9 = Blue alfalfa a	phid	Moapa 69	Fall dormancy and disease and insect resistance							
10 = Stem nemat		CUF 101	9	ratings are	e from Alfalfa Varieties, a publication of the					
10 – Otom nomal		001 101	Certified Alfalfa Seed Council, or from developers							

11 = Aphanomyces root rot race 1

12 = Southern root knot nematode

13 = Northern root knot nematode

Certified Alfalfa Seed Council, or from developers of

the varieties. Blank spaces indicate that the variety

has not been adequately tested.

Hobart Seed	800-866-6074	NE AES & USI	DΑ					402	2-4	72	-42	290		
Hobart Seed		Foundation Se	eed	Di	visi	on								
530 S. Main		University of N	lebr	ras	ka-	Lin	СО	ln						
Hobart, OK 7	73651	3115 North 70)th											
	1 2 3 4 5 6 7 8 9 10 11 12 13	Lincoln, NE 6	850	7-2	210	4								
SuperCuts	4 H H H H H - R - L R		1	2	3	4	5	6	7	8	9	10 ·	11 1	12 13
	705 500 0445	Perry	3	R	-	-	L	-	М	R	-	-	-	
KS AES & US								~ 4	~ -					
KSU - Found	ation Seed	NK						310	o-5	43	-21	' 07		
2200 Kimball	Ave.	Novartis Seed	ls, Ir	nc.										
Manhattan, K	S 66502	1060 Wheatla	nd [Эr.										
	1 2 3 4 5 6 7 8 9 10 11 12 13	Buhler, KS 67	7522	2										
Kanza														2 13
Riley	4 H L - M - H H	Ciba 2444												
MDC	515-733-5274	Fortress												
MBS	515-755-5274	Rushmore	4	. H	l R	Н	Н	Н	Н	R	-	-	Η	
MBS, Inc.	_	Pioneer						51	5-2	70.	-33	342		
225 West 1st	· • · · ·		المم	الدما	l.a	_		J 1 .	J- <u>Z</u>	.70	-00) + ∠		
Story City, IA		Pioneer Hi-Bre	ea II	nu.	, in	C.								
	1 2 3 4 5 6 7 8 9 10 11 12 13	Box 287												
PGI3212 Exp		7305 NW 62n	-											
PGI3392 Exp		Johnston, IA			_		_	_	_	_				
Mycogen	800-692-6436		1	2	3	4	5	6	7	8	9	10 ′ M	11 1	12 13
Mycogen See	***************************************	5454	4	· K				н		K	S	IVI	L	
		88C2PI2 Exp 90W3PR1 Exp	-	-						-	-	-	-	
P.O. Box 214	ate Center Curve	91CO1PR1 Exp										-		
		91CO2PR1 Exp			_					-	_	_	_	
St. Paul, MN		91I12PJ1 Exp	, - -	_	_	_	_	_	_	_	_	_	_	
TME Congretio	1 2 3 4 5 6 7 8 9 10 11 12 13 on 4 H H H H H - R R	011121 01 Exp												
	4													
i wir wuuupilet														

Variety characterization codes:	Fall dormand	cy ratings:	Pest resistance ratings:						
1 = Fall dormancy rating	Check variety	Rating	<u>Code</u>	Resistance class	% Resistant plants				
2 = Bacterial wilt	Norseman	1	S	Susceptible	0-5%				
3 = Verticillium wilt	Vernal	2	L	Low resistance	6-14%				
4 = Fusarium wilt	Ranger	3	M	Moderate resistance	15-30%				
5 = Anthracnose race 1	Saranac	4	R	Resistance	31-50%				
6 = Phytophthora root rot	DuPuits	5	Н	High resistance	>50%				
7 = Spotted alfalfa aphid	Lahontan	6	-	Not adequately tested					
8 = Pea aphid	Mesilla	7							
9 = Blue alfalfa aphid	Moapa 69	8		ancy and disease and in					
10 = Stem nematode	CUF 101	9	-	e from Alfalfa Varieties,					
11 = Aphanomyces root rot race 1				Alfalfa Seed Council, or es. Blank spaces indic					
12 = Southern root knot nematode				en adequately tested.	ato that the variety				
13 = Northern root knot nematode									
		/ t! I\							

(continued)

Sharp Bros. Seed Company													
Box 140													
Healy, KS 678	50												
,	1	2	3	4	5	6	7	8	9	10	11	12	13
AlfaLeaf II	4	R		Н	Н			Н	-	R		_	-
Shamrock	-	-	-	-	-	-	-	-	-	-	-	-	-
Sure	-	-	-	-	-	-	-	-	-	-	-	-	-
Star						78	5-3	46	-54	447	7		
Star Seed													
101 Industrial A	١ve												
Osborne, KS 6	747	73											
	1	2	3	4	5	6	7	8	9	10	11	12	13
A-100	-	-	-	-	-	-	-	-	-	-	-	-	_
Asset	4	Н	R	R	R	Н	R	R	-	-	М	-	-
Excalibur II	-	-	-	-	-	-	-	-	-	-	-	-	-
Spur	4	Н	R	Н	Н	Н	R	Н	-	М	R	-	М
Stamina	4	Н	R	Н	Н	Н	Н	Н	-	Н	R	-	Н
W-L Research						60	8-8	82	-4	100)		
W-L Research,	Inc).											
8701 Hwy. 14													
Evansville, WI	53	536	3-8	75	2								
,	1				5	6	7	8	9	10	11	12	13
Ace	4	Н	R	Н	Н	Н	М	R	R	Н	R	-	-
WL 252 HQ	2	Н	R	Н	Н	Н	М	R	L	R	L	-	-
WL 322 HQ	4	Н	R	Н	М	R	Н	Н	R	L	-	-	L
WL 323	4	Н	R	Н	Н	Н	М	R	-	Н	R	-	-
WL 414	6	R	R	Н	R	Н	Н	Н	Н	R	-	R	-

Variety characterization codes:	Fall dormancy	ratings:	Pest resistance ratings:						
1 = Fall dormancy rating	Check variety	<u>Rating</u>	<u>Code</u>	Resistance class	% Resistant plants				
2 = Bacterial wilt	Norseman	1	S	Susceptible	0-5%				
3 = Verticillium wilt	Vernal	2	L	Low resistance	6-14%				
4 = Fusarium wilt	Ranger	3	M	Moderate resistance	15-30%				
5 = Anthracnose race 1	Saranac	4	R	Resistance	31-50%				
6 = Phytophthora root rot	DuPuits	5	Н	High resistance	>50%				
7 = Spotted alfalfa aphid	Lahontan	6	-	Not adequately tested					
8 = Pea aphid	Mesilla	7							
9 = Blue alfalfa aphid	Moapa 69	8		ancy and disease and in					
10 = Stem nematode	CUF 101	9	•	e from Alfalfa Varieties,	•				
11 = Aphanomyces root rot race 1				Alfalfa Seed Council, or es. Blank spaces indic	•				
12 = Southern root knot nematode				es. Blank spaces indicate an adequately tested.	ale that the variety				
13 = Northern root knot nematode			nas not be	on adoquatory toolog.					

ELECTRONIC ACCESS

For those interested in accessing crop performance testing information electronically, try visiting our World Wide Web site. Most of the information contained in this publication is available for viewing or downloading. The URL is http://www.ksu.edu/kscpt.

Excerpts from the

UNIVERSITY RESEARCH POLICY AGREEMENT WITH COOPERATING SEED COMPANIES*

Permission is hereby given to Kansas State University to test our varieties and/or hybrids designated on the attached entry forms in the manner indicated on the test announcement. I understand that all results from Kansas crop performance tests belong to the University and to the public and shall be controlled by the University so as to produce the greatest benefit to the public. It is further agreed that the name of the University shall not be used by the company in any commercial advertising either in regard to this agreement or any other related matter.

These materials may be freely reproduced for educational purposes. All other rights reserved. In each case, give credit to the author(s), name of work, Kansas State University, and the date the work was published.

CONTRIBUTORS

MAIN STATION, MANHATTAN

Kraig Roozeboom, Associate Agronomist (Senior Author)

RESEARCH CENTERS

Joseph Moyer, Parsons Merle Witt, Garden City

EXPERIMENT FIELDS

William Heer, Hutchinson W. Barney Gordon, Belleville

NOTE: Trade names are used to identify products. No endorsement is intended, nor is any criticism implied of similar products not named.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service, Manhattan 66506
SRP 828
January 1999