## THE EFFECTS OF DIET THERAPY, BEHAVIOR MODIFICATION, AND EXERCISE ON WEIGHT REDUCTION AND SERUM LIPIDS

by

Joyce Petitjean Oldenburg

B. S., Kansas State University, 1977 M. S., Kansas State University, 1979

A MASTER'S THESIS

submitted in partial fulfilment of the

requirements for the degree

MASTER OF SCIENCE

Department of Foods and Nutrition

KANSAS STATE UNIVERSITY Manhattan, Kansas

1979

Approved by:

Major Professor

Sprc.Coll. LD 2668 .T4 1979 042 C.2

#### ACKNOWLEDGEMENTS

I would like to express my sincere appreciation of and gratitude to members of the research team: Barbara Moyer -- my co-partner and friend; Dr. William Zuti--who spent many long hours in the Exercise Physiology Laboratory testing research subjects; and Dr. Robert Reeves--who worked very hard in the co-ordination and planning of the serum lipid research. I would also like to thank Susan Davis and Jean Craig for their assistance in the laboratory. A special thank you is given to Dr. Jane Bowers for her support and encouragement duing my undergraduate and graduate work at Kansas State. I also would like to thank Dr. Art Dayton for his assistance in the statistical analysis of the data. I would like to acknowledge Peggy Weddle, who helped me type this thesis. I also would like to thank Eunice Bassler and other members of Manhattan Weslevan Church for their love and prayers. A very special thank you is given to Dr. Robert Reeves for his support and assistance in the preparation of this manuscript, and for his support and encouragement throughout my two years of graduate work at K-State. I would like to express personal appreciation for each member of my graduate committee: Dr. Jane Bowers, Dr. William Zuti, and Dr. Robert Reeves. I would like to express my sincere love and appreciation for my husband, Charlie, who has comforted, assisted, and encouraged me during this time. I also would like to thank my parents for their unfailing example, love and support. Above all, I would like to thank my Heavenly Father, Who continues to be my Source, my Guide, and my Goal.

## TABLE OF CONTENTS

INTRODUCTION
REVIEW OF LITERATURE
Causes of Obesity Diet Therapy as a Means of Weight Reduction Behavior Modification as a Means of Weight Reduction Physical Activity and Weight Reduction Weight Reduction and Body Composition Lean Body Weight and Urinary Creatinine Excretion Coronary Heart Disease Risk Factors Serum Lipids Physical Activity and Serum Lipids Body Weight and Serum Lipids Blood Pressure and Physical Activity
Blood Pressure and Body Weight Dietary Intake and Serum Lipids
METHODS AND PROCEDURES
Subjects Preliminary Testing Period Post Testing Period Experimental Testing Period Post Experimental Follow-up Statistical Treatment of the Data
RESULTS AND DISCUSSION
. Body Composition Urinary Creatinine Excretion Serum Lipids Blood Pressure Blood Serum Analysis Dietary Intake
SUMMARY AND CONCLUSIONS
BIBLIOGRAPHY
APPENDIX

### INTRODUCTION

Obesity occurs when an excess amount of adipose tissue accumulates on the body mass. This accumulation is a result of an energy imbalance: energy consumed exceeds energy expended. Overeating, a metabolic defect, a lack of physical activity, or a combination of these factors can induce a positive energy balance.

The number of obese individuals in the United States has increased to the point that obesity can be considered a national problem. The United States Public Health Service reports that 25 to 45 percent of the adult American population over 30 years of age is more than 20 percent overweight (1). Data obtained by life insurance companies have shown that life expectancy is increased in individuals who are lighter than average weight and decreased in individuals who are heavier than average weight. This suggests that any excess weight is associated with a detrimental impact on longevity (2).<sup>1</sup>

Statistical evidence has indicated that obese individuals also have a greater propensity for developing diseases such as diabetes mellitus, coronary heart disease, gall bladder disease, and hypertension (3). Because of these relationships and the tendency toward decreased life expectancy, the Select Committee on Nutrition and Human Needs has recommended curtailing energy intake and increasing physical activity for obese individuals (4).

<sup>&</sup>lt;sup>1</sup>The terms obesity and overweight are often used synonomously even though they are not really interchangable. By definition, an obese individual has accumulated an excess amount of adipose tissue on the body mass. An overweight individual is "over" or above average weight. An athlete, who has increased his muscle mass, may be heavier than normal for his height and be considered overweight, even though he may be carrying very little adipose tissue on the body mass.

Until recently, many professionals have taken a pessimistic view of the obese individual's chances of controlling his weight. Results from traditional, dietary approaches to obesity have been discouraging. Stunkard (5) has reported that of 100 obese patients admitted to a nutrition clinic, only 12 percent lost as much as 20 pounds and 28 percent never returned to the clinic. Two years after the end of treatment, only two of the initial 100 patients referred to the clinic had maintained their weight loss.

Stuart (6) was one of the first to successfully use behavioral techniques to help obese individuals change inappropriate eating behaviors. He reported that over a period of one year, 30 percent of the original sample had lost more than 40 pounds and 60 percent lost more than 30 pounds. This approach to obesity is based on the premise that obesity is an inappropriately learned behavior that responds to many environmental cues unrelated to physiological hunger. By using stimulus control techniques which change the antecedents to inappropriate food-related behaviors, an individual can change his eating behavior and the consequences of that behavior.

The importance of body composition to the study of nutrition, exercise performance, and responses to other physical stresses has been recognized. Most of the accepted procedures (fluid volume analysis, whole body isotopic counting, and densiometry) are either time consuming, or they require expensive equipment. Because of the large amount of creatine in muscle tissue and the direct proportionality of body creatine to urinary creatinine excretion, urinary creatinine excretion values have been used to estimate body compostion. Although urinary creatinine excretion values have been correlated with lean body mass, the accuracy and consistency with which this method can estimate body composition remains in question.

Risk factors for identifying individuals with a greater propensity for development of coronary heart disease include: depressed high density lipoproteins (7), elevated serum cholesterol levels, hypertension, smoking, physical inactivity, and obesity (8). The risk of developing coronary heart disease is not only proportional to the antecedent level of risk factors, but also the risk increases with the number of risk factors involved (9). The full implications of this statement become disheartening when considering the concomitant factors often associated with obesity: hypertension, high serum cholesterol levels, depressed high density lipoprotein levels, and physical inactivity.

Because of the relationship of obesity to reduced life expectancy, coronary heart disease, and the potential for infringement on personal happiness, a need exists to investigate the effectiveness of behavioral and dietary approaches to weight reduction and the effects of weight reduction and physical acitivity on body composition, blood pressure, and serum lipids.

The purpose of the present study, conducted jointly by the Department of Foods and Nutrition and the Department of Health, Physical Education and Recreation, was to investigate the effects of four methods of weight reduction on body composition, blood pressure, and serum lipids. The four methods of weight reduction were: 1) diet therapy, 2) diet therapy plus exercise, 3) behavior modification, and 4) behavior modification plus exercise.

Subjects in the diet therapy groups were given guidance in designing their own dietary regimens based on food choices from the "American Dietetic

Association Exchange Lists for Meal Planning" (10). In addition to dietary guidance, subjects in the diet therapy plus exercise group were given assistance in planning supplemental programs of physical activity (equivalent of 250 kcal energy expenditure, daily). Subjects in the behavior modification groups were to lose weight and maintain that loss by developing appropriate eating patterns rather than by following a special, energy-restricted diet. The behavior modification techniques employed were taken from: <u>Habits Not</u> <u>Diets. The Real Way to Weight Control</u> (11). Treatment of subjects in the behavior modification plus exercise group was identical to that of subjects in the behavior modification group except supplemental programs of physical activity (equivalent of 250 kcal energy expenditure, daily) also were included.

#### REVIEW OF LITERATURE

Obesity is defined as an excess accumulation of adipose tissue on the body mass. Brozeck et al. (12) have suggested that body fat compose 16 to 19 percent of the total body mass for a normal man and 19 to 23 percent for a normal woman.

### Causes of Obesity

Excess body fat accumulates when a positive energy balance occurs: energy consumed exceeds energy expended. This is usually a result of 1) simple overeating, 2) a metabolic defect, 3) a lack of physical activity, or 4) a combination of those factors.

In a study conducted by Miller and Parsonage (13), 29 women who claimed that they could not lose weight were isolated in a country house and fed 1,500 kcal per day for three weeks. Nine women maintained within  $\pm$  1 kilogram and were characterized by low basal metabolic rates and long previous histories of dieting. The remaining women did lose weight in varying amounts. These findings suggest that among groups of potential dieters, there will be some who have become metabolically adapted to a low energy diet, and others whose inability to lose weight is illusory.

Many controlled studies show significantly reduced physical activity in the obese. Stefanik et al. (14) assessed 14 obese boys and 14 controls at school and during summer camp. He found that the obese boys preferred the less active exercises. A similar study was conducted by Bullen et al. (15) with girls. Motion pictures were taken of obese teenage girls participating in swimming, volleyball and tennis. The pictures showed that the obese girls were less active than non-obese girls even though all the girls spent the same amount of time at the sport. Mayer (16) has reviewed a number of studies which have shown that obesity is more closely related to inactivity than to energy intake.

### Diet Therapy as a Means of Weight Reduction

The traditional medical approach to obesity has been diagnosis and referral to a dietician or nutritionist who can provide dietary guidance. The client usually receives excellent advice about the nutritive value of various foods and the proper amounts to include in his prescribed diet. Unfortunately, the client often becomes frustrated with the dietary restrictions imposed upon him and ignores everything he has been taught.

Stunkard and McLaren-Hume (17) and Shipman and Plesset (18) have reported a dropout rate of 20-80 percent for those entering an educational dietary counseling program. Of those individuals remaining in treatment, only 25 percent were able to lose as much as 20 pounds and only 5 percent lost 40 pounds or more. Stunkard (5) also has reported that of 100 consecutive obese out-patients admitted to a nutrition clinic, only 12 percent lost as much as 20 pounds and 28 percent never returned to either the nutrition clinic or the initial referring clinic. Two years after the end of treatment, only two of the initial 100 patients referred to the clinic had maintained their weight loss.

## Behavior Modification as a Means of Weight Reduction

Studies conducted by Stuart and Davis (19), Stunkard (5), and Levitz (20) support the hypothesis that behavior modification is the most effective way to lose weight permanently. Behavior modification helps the obese person change the habits and behavior patterns associated with the act of obtaining and consuming food. One can think of all behaviors as determined by its antecedents: those events which take place prior to the occurrence and consequences of a behavior (Figure 1).



FIG. 1. Schematic representation of the determinants of simple behavior (21).

If the consequences of a behavior are positive (covert: \$5 reward or overt: self-esteem), the probability that the behavior will be repeated increases.

Ferster (22) applied these behavioral concepts to the problem of obesity. He reasoned that if the obese had a <u>learned</u> overweight-eating style that was responsible for their excess weight, then by using stimulus control techniques, which change the antecedents to such behaviors, the therapist should be able to help the client change his eating style and, therefore, reduce weight. For example, if a client experiences a desire to eat a doughnut, in response to an environmental cue or stimulus (walking by the bakery on the way home from work), the first step is to change the environment (taking a different route home), thus decreasing his exposure to a cue that usually provokes an eating response. This approach assumes that all behaviors are learned and maintained through interactions between the individual and relevant persons and situations in his environment.

Stuart (6) was the first to successfully apply the behavioral technology developed experimentally to control various aspects of eating habits. In 1967, he reported the results of his treatment of 10 women. Over a period of one year, two of the ten women dropped out of treatment, 30 percent of the original sample lost more than 40 pounds and 60 percent lost more than 30 pounds.

The attrition rate is usually lower in behavioral programs than in traditional weight loss programs. Brightwell (23) has suggested that motivation is easier developed in a behavior modification program because the program is in no way restrictive; there is really nothing clients are not allowed to do, only things they must do in addition to their normal daily behavior (eat slowly, keep food out of sight, plan ahead, etc.). Most find this easier to tolerate than rules and regulations which directly restrict dietary intake.

The behavior modification approach to weight control focuses on changing people's eating habits and does not include specific dietary guidance. Ritt et al. (24) conducted a study to learn whether behavior modification techniques result in the adoption of nutritionally sound dietary patterns. Originally, the diets of 15 individuals met two-thirds of all Recommended Dietary Allowances (RDA). After a 20-week behavior modification program, the only nutrients below this level were iron, thiamine, and calcium. Other changes involved an increase in nutrient density (amount of a nutrient found in a food/the number of kcal in the same food x 1,000) for protein, fiber, phosphorous, iron, vitamin A, thiamine, riboflavin, niacin, ascorbic acid and cholesterol. Nutrient density for carbohydrates, fat, and calcium decreased.

## Physical Activity and Weight Reduction

Traditionally physical activity has been ignored in most weight control programs because of two common misconceptions: 1) exercise plays an insignificant role in total body energy requirements and 2) an increase in physical activity will result in an increased appetite (19).

Physical activity, in fact, is the most variable factor determining the total energy requirement. Inactive individuals may require an energy increase of only 40-50% above their basal needs to satisfy the total energy requirement, whereas very active persons may need increases of 100% or more above basal needs (Figure 2) (19).

Total Energy Needs for the Sedentary:

basal

physical activity

Total Energy Needs for the Active:

basal

physical activity

FIG. 2. Diagrammatic representation illustrating the effects of physical activity on total energy needs (19).

Mayer (16) has shown that subjects at a truly sedentary level of activity eat more than those who participate in regular physical activity. Research with both humans and animals suggests that the physiological mechanisms which act to balance food intake with energy expenditure work only within normal ranges of physical activity (16).

## Weight Reduction and Body Composition

Changes in body weight and body composition of overweight individuals as a result of increased physical activity and decreased energy intake seem to be dependent on the nature and magnitude of the energy deficit (25). In a study conducted by Zuti and Golding (25), exercise, diet and exercise plus diet weight reduction programs were used to create an energy deficit. Methods using exercise produced a greater increase in lean body weight, with a greater percentage of fat loss. Treatment by dietary control resulted in a loss of lean body mass. Subjects in the exercise-only group had a slightly greater increase in lean body weight than subjects in the exercise plus diet group.

Keys (26) has found that 98 percent of the weight loss is usually adipose tissue when a diet plus exercise program is employed while only 75 percent of the weight loss is adipose tissue when a diet-only program is followed.

## Lean Body Weight and Urinary Creatinine Excretion

The precursor of creatinine is creatine which is formed in the liver from three amino acids: arginine, glycine, and methionine. After being discharged in the blood, creatine is absorbed by the body in varying amounts (27). Approximately 98 percent is absorbed by the muscle tissues in the form of creatine-phosphate (28). Creatinine is formed from the hydrolysis of dephosphorylated creatine-phosphate and is excreted in the urine. The fact that creatine is found primarily in muscle tissue, provides the basis for describing urinary creatinine excretion (UCE) as a function of lean body weight.

Forbes and Bruining (29) have reported a very high correlation between LBW as determined by potassium-40 counting and UCE (r=.988). The effect of technical error was reduced by averaging the results of two or three  ${}^{40}$ K assays on each subject and by making consecutive three-day collections of urine.

Boileau et al. (30), however, have reported lower correlation co-efficients between LBW and UCE (r=.73 before conditioning and r=.57 after conditioning). A large within subject variation was found. Miller and Blyth (31) have reported individuals exhibiting as much as 20 percent variation in daily UCE.

Table 1 shows the relationship of UCE to selected measurements of body composition reported by various authors.

#### TABLE 1

Relationship of urinary creatinine excretion to selected measurements of body composition reported by various authors (30)

Author	Method	Correlation Coefficient
Miller & Blyth (31)	Body density	. 83
Best et al. (32)	Skinfolds	.61
Roessler et al. (33)	40 <sub>K</sub>	.56
Forbes & Bruining (29)	40 <sub>K</sub>	.98
Muldowney et al. (34)	42K dilution	. 90
Boileau et al. (30)	Water density	.73 <sup>a</sup>
		.57 <sup>b</sup>

<sup>a</sup>Before physical conditioning <sup>b</sup>After physical conditioning

Although the validity of employing UCE to estimate body composition is supported by the significance of its relationship to LBW, the accuracy and consistency with which this method can estimate body composition remains in question (35, 36, 37).

## Coronary Heart Disease Risk Factors

America is one of the nations with the highest death rates from coronary heart disease (CHD) (38). Certain risk factors for identifying individuals with a greater propensity for development of CHD have been studied internationally. The most prominent of these risk factors include: elevated serum cholesterol and blood pressure levels and smoking. Obesity, dietary intake, elevated serum triglyceride levels and physical inactivity are recognized as risk fators, but their relative contribution to the development or progression of CHD is less certain (8). The risk of developing CHD is not only proportional to the antecedent level of risk factors but also the risk increases with the number of risk factors involved (9).

### Serum Lipids

In a nine year follow-up of 3,168 men, Carlson et al. (39) have found that the rate of coronary heart disease (CHD) increases linearally with increasing triglyceride (TG) or cholesterol levels. Combined elevation of both lipids carries the highest risk.

Epidemiological studies of various population groups have suggested to many investigators that elevated serum cholcsterol is one of the major factors associated with CED (40). Truett et al. (41) have found that even mildly raised levels of serum cholesterol are considered to be risk factors of ischemic heart disease.

In the Western Collaborative Study, subjects with serum TG levels greater than 176 mg/dl had an annual incidence of CHD over three times that of subjects with levels under 100 mg/dl (42). Carlson (43) and Albrink et al. (44) have reported that serum TG were elevated in 80 percent of men with clinically evident CHD.

In 1975, Miller and Miller (45) postulated a negative correlation between levels of circulating high density lipoprotein and incidence of CHD. This hypothesis has received support from the epidemological studies of Castelli et al. (46) (Table 2) and Berg et al. (47). Gordon et al. (48) have reported that women who are obese, diabetic and have low levels of HDL have a very high risk of developing CHD.

TABLE 2

Prevalence of coronary heart disease by high density lipoprotein-cholesterol level, men aged 50-69 (46)

Rate/1000
180.0
123.6
94.6
77.9
77.9

The anti-atherogenicity of HDL may be explained by the hypothesis that plasma HDL delivers all of the cholesterol derived from tissues, including the arterial wall, to the liver for catabolism, via the lecithin-cholesterol acyl transferase mechanism (49).

## Physical Activity and Serum Lipids

Several studies have suggested that physical activity has no significant effect on serum cholesterol (50,51,52). Lewis et al. (53) conducted a study investigating the effects of physical activity on weight reduction in obese women. After 17 weeks of increased physical activity (2.5 miles of jogging and one hour of calisthenics per week), mean plasma cholesterol concentrations were not significantly changed.

Goode et al. (54) conducted a two-week study in which male subjects were exercised by treadmill to increase energy expenditure by one-sixth. Serum cholesterol levels of the exercisers did not differ significantly with the cholesterol levels of the control subjects, but the serum TG levels were significantly lower.

Other studies also have shown an inverse relationship between levels of physical activity and serum TG concentrations. Garcia-Palmieri et al. (55) reported a correlation of -.600 between serum TG levels and degree of physical activity. Hollosky et al. (51), Hurter et al. (56), and Wood et al. (57) have found fasting plasma TG concentrations to be lower in physically well-trained men than in their sedentary counterparts.

The reduction in serum triglyceride concentrations could be explained by the fact that epinephrine and norepinephrine increase both the lipase activity of adipose tissue (58) and the breakdown of triglyceride in isolated muscle preparations (59) and strenuous exercise is known to increase the discharge of these hormones (60).

High density lipoprotein concentrations have been reported to increase with physical activity (61,53). Lopez et al. (62) have found that after a seven-week program of brief, intense exercise, high density lipoprotein (HDL) increased in medical students. Lehtonen and Viikari (50) have made a positive correlation between the number of kilometers run weekly and the amount of HDL present in the blood. Wood et al. (57) have reported that HDL levels increased in men who did long distance running in their leisure time.

Miller et al. (63) conducted a study in which 27 urban and 25 rural Jamaican men, between the ages of 40 and 76 years, were assessed for body fat,

total serum cholesterol, HDL, low density lipoprotein (LDL), triglyceride, and blood glucose. Rural farmers engaged in manual labor had less body fat, and lower fasting levels of cholesterol, LDL, and triglyceride than urban businessmen. Mean HDL levels were considerably higher for farmers than businessmen. The results of this study imply that physical activity may be beneficial when considering risk factors often associated with greater propensity for coronary heart disease development.

## Body Weight and Serum Lipids

Obesity is often associated with low levels (less than 35 mg/dl) of high density lipoprotein (7). Results from the Framingham Study (48) have suggested that high density lipoprotein (HDL) is correlated negatively (r=-.25) with relative weight (actual weight/ideal weight X 100).

As noted earlier, rural Jamaican farmers who were leaner than their sedentary counterparts had considerably higher mean HDL levels, and lower triglyceride, total serum cholesterol, and low density lipoprotein levels (Table 3) (63).

#### TABLE 3

Plasma lipoprotein lipids and triceps skinfold measurements (63)

Parameters	Urban (n=27) Mean	Rural (n≖25) Mean
Cholesterol (mg/dl)	222.0	194.0
High density lipoprotein (mg/dl)	63.0	83.0
Low density lipoprotein (mg/dl)	130.0	94.0
Triglyceride (mg/dl)	146.0	88.0
Tricep skinfold (mm)	11.5	5.2

Kannel et al. (64) have investigated the effects of obesity on serum lipid levels in 5,209 men and women. Results of the Framingham Study suggest that relative weight is correlated with serum cholesterol and triglyceride and inversely correlated with high density lipoprotein cholesterol. In general, relative weight was related more closely to blood lipid levels among men than women of the same age. The strength of that association was greater in younger individuals than older.

Early research, conducted by Walker (65), has suggested that weight reduction can depress elevated serum cholesterol levels. Berkowitz (66) has found that weight reduction improved serum cholesterol concentration in 100 obese men and women, but the specific effects of that improvement were erratic.

## Blood Pressure and Physical Activity

Lewis et al. (53) conducted a study in which increased physical activity, consisting of jogging two and one-half miles and one hour of calisthenics per week, was the primary focus of a 17-week weight reduction program. Mean resting heart rate and systolic blood pressure were slightly lower at the time of re-evaluation. Heart rate decreased from 65.5 to 62.4 beats per minute, and systolic blood pressure decreased from 116.8 to 112.0 mm/Hg.

Montoye et al. (67) assessed the physical activity patterns of 1,700 adult males. Energy cost was estimated and expressed as a ratio of the work metabolic rate divided by the basal metabolic rate (WMR/BMR). Men were then grouped according to estimated levels of physical activity (highest, intermediate, and lowest). In general, the most active men had the lowest mean systolic and diastolic blood pressures (Table 4).

### TABLE 4

Physical activity group	Mean systolic (mm/Hg)	(n)	Mean diastolic (mm/Hg)	(n)
Highest	137.3	275	77.9	274
Intermediate	137.9	816	78.5	815
Lowest	140.4	270	80.8	270

Systolic and diastolic blood pressures and physical activity level (67)

Vitale's (68) description of the physiological effects of physical activity on blood pressure would support the findings of Lewis et al. (53). During exercise, collateral circulation between all major arteries is increased and there is better diffusion between the capillaries and the working tissues. The improved blood flow takes an additional work load off the heart. The enlargement and increase in the number of blood vessels decrease resistance to blood flow and lower abnormally high blood pressure, both at rest and during exercise.

#### Blood Pressure and Body Weight

Wilcox (69) has reported that body weight was correlated with systolic  $(\pi=.42)$  and diastolic  $(\pi=.38)$  blood pressure in 70 massively obese women.

In a study conducted by Montoye et al. (67), approximately 1,700 men (16 to 54 years old) were assessed for systolic and diastolic blood pressure, body fatness (determined by the sum of four skinfolds: triceps, subscapular umbilical and suprailiac), and level of physical activity (determined by the work metabolic rate/basal metabolic rate ratio). Men were grouped according to level of physical activity (highest, intermediate and lowest). Men who were leaner and more physically active had lower blood pressures than men who were heavier and less active.

The Framingham study (70) has shown that changes in relative weight were paralleled by changes in systolic blood pressure. These changes were small and unrelated to initial relative weight. Changes were less pronounced in women than in men.

### Dietary Intake and Serum Lipids

Serum cholesterol concentrations in population studies have been correlated with the proportion of dietary kilocalories derived from fat (71). Saturated fats appear to be the main dietary component responsible for this relationship (72).

Nutrient intake in relation to serum triglyceride and cholesterol concentrations has been investigated by Nichols et al. (73). Twenty-four hour dietary recall interviews were conducted among 2,000 men and women in the community of Tecumseh, Michigan. Cholesterol and triglyceride levels were unrelated to quantity, quality, or proportion of fat, carbohydrate or protein consumed. Balart et al. (8) have reported similar findings. Daily dietary intakes of cholesterol and carbohydrate were not correlated with serum cholesterol concentration.

The present study included two weeks of preliminary testing, 12 weeks of experimental treatment, two weeks of post testing, and a three-month follow-up evaluation.

#### Subjects

Volunteer subjects were solicited from the community via local newspaper advertisement. Study subjects were premenopausal, non-pregnant, adult women at least 15 pounds overweight according to acturarial statistics for height and weight (74). Approximately 58 women qualified for participation in the study. These subjects were pretested and randomnly assigned to one of two experimental groups: 1) diet therapy or 2) behavior modification. These two groups were subdivided into exercise and non-excercise groups.

#### Informed Consent Forms

This study was reviewed in accordance with University policy and the United States Department of Health Education and Welfare regulations regarding human subject research. Prior to preliminary testing, each subject was advised of all procedures and methods to be used. A signed consent form (Appendix) was then obtained from each subject.

## Preliminary Testing Period

Subjects were evaluated for the following parameters: 1) selected anthropometric measurements, 2) body composition, 3) urinary creatinine excretion, 4) systolic and diastolic blood pressures, 5) serum lipids (serum triglycerides, high and low density lipoproteins, total cholesterol, and high density lipoprotein/cholesterol ratio), and 6) three-day dietary food intakes.

## Anthropometric Measurements and Body Gomposition

Body density, height, weight, and selected anthropometric measurements were taken in the Exercise Physiology Research Laboratory at Kansas State University, Manhattan, Kansas. Subjects were underwater weighed to determine body density. Body density was converted to percentage fat using the equation devised by Brozeck et al. (12):

percentage fat = 100 X  $\frac{4.570}{\text{density}}$  - 4.142

Total body fat was calculated by multiplying total body weight by percentage fat. Lean body weight was determined by subtracting total body fat from total body weight. Detailed methods and procedures for this portion of the investigation are discussed in the M.S. thesis by Moyer (1979).

## Urinary Creatinine Excretion

Each subject collected a complete, 24-hour urine sample in a sealable container containing approximately 10 ml of toluene (a preservative). Urine volume was measured and urinary creatinine excretion was determined by the method of Sieverd (75).

#### Blood Pressure

Systolic and diastolic blood pressures were taken with a stethascope and sphygmomanmeter. Subjects were seated in a chair during blood pressure evaluation.

## Serum Blood Lipids

Fasting blood samples were taken by venipunture under medical supervision at Kansas State University, Manhattan, Kansas. The absence of chylomicrons in the electrophoresis pattern together with the standing turbidity refrigerator test served to indicate that a fasting blood sample had been obtained. Prior to veripuncture, each subject completed a questionnaire pertaing to factors affecting serum lipid levels and incidence of coronary heart disease (Appendix). Blood serum samples were sent to Dr. Manford Morris, Department of Pediatrics; University of Arkansas Medical Sciences, for lipoprotein electrophoresis of triglycerides, high density lipoprotein cholesterol, and total serum cholesterol. Total cholesterol was determined by the method of Rudel et al. (76), triglycerides by the method of Sardesai and Manning (77), agarose electrophoresis by method of Nobel (78), and high density lipoprtein cholesterol by the method of Lopes-Virella (79). Blood Serum Analysis

Blood serums were analyzed by SMA-12 auto-analyzer at the College of Veterinary Medicine, Kansas State University, Manhattan, Kansas. Blood serum constituents analyzed include: glucose, urea, creatinine, calcium, phosphorous, alkaline phosphatase, glutamic pyruvic transaminase, sodium, potassium, carbon dioxide, total protein, albumin, and chloride.

## Three-day Dietary Intakes

Each subject recorded quantity and type of food eaten for one week (Appendix). From this one-week period, three days were randomnly selected to assess dietary intake. Daily nutrient intake was determined as the mean of the three-day dietary assessment. Nutrients assessed include: kilocalories; protein, fat, carbohydrate, fiber, ash, saturated fatty acids, polyunsaturated fatty acids, monosaturated fatty acids (gm/day); sodium, potassium, dietary cholesterol (mg/day); and calcium, iron,vitamin A, ascorbic acid, phosphorous, thiamine, riboflavin, and niacin (percentage of Recommended Dietary Allowances). The DIETETIC COM-PAK program, written by the Department of Nutrition and Dietetics at the University of Missouri Medical Center, and modified by the Department of Dietetics, Restaurant and Institutional Management, Kansas State University, was used to compute nutrient intake.

Several subjects did not keep complete, accurate records. Quantities of food consumed were often vague ("three handfuls of nuts") or completely omitted ("mashed potatoes and gravy"). Consequently, preliminary three-day dietary intakes were not assessed.

## Experimental Testing Period

Subjects were assigned randomnly to one of two experimental groups for 12 weeks: 1) diet therapy or 2) behavior modification. These two groups were subdivided into exercise and non-exercise groups. Women already participating in some form of leisure physical activity were assigned to one of the exercise groups.

During the 12-week experimental period, subjects attended weekly instructional meetings lasting approximately one hour. Subjects were weighed (street clothes, without shoes) on a Homs full-capacity balance beam scale prior to the beginning of each meeting.

Selected presentations were given to all subjects during two of the 12 meetings. Topics discussed include: factors associated with increased risk of coronary heart disease development and factors associated with the origin and propagution of obesity.

During the twelfth week of the experimental treatment period, subjects were instructed to carefully measure and define all food and drinks consumed

<sup>2</sup>Douglas Homs Corp., Belmont, California.

during the next seven days. From this one-week period, three days were selected randomnly to assess dietary intake. Daily nutrient intake was determined as the mean of the three-day dietary assessment.

## Diet Therapy Group

The goal of the diet therapy group was to lose weight by restricting total energy intake. Individual energy requirements were estimated for each subject based on the energy needs for basal metabolic rate, specific dynamic action, and physical activity. To achieve a weight loss of approximately one pound a week, 500 kcal was subtracted from the estimated total energy needs of each subject (Appendix).

Women in the diet therapy group received the "American Dietetics Association Exchange Lists for Meal Planning" (10). This exchange list provided subjects with a guideline to assist them in designing their dietary regimens which were to be followed throughout the study. The number of food exchanges allowed from each food group was determined by the individual's energy needs. Approximately 20 percent of the total energy allowance was derived from protein, 30 percent from fat, and 50 percent from carbohydrate. (An example 1200 kcal diet is shown in the Appendix.)

A short questionnaire (Appendix) was completed by women in this group six weeks after the beginning of the experimental period. The primary purpose of the questionnaire was to evaluate subject compliance to an energyreduced, food-exchange diet.

### Diet Therapy and Exercise Croup

In addition to decreasing energy intake by dietary modification, subjects in the diet therapy plus exercise group were instructed to increase energy expenditure by developing individualized programs of physical activity. An exercise physiologist assissted subjects in program planning. During the last week of the experimental period, each subject kept a physical activity record. MET<sup>3</sup> and energy expenditure (kcal) levels were estimated using tables devised by Fox et al. (80).

## Behavior Modification Group

Subjects in the behavior modification group were to lose weight and maintain weight loss by developing appropriate eating patterns rather than following a special, restrictive diet. Weekly homework assignments, taken from Ferguson's manual (11), were given to subjects in this group. The purpose of these assignments was to teach subjects to reduce energy intake by modifying their obesity-related behaviors. Assignment topics given include: 1) habit awareness, 2) cue elimination, 3) changing the act of eating, 4) behavior chains and alternate activities, 5) behavioral analysis, progress, and problem solving, 6) pre-planning, 7) cue elimination-II 8) energy use 9) snacks, cues, and holidays, and 10) environmental support. <u>Behavior Modification and Exercise Group</u>

Treatment of this group was identical to that of the behavior modification group except physical activity was added to the program as described in the diet therapy and exercise group section.

## Post Testing Period '

A post-testing period of two weeks followed the 12-week experimental treatment period. During this time, all parameters were re-evaluated.

<sup>3</sup>One MET level is defined as the amount of oxygen consumed during resting metabolism which is equivalent to 3.5 ml 0<sub>2</sub>/kg/min.

## Post Experimental Follow-Up

Twelve weeks after completion of the post-testing period, subjects were evaluated for changes in anthropometric measurements and body composition (total body weight, percentage fat, lean body weight, and total body fat).

## Statistical Treatment of the Data

The analysis of variance test (31) was used to determine if any significant differences had occurred among subjects in the four experimental groups between preliminary and post treatment and post treatment and follow-up evaluation. The 0.05 level of probability was selected as the statistical criterion for significance. The paired t-test (81) was used to determine if tue mean change or difference from one evaluation period to the next, for subjects within a group, was significant (P<.05). The attrition rate was slightly less for subjects in the two behavior modification groups (41%) than for subjects in the two diet therapy groups (45%). A short questionnaire (Appendix) evaluated subject compliance to an energy reduced, food-exchange diet. Evaluation of completed questionnaires suggested that subjects in the diet therapy groups were not closely following their diets. The average score was 3.5 on a scale from 1 to 5 with 1 representing very good adherence to the diet and 5 representing very poor adherence to the diet. Brightwell (23) has suggested that motivation is more easily developed in behavior modification weight loss programs than in traditional weight loss programs. The results of the present study tend to support Brightwell's observation.

## Body Composition

Mean pre and post treatment and follow-up values for body composition parameters are presented in Table 5. The two series of difference values reflect changes which occurred between pre and post treatment measurement (12-week experimental period) and changes which occurred between post treatment and follow-up (12 weeks) evaluation. Subjects served as their own controls, and so mean differences or changes for each group are based solely on the values of subjects particiapting in both pre and post treatment tests and post treatment and follow-up measurements. Difference-I values show that only subjects in the behavior modification (BM) group lost a significant ( $P\leq.05$ ) amount of weight (-6.9 pounds). Subjects in the behavior modification plus exercise (BM+EX) group lost 4.9 pounds,

TABLE 5

Fre and post treatment and follow-up values of selected body composition parameters for obese (mean-33% fat) women aged 20-52 years

Parameters				Gr	oups				
	BM+EX <sup>a</sup>	(u)	BMb	(u)	DT+EXC	(u)	DTd	(u)	
Weight (1b)									
Pre	178.4+7.8 <sup>e</sup>	(6)	147.0+8.3	(8)	137.9+ 9.6	(9)	162.5+ 8.3	(8)	
Post	169.7+7.2	(8)	141.17.7	(2)	139.0H 9.1	(2)	153.0H 7.7	(1)	
Follow-up	170.47.6	(8)	$141.1\overline{+}7.6$	(8)	139.0+ 8.8	(9)	161.0+ 7.6	(8)	
Difference - I <sup>f</sup>	-4.0	(8)	-6.9*	22	$+1.2^{-}$	(4)	-1.7	(9)	
Difference - 118	+0.6	(8)	+1.9	6	-0.4	(2)	+1.6	(9)	
Fotal body fat (1b)									
Pre	71.2+5.6	(6)	55.0+6.0	(8)	45.8+ 8.5	(4)	64.4+ 6.9	(9)	
Post	64.5+4.8	(8)	50.8+5.6	(9)	50.54 6.8	(4)	52.1+ 5.6	(9)	
Follow-up	63.477.6	(2)	44.778.5	(4)	52.4+17.0	5	57.2+12.0	(2)	
Difference - I	-3.7	(8)	-4.9*	(9)	+1.1	(E)	-1.0	(2)	
Difference - II	-0.5	(2)	+2.8	(3)	-0.4	(1)	+0.2	(2)	
ean body weight (1b)									
Pre	107.2+3.8	(6)	91.5+4.1	(4)	92.5+ 5.7	(4)	97.9+ 4.7	(9)	
Post	$105.2 \pm 3.9$	(8)	90.144.5	(9)	87.24 5.5	(4)	96.94 4.5	(9)	
Follow-up	$103.8 \pm 6.2$	(2)	91.476.9	(4)	94.7 <del>1</del> 13.9	(1)	100.1+ 9.8	(2)	
Difference - I	-0.3	(8)	-1.4	(9)	+0.2	(3)	-0.4	(2)	
Difference - II	+0.3	(2)	+1.3	(3)	+0.5	(1)	+1.8	(2)	
ercentage body fat									
Pre	39.3+1.8	(6)	37.4+1.9	(8)	33.1+ 2.7	(4)	39.1+ 2.2	(9)	
Post	$37.5\overline{+1.8}$	(8)	35.8+2.1	(9)	36.7+ 2.6	(4)	34.8+ 2.1	(9)	
Follow-up	37.1+2.8	(2)	32.3+3.1	(4)	35.64 6.3	E	36.5+ 4.4	(2)	
Difference - I	$-1.2^{-1.2}$	(8)	-1.9*	(9)	+0.5	(3)	-0.3	(2)	
Difference - II	+0.1	(5)	+0.4	(3)	-0.3	(1)	-0.2	(2)	
<sup>a</sup> BM+EX=behavior modifica	tion plus exe	srcise.	bBM=behavior	modifi	cation. <sup>c</sup> DT+EX	=diet	therapy plus	exercise.	

SDifference-II represents the change or difference between post and 12-week follow up measureeStandard error. <sup>I</sup>Difference-I represents the change or difference between pre and post Note: treatment period was 12 weeks in duration. \*Significant at the 0.05 level. measurement values. <sup>d</sup>DT=diet therapy. ment values.

subjects in the diet therapy plus exercise (DT+EX) group gained 1.2 pounds, and subjects in the diet therapy (DT) group lost 1.7 pounds. Assessment of physical activity records indicated that subjects in the two exercise groups expended approximately 250 kcal, daily, in some form of additional, leisure physical activity. In general, subjects in the two behavior modification groups lost more weight than subjects in the diet therapy groups. Studies conducted by Stuart (6), Stundkard (5), and Levitz (20) also have supported the hypothesis that behavioral approaches to obesity are more successful in terms of weight loss than dietary educational programs.

Difference-II values (Table 5) indicate that no significant changes occurred in body composition between post treatment and follow-up evaluation, however, subjects in the EM+EX group were slightly more successful at maintaing weight loss (+.6 pounds) than subjects in the EM group (+1.9 pounds) and subjects in the DT group (+1.6 pounds). Subjects in the DT+EX group lost some (-.4 pounds) of the weight they had gained over the experimental period. The length of time elapse allowable from post treatment measurement to follow-up evaluation was limited. A two or three year time span between evaluation periods would have been a more interesting and reliable test.

Harris and Hallbauer (82) have conducted a similar 12-week study investigating the effects of diet therapy, behavior modification and exercise on weight reduction. The results of their study indicated that subjects in a behavior modification plus individualized exercise program lost an average of 7.0 pounds while subjects in the behavior modification only program lost 6.0 pounds. The control group, receiving only dietary

counseling, lost just 4.5 pounds. No significant differences occurred between subjects in the three groups. Results of a seven month follow-up showed that subjects in the behavior modification group maintained their previous loss. Subjects in the behavior modification plus exercise group lost two more pounds, and subjects in the control group gained back 7.5 pounds.

In contrast to the study conducted by Harris and Hallbauer (82), subjects in the EM+EX group did not lose more weight than subjects in the EM group over the 12-week experimental period. The reason for this is unknown. Individual motivation, body metabolism, or a combination of these factors may have accounted for some of the differences in group performance. Both studies have suggested that subjects in the behavior modification programs lose more weight than subjects in diet therapy programs, and that exercise facitlates weight loss maintenance.

The data in Table 5 show that only subjects in the EM group lost a significant ( $P \leq .05$ ) amount of total body fat (-4.9 pounds), and percentage fat (-1.9%) between pre and post experimental treatment. Total body weight was correlated positively with total body fat (r=.96) and percentage fat (r=.89). No significant changes occurred in lean body weight for subjects in any of the four experimental groups. Detailed results of anthropometric and body composition parameters are discussed in the M.S. thesis by Moyer (1979).

#### Urinary Creatinine Excretion

Normal urinary creatinine excretion (UCE) values for adults range from 1-1.8 gm/24 hr (75). In general, mean UCE values (Table 6) for

TABLE 6

Pre and post treatment values of urinary creatinine excretion (UCE) for obese (mean=38% fat) women aged 20-52 years

			Grou	bs					
JEINARY CFEALININE EXCFECION (g/24hr)	BM+EX <sup>a</sup>	(u)	BMb	(u)	DT+EX <sup>c</sup>	(u)	DTd	(u)	
Pre	1.58 <u>+</u> .11 <sup>e</sup>	6	1.284.11	(8)	0.984.12	(9)	1.35±.11	(8)	
Post	1.31+.09	(6)	1.20±.11	(9)	1.124.11	(9)	1.34±.09	(6)	
Difference <sup>f</sup>	-0.278*	(1)	+0.01	(9)	+0.148	(9)	-0.10	(8)	

measurement values. BUCE difference values for subjects in these groups are significantly different from one another at the P<u>C</u>0.05 level. \*Significant at the P<u>C</u>0.05 level. Note: treatment period was 12 weeks in duration. <sup>c</sup>DT+EX=diet therapy plus exercise. <sup>t</sup>Difference represents the change or difference between pre and post bBM=behavior modification. <sup>a</sup>BM+EX=behavior modification plus exercise. eStandard error. dDT=diet therapy.

all four groups were within normal limits. Urinary creatinine excretion was significantly ( $P\leq.05$ ) decreased from pre to post treatment evaluation for subjects in the EM+EX group. (Difference values in Table 6 indicate this phenomenon). The change in urinary creatinine excretion for subjects in the EM+EX group (-.27 g/24 hr) was significantly ( $P\leq.05$ ) different from the change in UCE for subjects in the DT+EX group (+.14 g/24 hr). Observation by Srivastava (83) have suggested that urinary creatinine excretion increases following physical activity. This observation would account for the slight increase found in subjects in the DT+EX group, but his work would not substantiate the significant ( $P\leq.05$ ) decrease in urinary creatinine excretion exhibited by subjects in the EM+EX group.

Many investigators (30,31,34) have shown that urinary creatinine excretion is related directly to lean body weight. Correlations ranging from .56 (33) to .98 (29) have been reported. The results of the present study indicated that urinary creatinine excretion was slightly correlated with lean body weight (as determined by water densiometry) before ( $r_{pre}$ =.58) and after ( $r_{nost}$ =.53) experimental treatment.

The somewhat low correlation and the significant  $(P\leq.05)$  decrease in UCE by subjects in the BH+EX group may have been the result of incomplete urine collection, UCE variation or a combination of these factors. Forbes and Druining (29) have reported an average co-efficient variation of 6.9 percent for 34 adults and children. The co-efficient variation was based on three, 24 hour urine collections. The use of UCE to determine body composition of individuals seems to be limited by the likelihood of large within subject variation.

## Serum Lipids

Mean pre and post treatment values for fasting serum triglyceride, cholesterol, low density lipoprotein  $\left(LDL\right)^4$ , high density lipoprotein (HDL) and HDL/cholesterol ratio are presented in Table 7.

## Cholesterol and Triglyceride

The data indicate that initial serum lipid values for subjects in both exercise groups were slightly lower than for subjects in the nonexercise groups. One of the factors considered prior to group assignment was current participation in leisure physical activity. Ideally, occupational as well as leisure physical activity would have been considered in group assignment. Miller et al. (63) also have noted this seemingly beneficial effect of exercise. Jamaican farmers engaged in manual labor as a result of their occupations had lower fasting serum levels than urban businessmen not engaged in occupational physical activity.

The present study indicates that subjects in the BM+EX group significantly ( $P_{\leq}$ .05) decreased their cholesterol levels by 17.7 mg/dl. No significant changes occurred for subjects in the other three groups.

Lewis et al. (53) however, have found that physical activity had no effect on cholesterol levels. Twenty-two obese women jogged two and onehalf miles and participated in one hour of calisthenics each week for 17 weeks. Mean plasma cholesterol concentrations did not change significantly from the initial to the final evaluation. Lewis et al.'s (53) research supports the response noted for subjects in the DT+EX group.

<sup>&</sup>lt;sup>4</sup>LDL was calculated using Fredrickson's formula: LDL = total cholesterol - HDL cholesterol - triglyceride/5. This formula is only applicable when triglyceride levels are less than 400 mg/dl. All subjects in the present study had levels less than 400 mg/dl.

Tintdo in Table 7

arameters				Group	S			
	BM+EX <sup>a</sup>	(u)	BMb	(u)	DT+EX <sup>C</sup>	(u)	DTd	(u)
riglyceride (TG-mg/d1)								
Pre	66.9+12.30	(8)	83.8+12.30	(8)	58.9+13.10	6	98.0+11.50	(6)
Post	55.94 8.50	(8)	44.3= 9.80	(9)	59.7+ 9.80	(9)	79.0H 8.00	(6)
Difference	-13.4	(2)	-14.7	(9)	+5.2	(9)	-19.0*	(6)
holesterol (mg/dl)								
Pre	186.8+10.10	(8)	196.9+10.10	(8)	176.7+10.80	(2)	202.1+ 9.50	(6)
Post	168.1+11.40	(8)	$192.7\overline{+}13.10$	(9)	182.5+13.10	(9)	200.2+10.70	(6)
Difference	-17.7*	(2)	0.0	(9)	+6.0	(9)	-1.9	(6)
ow density lipoprotein8	(LDL-mg/d1)							
Pre	125.9+ 9.10	(8)	127.4+ 9.10	(8)	119.4+ 9.80	(2)	138.4+ 8.60	(6)
Post	109.6+10.50	(8)	128.6+12.20	(9)	118.2+12.20	(9)	139.0F 9.90	(6)
Difference	-13.9	6	-11.6	(9)	-1.0	(9)	-0.6	(6)
igh density lipoprotein	(HDL-mg/d1)							
Pre	47.5+ 3.40 <sup>e</sup>	(8)	52.8+ 3.40	(8)	45.6+ 3.70	6	44.1+ 3.20	(6)
Post	47.47 3.50	(8)	55.24 4.00	(9)	52.3+ 4.00	(9)	45.47 3.30	(6)
Differencef	-1.1	(2)	0.0	(9)	+6.0	(9)	+1.3	(6)
DL/cholesterol ratio								
Pre	0.26+0.02	(8)	0.27+0.02	(8)	0.26+0.02	(2)	0.22+0.02	(6)
Post	0.29+0.02	(8)	$0.29 \pm 0.02$	(9)	0.29+0.02	(9)	$0.23 \pm 0.02$	(6)
Difference <sup>I</sup>	+0.03*	6	-0.01	(9)	+0.02*	(9)	+0.01	(6)

treatment measurement values. Scalculated by Fredrickson et al. (84) formula:LDL = cholesterol - HDL - TG/5. <sup>a</sup>BM+EX=behavior modification plus exercise. <sup>b</sup>BM=behavior modification. <sup>c</sup>DT+EX=diet therapy plus exercise. <sup>I</sup>Difference represents the change or difference between pre and post \*Significant at the  $P \leq 0.05$  level. Note: treatment period was 12 weeks in duration. eStandard error. dDT=diet therapy.

Mann et al. (85) have reported that decreases in weight have been associated with decreases in serum cholesterol. Berkowiz (66) has shown that weight reduction lowers serum cholesterol, but the specific effects are erratic. Similarly, early studies by Walker (65) have indicated that weight reduction has no significant effect on serum cholesterol unless levels are initially elevated. In the present study, total body weight was not correlated with serum cholesterol. All average group cholesterol levels were less than 220 mg/dl, however, several individual cholesterol values were elevated (Appendix). In a few of these individual cases, cholesterol corcentrations decreased with weight reduction (Subject 9010: 167.2 to 164.9 pounds, cholesterol-238 to 155 mg/dl; Subject 9014: 179.8 to 176.8 pounds, cholesterol-234 to 187 mg/dl; Subject 9043: 231.3 to 222.3 pounds, cholesterol-260 to 253 mg/dl).

Kannel et al. (64) have investigated the effects of obesity on serum lipid levels in 5,209 men and women. Results of this study suggest that relative weight is correlated (r=.10) with cholesterol concentrations in middle-aged women. In the present study, serum cholesterol levels were not correlated with total body fat mass. The sample size of the present study was very small compared to the population of the Framingham Study. This may have partially accounted for the discrepancy between the two investigations.

The data in Table 7 show that subjects in the DT group significantly  $(P\zeta,05)$  decreased triglyceride concentration. The reason for this is unknown.

Acquisition of complete, initial dietary intake records would have allowed the assessment of changes in nutrient intake and the effects of those changes on serum cholesterol and triglyceride levels. The present study was limited in this respect.

### Righ Density and Low Density Lipoprotein (HDL, LDL)

Epidemiological studies have shown that low levels of HDL and high levels of LDL are rélated directly to coronary heart disease prevalence. Gordon et al. (7) have reported that individuals with HDL levels less than 35 mg/dl have eight times the risk of developing coronary heart disease when compared with individuals having HDL levels greater than 65 mg/dl. The Cooperative Lipoprotein Phenotyping Study (46) has indicated that individuals with LDL levels greater than 130 mg/dl have twice the risk of developing coronary heart disease in comparison to individuals having levels less than 140 mg/dl. Mean HDL and LDL values for subjects in the present study (Table 7) were greater than 35 mg/dl and less than 140 mg/dl, respectively. No significant changes occurred in LDL or HDL concentrations between pre and post treatment evaluation.

The relationship between HDL and physical activity has been studied to some extent by Lopez et al. (62), Enger (61) and Lehtonen and Viikari (50). Increases in physical activity are often associated with increases in HDLcholesterol. Lewis et al. (53) have investigated the effects of physical activity on weight reduction in 22 obese (mean = 40% fat) middle-aged women. After a 17-week exercise program consisting of 2.5 miles of jogging and one hour of calisthenics per week, mean HDL levels did not change significantly, but the HDL/cholesterol ratio was increased significantly (P $\leq$ .05) (pre=.36, post=.43, increase =.05). Similarly, the results of the present study (Table 7) show that HDL values were not altered significantly by experimental treatment, but the HDL/cholesterol ratio was significantly (P $\leq$ .05) increased for subjects in the two exercise groups (BH+EX = +.03, DT+EX = +.02).

The inverse relationship of HDL-cholesterol to coronary heart disease is a relatively new approach to assessment of coronary heart disease development. Whether the HDL level is a guide to therapy as well as a risk indicator remains to be determined. The full effects of physical activity, nutrient intake, serum lipids and body composition on coronary heart disease risk and life expectancy are still unknown. More conclusive research is needed.

### Blood Pressure

Table 8 summarizes the mean pre and post treatment values for all four experimental groups. Difference values indicate changes which may have occurred between initial and final evaluation. No significant changes were found in either systolic or diastolic blood pressures for subjects in any of the study groups. Diastolic blood pressure was correlated (r=.48) with body weight. Wilcox (69) has reported a slightly lower correlation (r=.38) in 70 massively obese women (mean age=34 years, mean weight=228 pounds, mean height=65 inches).

## Blood Serum Analysis

Mean serum constituent values are presented in Table 9. Initial analysis of those values indicated that serum constituents for all subjects were within normal limits.

#### Dietary Intake

The post treatment data in Table 10 indicate that all four experimental groups met two-thirds of the Recommended Dietary Allowances for all nutrients but iron. Subjects in all four groups ingested about 1300 kcal with 20 percent of the energy being derived from protein, 30 percent from fat, and 50 percent from carbohydrate. Even though average energy intakes were not significantly different among subjects in the four groups, the data indicate that subjects in

TABLE 8

Pre and post treatment values for systolic and diastolic blood pressures in obese (mean=30% fat) women aged 20-52 years

								-	11
Blood pressure	BMHEXa	(u)	BMb	(n)	B DT+EX <sup>C</sup>	(u)	pTd	(u)	
Systolic blood pressure (	(mm/Hg)								
Pre 12 weeks Post 12 weeks Difference <sup>f</sup>	124.5 <u>+</u> 4.8 <sup>e</sup> 113.4 <u>+</u> 3.7 -8.3	(9) (9) (9) (9) (9) (9) (9) (9) (9) (9)	$120.9+4.4 \\ 108.3+4.3 \\ -7.6$	(2)	106.0+5.2 116.3+3.9 +10.8	666	$\frac{111.8+4.1}{107.1+3.5}$	868	
Diastolic blood pressure	(mm/Hg)								
Pre 12 weeks Post Difference	85.0+3.3 77.1+3.1 -4.7	(9) (8) (9)	75.1 $\pm$ 3.1 72.5 $\pm$ 3.4 -1.4	(2)	68.0 <u>+</u> 3.6 71.9 <u>+</u> 3.2 +4.4	838	76.8+2.9 75.9+2.8 +1.4	868	

<sup>C</sup>DT+EX=diet therapy plus exercise. <sup>a</sup>BM+EX=behavior modification plus exercise. <sup>D</sup>BM=behavior modification. <sup>D</sup>DT+EX=diet therapy plus exvici DT=diet therapy. <sup>E</sup>Standard error. <sup>D</sup>Difference represents the change or difference between pre and post treatment period was 12 weeks in duration. Note: measurement values. dDT=diet therapy.

TABLE 9

Pre treatment blood serum constituent values for obese (mean=38% fat) women aged 20-52 years

	(0-II) VIIII	$BM^{\prime}$ (n=8)	DT+EX <sup>C(n=</sup> ))	DTd /
lucose (mg/dl)	81.5 <u>+</u> 1.60 <sup>e</sup>	74.6+2.71	77.7+1.73	77.3+2.99
rea (mg/d1)	12.7+1.20	13.8+1.01	-11.4+1.19	13. 3+0. 84
reatinine (mg/dl)	0.8+0.03	0.8+0.03	-0.8+0.02	0.840.03
alcium (mg/dl)	8.5+0.09	8.5+0.14	8.7+0.11	11 0+7 8
iosphorous (mg/d1)	3.6+0.08	3.7 <u>+</u> 0.11	3. 7+0.13	3.6+0.14
kaline phosphatase (I.U.)	68.6 <u>+</u> 5.34	65.3+3.75	61.4+7.04	18 0+6 69
utamic pyruvic (units/ml) transaminase	$14.4 \pm 3.27$	$11.6 \pm 1.28$	11.9+1.29	17.1+3.76
dium (mEq/L)	138.1±0.47	139.0+0.65	138.640 37	130 110 75
tassium (mEq/L)	4.4+0.20	ar 040 %		C/ .UT1.6C1
the state of the s	1	01.016.4	4.6+0.17	4.6+0.10
TDON GIOXIDE (MM/L)	27.9+1.01	26.1+0.72	26.6+0.92	26.8+1.18
tal protein (mg/d1)	6.9 <u>+</u> 0.15	7.0+0.07	6.9+0.16	7.1+0.15
(ID/mg) uimnc	$4.2 \pm 0.12$	4.3+0.05	4.2+0.10	4.3+0.07
loride (mEq/L)	106.0+0.94	$105.6\pm0.56$	106.9+1.52	105.7+0.69

	women
	fat)
	38%
	8
	(mean
	obese
	for
	Intake
	nutrient
	of
	values
10	treatment
TABLI	Post

aged 20-52 years

		Grou	ps		
Wutrient (daily intake)	BM+EX <sup>a</sup> (n=9)	вм <sup>b</sup> (n=8)	DT+EX <sup>C</sup> (n=7)	DT <sup>d</sup> (n=9)	
Kilocalories	1506+143.3 <sup>e</sup>	1165+143.3	1344+143.3	1305+134.4	
Protein (gm)	727 5.3	557 5.3	50.4 5.3	584 4.9	
Fat (gm)	737 8.1	42 <b>7</b> 8.1	55+ 8.1	567 7.5	
Carbohydrate (gm)	135+ 20.0	145+ 20.0	163+ 20.0	146+ 18.7	
Fiber (gm)	2+ 0.4	2+ 0.4	2+ 0.4	2+ 0.4	
Calcium (% of RDA)	77+ 12.6	51+ 12.6	61+ 12.6	827 11.8	
Phosphorous (% of RDA)	120+ 11.4	96+ 11.4	99+ 11.4	109+ 10.6	
Iron (% of RDA)	56+ 6.9	57+ 6.9	46+ 6.9	477 6.4	
Sodium (mg)	2511+302.9	1510-302.9	2207-302.9	18067283.3	
Potassium (mg)	1513+158.2	1324+158.2	1531+158.2	14157148.0	
Vitamin A (% of RDA)	111- 23.2	81+ 23.2	69+ 23.2	787 21.7	
Thiamine (% of RDA)	79+ 9.9	68+ 9.9	717 9.9	747 9.2	
Riboflavin (% of RDA)	98+ 12.2	89+ 12.2	69+ 12.2	837 11.4	
Niacin (% of RDA)	98+ 11.1	927 11.1	$75\overline{+}11.1$	$82 \pm 10.5$	
Ascorbic acid (% of RDA)	139- 31.1	150- 38.1	145+ 38.1	$147 \pm 35.6$	
SFA <sup>f</sup> (gm)	17+ 2.4	8+ 2.4	15+ 2.4	157 2.3	
UFAB (gm)	20+ 2.7	10+ 2.7	147 2.7	157 2.5	
PUFA <sup>h</sup> (gm)	10+ 2.1	47 2.1	77 2.1	6∓ 2.0	
Dietary cholesterol (gm)	340+ 65.5	$242 \pm 65.5$	175+ 65.5	251+61.3	

<sup>a</sup>BNHEX-behavior modification plus exercise. <sup>b</sup>BN-behavior modification. <sup>c</sup>DTHEX-diet therapy plus exercise. <sup>d</sup>DT-diet therapy. <sup>e</sup>Standard error. <sup>f</sup>SFA-saturated fatty acids. <sup>B</sup>UFA-monounsaturated fatty acids. <sup>b</sup>DUFA-polyunsaturated fatty acids.

the Bif group consumed the least amount of energy (1,165 kcal) and lost the greatest amount of weight (-6.9 pounds). Subjects in the two behavior modification groups adopted nutritionally sound dietary patterns without specific dietary guidance. Ritt et al. (24) have reported similar results.

This study was primarily limited by subject compliance. Individual motivation, sincerity, honesty and willingness to participate greatly influenced the outcome of the study. Because of the insignificant change in weight by subjects in the diet therapy groups and the poor adherence to dietary regimens, daily energy intakes may have been higher than mean values indicate.

Serum cholesterol concentrations in population studies have been correlated with the proportion of energy derived from fat (71). Saturated fats appear to be the dietary component responsible for this relationship (72). Nutrient intake in relation to serum triglyceride and cholesterol levels have been investigated by Nichols et al. (73). Twenty-four hour dietary recall interviews were conducted among 2,000 men and women in the community of Tecumseh, Michigan. Serum lipid levels were unrelated to quantity, quality or proportion of fat, carbohydrate or protein consumed. Similarly, in the present study, no correlations were found between serum lipid levels and carbohydrate, fat, protein or cholesterol intakes.

## SUMMARY AND CONCLUSIONS

The present study was conducted to investigate the effects of diet therapy, behavior modification, and exercise on weight reduction, body composition, serum lipids, and blood pressure. Following are the four experimental groups: 1) behavior modification plus exercise (BM+EX), 2) behavior modification (BM), 3) diet therapy plus exercise (DT+EX), and 4) diet therapy (DT).

The success of the weight reduction treatments was measured by the attrition rate and the total weight loss for the 12-week period. Lean body weight loss was considered undesirable. The attrition rate for subjects in the two behavior modification groups was slightly less than the attrition rate for subjects in the two diet therapy groups. Subjects in the two diet therapy groups indicated fair to poor adherence to their respective diets. Subjects in the EM group lost a significant ( $R\leq.05$ ) amount of weight (-6.9 pounds) while weight loss was not significant for subjects in the other three groups (BM+EX: -4.0 pounds; DT+EX: +1.2 pounds; DT: -1.7 pounds).

A significant ( $\underline{P} \leq .05$ ) reduction in total body fat (-4.9 pounds), and percentage fat (-1.9 percent), as determined by water densiometry, was noted for subjects in the EM group. Total body weight was correlated with total body fat (r=.96) and percentage fat (r=.89). No significant changes in lean body weight occurred for subjects in any of the four experimental groups. A 12-week follow-up evaluation indicated that weight and body composition parameters were not significantly changed from post to follow-up evaluation.

Mean urinary creatinine excretion values for subjects in the EM+EX group decreased significantly (P $\leq$ .05) from pre to post treatment evaluation (-.27 g/24 hr), and the change in urinary creatinine excretion values for members of this group was significantly ( $P\leq.05$ ) different from the change in urinary creatinine excretion values for subjects in the DT+EX group (+.14 g/24 hr). In general, mean values for all groups were within normal limits. Lean body weight was slightly correlated ( $r_{pre}$ =.58,  $r_{post}$ =.53) with urinary creatinine excretion.

Subjects in the BM+EX group significantly ( $P\leq.05$ ) decreased serum cholesterol levels (-17.7 mg/dl) from pre to post treatment evaluation. A significant ( $P\leq.05$ ) decrease in serum triglyceride was noted for subjects in the DT group. Low density lipoprotein and high density lipoprotein (HDL) serum lipids were not significantly changed for subjects in any of the four groups, however, the HDL/cholesterol ratio was significantly ( $P\leq.05$ ) increased for subjects in the two exercise groups (BM+EX: +.03; DT+EX: +.02).

No significant changes were found in either systolic or diastolic blood pressures for subjects in any of the experimental groups. Diastolic blood pressure was correlated (r=.48) with total body weight.

Selected blood serum constituents were within normal limits for all subjects.

Average energy intakes were not significantly different among subjects in the four experimental groups, however, subjects in the EM group consumed the least amount of energy (1,165 kcal) and lost the most weight (-6.9 pounds). Twenty percent of the total energy consumed by subjects in all four groups was derived from protein, 30 percent from fat, and 50 percent from carbohydrate. Subjects in all experimental groups met two-thirds or more of the Recommended Dietary Allowances for all nutrients but iron. No correlation was found between serum lipid levels and carbohydrate, fat, protein, or cholesterol intake.

Based on the results of the present investigation, the following conclusions have been made. Motivated individuals, utilizing a balanced food-exchange diet for weight reduction, can lose weight and maintain good nutritional status, however, motivation is not easily maintained in this kind of program. In contrast, motivation is more easily developed in a behavior modification program which attempts to change the consequences of an individual's behavior by reducing the number of antecedents that lead to inappropriate eating behavior. Secondly, daily physical activity representing a 250 kcal energy expenditure may reduce serum cholesterol and increase the high density lipoprotein/cholesterol ratio.

#### BIBLIOGRAPHY

- U.S. Public Health Service, Division of Chronic Diseases. Obesity and Health; A source book of current information for professional health personnel. Public Health Service Publication No. 1485. Washington D.C., U.S. Government Printing Office, 1966.
- Society of Actuaries. Build and blood pressure study. Vol. I and II. Chicago: Society of Actuaries, 1959.
- Williams, S.R. Nutrition and diet therapy. St. Louis: C.V. Mosby Co., 1973, p. 484.
- Select Committee on Nutrition and Human Needs, U.S. Senate. Dietary goals for the United States. U.S. Government Printing Office, Washington, D.C. Feb., 1977.
- Stunkard, A. New therapies for the eating disorders. Arch:.Genz.Psychiat:, 26:391, 1972.
- Stuart, R.B. Behavioral control of overeating. Behav. Research Ther. 5: 357, 1967.
- Gordon, T., W.P. Castelli, M.C. Hjortland, W.B. Kannel and T.R. Dawber. HDL as a protective factor against CHD: The Framingham Study. Am. J. of Med. 62:707, 1977.
- Balart, L., M.C. Moore, L. Gremillion, and A. Lopez. Serum lipids, dietary intakes, and physical exercise in medical students. J. Am. Dietet. Assoc. 64:42, 1974.
- Kannel, W.B., T.R. Dawber, G.D. Friedman, W.E. Cannon and P.M. McNamara. Risk factors in coronary heart disease. An evaluation of several serum lipids as predictors of CHD. The Framingham Study. Ann. Inter. Med. 61:888, 1964.
- Exchange Lists for Meal Planning, American Diabetic Association, Inc., and the American Dietetic Association, 1976.
- Ferguson, J.M. Habits not diets: The real way to weight control. Palo Alto, California: Bull Publishing Co., 1976.
- Brozeck, J.F. Grande, J.T. Anderson and A. Keys. Densiometric analysis of body composition: revision of some quantitive assumptions. Ann. N.Y. Acad. Sci. 110:113, 1963.
- 13. Miller, D.S., and S. Parsonage. Resistance to slimming. Lancet. 72:773, 1975.
- Stefanik, F.A., F.P. Heald, Jr. and J. Mayer. Calorie intake in relation to energy output of obese and non-obese adolescent boys. Am. J. Clin. Nutr. 7:55, 1959.

- Bullen, B.A., R.B. Read and J. Mayer. Physical activity of obese and nonobese girls appraised by motion picture sampling. Am. J. Clin. Nutr. 14:211, 1964.
- Mayer, J. Overweight. Causes, cost, control. Englewood Cliffs, N.J. Prentice-Hall, 1968.
- Stunkard, A.J. and M. McLaren-Hume. The results of treatment for obesity: a review of the literature and report of a series. Arch. Internal Med. 103:79, 1959.
- Shipman, W.G., and M.R. Plesset. Anxiety and depression in obese dieters. Arch. Gen. Psychiatry. 8:530, 1963.
- Stuart, R.B., and B. Davis. Slim chance in a fat world: Behavioral control of obesity. Champaign, Illinois: Research Press, 1972.
- Levitz, L.S. Behavior therapy in treating obesity. J.Am. Dietet. Assoc. 62:22, 1973.
- Ferguson, J.M. Dietitians as behavior-change agents. J. Am. Dietet. Assoc. 73:231, 1978.
- Ferster, C.B., J.S. Nurberger and E.B. Levitt. The control of eating. J. Mathetics. 1:87, 1962.
- Brightwell, D.R. Treating obesity with behavior modification. Postgrad. Med. 55:52, 1974.
- Ritt, R.S., H.A. Jordan and L.S. Levitz. Changes in nutrient intake during a behavioral weight control program. J. Am. Dietet. Assoc. 74:325, 1979.
- Zuti, W.B. and L.A. Golding. Effects of diet and exercise on body composition of adult females during weight reduction. Med. and Sci. in Sports. 5:62, 1973.
- Keys, A. Body composition and its change with age and diet. In: Weight Control, edited by E.S. Eppright, R. Swanson, and C.A. Inverson. Ames: Iowa State College Press, 1955.
- Hoberman, H.D., E.A.H. Sims and J.H. Peters. Creatine and creatinine metabolism in the normal male adult studied with the aid of isotopic nitrogen. J. Biol. Chem. 172:45, 1948.
- Borsook, H., and J.W. Dubnoff. The hydrolysis of phosphocreatine and the origin of urinary creatinine. J. Biol. Chem. 168:493, 1947.
- Forbes, G.B., and G.J. Bruining. Urinary creatinine excretion and lean body mass. Am. J. Clin. Nutr. 29:1359, 1976.
- Boileau, R.A., D.H. Horstman, E.R. Buskirk and J. Mendez. The usefulness of urinary creatinine excretion in estimating body composition. Med. and Sci. in Sports. 4:85, 1972.

- Miller, A.T., and C.S. Blyth. Estimation of lean body mass and body fat from basal oxygen consumption and creatinine expiration. J. of App. Physiol. 5:73, 1972.
- Best, W.R., W.J. Kuhl, Jr. and C.F. Consolazio. Comparison of physical and metabolic estimates of active protoplasmic mass. Fed. Proc. 12:178, 1953.
- Roessler, G.S., and B.G. Dunavant. Comparative evaluation of a wholebody counter potassium 40 method for measuring lean body mass. Amer J. Clin. Nutr. 30:1171, 1967.
- Muldowney, F.P., J. Crooks and M.M. Bluhm. The relationship of total exchangeable potassium and chloride to lean body mass, red cell mass, and creatinine excretion in man. J. Clin. Invest. 46:1375, 1957.
- Bailey, R.R., and H.E. DeWardener. Creatinine excretion. Lancet. 1:145, 1970.
- Bleiler, R.E., and H.R. Scheal. Creatinine excretion: Variability and its relationships to diet and body size. J. Lab. Clin. Med. 59:945, 1962.
- Scott, P.J. and P.J. Hurley. Demonstration of individual variation in constancy of 24-hour urinary excretion. Clin. Chem. Acta. 21:411, 1968.
- McGill, H.C., Jr., and J.P. Strong. The geographic pathology of atherosclerosis. Ann. N.Y. Acad. Sci. 149:923, 1968.
- Carlson, L.A., and L.E. Bottiger. Ischemic heart disease in relation to fasting values of plasma triglyceride and cholesterol. Stockholm Prospective Study. Lancet. 1:865, 1972.
- Kagan, A., T.R. Dawber, W.B. Kannel and N. Revotskie. The Framingham Study: A prospective study of coronary heart disease. Fed. Proc. 21:52, 1962.
- Truett, J.J. Cornfield and W.B. Kannel. A multivariate analysis of the risk of coronary heart disease in Framingham. J. Chron. Dis. 20:511, 1967.
- Rosenman, R.H., M. Friedman, R. Straus, C.D. Jenkins, S.J. Zyzanski and M. Wurm. Coronary heart disease in the western collaborative group study. J. Chronic Diseases. 23:173, 1970.
- Carlson, L.A. Serum lipids in men with myocardial infarction. Acta. Med. Scand. 167:399, 1960.
- 44. Albrink, M.J., J.W. Meigs and E.B. Man. Serum lipids, hypertension and coronary artery disease. Amer. J. Med. 31:4, 1961.
- Miller, G.T., and N.E. Miller. Plasma-high-density lipoprotein and development of ischemic heart disease. Lancet. 1:16, 1975.

- 46. Castelli, W.P., J.T. Doyle, T. Gordon, C.G. Hames, M.E. Hjortland, S.B. Hulley, A. Kagan and W.J. Zukel. HDL cholesterol and other lipids in coronary heart disease: The Cooperative Lipoprotein Phenotyping Study. Circulation 55:767, 1977.
- Berg, K., A. Borresen and G. Dahlen. Serum HDL and atherosclerotic HDL. Lancet. 1:499, 1976.
- Gordon, T., W.P. Castelli, M.C. Hjortland, W.B. Kannel and T.R. Dawber. Diabetes, blood lipids, and the role of obesity in coronary heart disease risk for women: The Framingham Study. Ann. Internal Med. 87:393, 1977.
- Glomset, J.A. The plasma lecithin-cholesterol acyl transferase reaction. J. Lipid Res. 9:155, 1968.
- Lehtonen, A. and J. Viikari. Serum triglycerides and cholesterol and serum high-density lipoprotein cholesterol in highly physically active men. Acta. Med. Scand. 204:111, 1978.
- Holloszy, J.D., J.S. Skinner, G. Toro and T.K. Cureton. The effects of a six month program of endurance exercise on serum lipids of middle-aged men. Am. J. Cardiol. 14:753, 1964.
- Kilbom, A., L. Hartley, B. Saltin, J. Bjure, G. Grimby and I. Astrand. Physical training in sedentary middle-aged and older men. Scand. J. Clin. Lab. Invest. 24:315, 1969.
- Lewis, S., W.L. Haskell, P.D. Wood, N. Manoogian, J.E. Bailey and M. Pereira. Effects of physical activity on weight reduction in obese middleaged women. Am. J. Clin. Nutr. 29:151, 1976.
- Goode, R.C., J.B. Firstbrook and R.J. Shephard. Effects of exercise and a cholesterol-free diet on human serum lipids. Can. J. of Physiol. and Pharm. 44:575, 1966.
- Garcia-Palmieri, M.R., R.Costas, Jr., J. Schiffman, A.A. Colon, R. Torres and E. Kazario. Interrelationship of serum lipids with relative weight, blood glucose, and physical activity. Circulation 45:829, 1972.
- Hurter, R., M.A. Peyman, J. Suale and C.W.H. Barnett. Some immediate and long term effects of exercise on the plasma-lipids. Lancet. 2:671, 1972.
- Wood, P.D., W. Haskell, H. Klein, S. Lewis, M.P. Stern and J. Farquhar. The distribution of plasma lipoproteins in middle-age male runners. Metabolism. 25:1249, 1976.
- Rony, H.R., and T.T. Ching. Studies on fat metabolism: effect of certain hormones on fat transport. Endocrinology. 14:355, 1930.
- Randle, P.J., P.B. Garland, C.N. Hales and E.A. Neusholme. The glucose fatty acid cycle. Its role in insulin insensitivity and the metabolic disturbances of diabetes mellitus. Lancet. 1:785, 1963.

- Chapman, C.B., and J.H. Mitchell. The physiology of exercise. Sci. Am. 212:88, 1965.
- 61 Enger, S.C., K. Herjornsen, J. Erikssen and A. Fretland. High density lipoproteins and physical activity: the influence of physical exercise age & smoking on HDL-cholesterol and the HDL/total cholesterol ratio. Scand. J. Clin. Lab. Invest. 37:251, 1977.
- Lopez, S.A., R. Vial, L. Balart and G. Arroyave. Effects of exercise and physical fitness on serum lipids and lipoproteins. Artherosclerosis. 20:1, 1974.
- Miller, G.J., N.E. Miller and M.T. Ashcroft. Inverse relationship in Jamaica between plasma HDL cholesterol concentration in coronary-disease risk as predicted by multiple risk factor status. Clin. Sci. and Mol. Med. 51, 475, 1976.
- Kannel, W.B., T. Gordon and W.P. Castelli. Obesity lipids and glucose intolerance. The Framingham Study. Am. J. Clin. Nutr. 32:1238, 1979.
- Walker, W.J. Relationship of adiposity to serum cholesterol levels in human subjects. Am. Intern. Med. 39:705, 1953.
- Berkowitz, D. Metabolic changes associated with obesity before and after weight reduction. J. Am. Intern. Med. 39:705, 1953.
- Montoye, J.H., H.L. Metzner, J.B. Keller, B.C. Johnson and F.H. Epstein. Habitual physical activity and blood pressure. Med. and Sci. in Sports. 4:175, 1972.
- Vitalle, F., Effects and benefits of exercise. In: Individualized Fitness Programs. Englewood Cliffs: Prentice Hall, Inc., 1973, p. 58.
- Wilcox, R.G. Serum lipid concentrations and blood pressure in obese women. Br. Med. J. 1:1513, 1978.
- Ashley, F.W. and W.B. Kannel. Relation of weight change to changes in atherogenic traits: The Framingham Study. J. Chron. Dis. 27:103, 1974.
- Connor, W.E., D.B. Stone and R.E. Hodges. The interrelated effects of dietary cholesterol and fat upon human serum lipid levels. J. Clin. Invest. 43:1691, 1964.
- Atherosclerosis Study Group. Primary prevention of the atherosclerotic diseases. Report of Inter-Society Commission for Heart Disease Resources. Circulation 42 (suppl. A):1-44, 1970.
- Nichols, A.B., C. Ravenscroft, D.E. Lamphiear and L.D. Ostrander, Jr. Daily nutritional intake and serum lipid levels. The Tecumseh Study. Am. J. Clin. Nutr. 29:1384, 1976.
- 74. Metropolitan Life Insurance Co. Public health report. 69:1085, 1954.

- Seiverd. Chemistry for medical technologists. White & Frankel, St. Louis, Mosby Co. 1965, pp. 121-124.
- Rudel, L.L., J.A. Lee, M.D. Morris and J.M. Felts. Characterization of plasma lipoproteins separated and purified by agarose - column chromatography. Biochem. J. 1398, 1974.
- Sardesai, V.M. and J.A. Manning. Determination of serum triglycerides. Clin. Chem. 14:156, 1968.
- Nobel, R.P. Electrophoretic separation of plasma lipoproteins in agarose gel. J. Lipid Research. 9:693, 1968.
- Lopes-Virella, M.F. Cholesterol determination in high-density lipoproteins separated by three methods. Clin. Chem. 23:882, 1977.
- Fox, S.M., J.P. Naughton and P.A. Gorman. Physical activity and cardiovascular health. The exercise prescription; frequency and type of activity. Mod. Concepts Cardiovas. Dis. 41:6, 1972.
- Snedecor, G.W. and W.G. Cochran. Statistical methods. 6th ed. Iowa State University Press, 1967.
- Harris, M.B. and E.S. Hallbauer. Self-directed weight control through eating and exercise. Behav. Res. & Therapy. 11:523, 1973.
- Srivastava, S.S., K.V. Mani, C.M. Soni and J. Bahti. Effect of muscular exercise on urinary excretion of creatinine. Ind. J. Med. Res. 55:953, 1967.
- Fredrickson, D.S. and R.I. Levy. Familial Hyperlipoproteinemia. In: The Metabolic Basis of Inherited Disease, edited by J.B. Stanburg, J.B. Wyngaarden and D.S. Fredrickson. (3rd ed.) McGraw-Hill, 1972, p. 551.
- Mann, G.V. The influence of obesity on health. New Engl. J. Med. 291:178, 1974.

APPENDIX

#### CONSENT FORM

have volunteered to

participate in an experiment designed to study the effectiveness of different methods of weight control. I have been asked to complete the study which will conclude Dec. 15, 1978, but I should feel free to withdraw at any time. A follow-up test will be done in March of 1979 if I can participate. I understand that I may refuse to undergo any of the testing procedures without prejudice.

Participation will require the following physical measurements and tests:

- (1) Body weight--taken initially and on a weekly basis
- (2) Body height -- taken initially
- (3) Skinfolds -- taken initially, at the midpoint, and end
- (4) Underwater weighing--taken initially, at the midpoint, and end. (Weighing will be done while in a swimming suit.)
- (5) Bone diameters and girth measurements will be taken initially
- (6) Blood pressure -- initial, midpoint, and final
- (7) Three day dietary food records will be obtained three times during the study
- (3) A 24-hour urine collection--taken initially, at the midpoint and final stages of testing
- (9) Fasting blood will be obtained by veinipuncture at the beginning and end of the study period. This will be done under the supervision of medical or allied health personnel.

All final testing will be done before the end of the 1978 school semester. A follow-up test will be planned for March 1979, using the same testing procedures as above.

I will be required to attend 12 weekly instructional and educational meetings, lasting from 30 to 60 minutes.

I will feel free to ask questions about testing procedures. All results will be kept strictly confidential. I will have an opportunity at the end of the study to find out the conclusions.

To my knowledge, I am in a good state of health. I am not pregnant and do not have any cardiovascular problem, infectious disease or metabolic condition which could limit my participation in the study. If pregnancy occurs, I will voluntarily withdraw from the study.

I have read and understand the above statement. I hereby voluntarily consent to participate.

Date \_\_\_\_\_

Signature

Witness

53

		. Department of	Junattan, Kansi	Nutrition ustin Hall
		P	hone: 913 :	532-5508
	BLOOD LIPID SU	RVEY		
Name	: Lo	cal Physician		
Telep	hone Number			
deigh	t			
Veigh	t			
ige _		Date Form Complete	d	
L. A	re you now on diet therapy for hyperli holesterol diet?	pidemia or a low	Yes	No
2. H	ave you ever been advised by your phys holesterol diet?	ician to eat a low	Yes _	No
. A	re you taking drugs for hyperlipidemia	?	Yes	No
	If yes indicate drug and dosage			
	a) clofibrate (Atromid-S)			
	b) thyroxine			
	c) nicotinic acid			
	d) cholestyramine			
	e) other			
. A:	re you taking oral contraceptives?		Yes	No
	If yes indicate type (name)			
. A:	re you taking any androgenic steroid?		Yes	No
	If yes indicate drug and dosage			
	a) fluoxymesterone			
	b) oxymethalone			
	c) norandrolone			
	d) testosterone			
	e) other			
. Do	o you smoke?		Yes	No
	If yes, please indicate			
	cigarettes packs	day		
	cigars			

pipe

				34
7.	Have you ever taken any medicine for heart trouble	Yes	No	-
	If yes, type			
	Do you take it now? Yes No			
8.	Have you ever been told that you have high blood pressure?	Yes	No	
	If yes, what was your age			
0	Num un solar and adde for bigh blood programs?	Vec	No	
9.	Have you ever taken any medicine for high blood pressure:	ies		-
	Ear how long			
	Do you take it now Yes No			
	Do you take it now ites no			
10.	Have you ever had a heart attack?	Yes	. No	-
	If yes, at what age?			
11.	Is your mother alive?	Yes	No	-
	If yes, a) age now			
	b) Has she ever been thought to have neart			
	Vec (decertibe)			
	If no a) cause of death?			
	b) are at death			
	b) age at death			
12.	Is your father_alive?	Yes	. No	-
	If yes, a) age now			
	b) Has he ever been thought to have heart disease?			
	Yes (describe)No			
	If no, a) cause of death?			
	b) age at death	•		
13.	How many brothers and sisters have you who are alive now?			
	Brothers? Ages?			
	Sisters? Ages ?			
14.	Have any of them ever had heart trouble?	Yes	No	_
	not app	licable		
	It yes, give details as follows	d Natur	e of T	llnes
	a)	- HELDI		
	b)			
	c)			
	d)			

15.	Have you ever had blood drawn for a test for cholesterol, triglycerides or lipoprotins (fats)?	Yes	No
	If yes, please record the known values		
	cholesterol date		
	triglycerides		
	lipoproteins		
16.	Have your children had blood drawn for a test of cholesterol, triglycerides or lipoproteins? Not known	Yes	No
	If yes, please record the known values		
	Son Daughter Age		
	cholesterol		
	triglyceride		
	lipoproteins		
17.	Has anyone in your family ever had diabetes (sugar disease)?	Yes	No
	If yes, which of your relatives (e.g. brother, cousin, parent)		
18.	If you are married, does your spouse suffer from		
	a) heart trouble Yes No		
	b) high blood pressure Yes No		
19.	What medications are you presently taking?		

Please do no write below this line.

NAME:			 	
DATE:		<u>.</u>	 	 
Serum chole	sterol mg/dl:		 	 
Serum trigl	yceride mg/dl:		 	 
Lipoprotein	electrophores	is:		
Check	one			
	Normal		 	 
	Type I		 	 
	Type II		 	 
	Type III		 	 
	Type IV			
	Type V		 	 
. Chylomi	crons present			
	Yes		 	 
	No		 	 

Other abnormality (specify)

# DIETARY RECORD

Name	Date	Day
Place eaten:	Dinner	Supper
Breakfast	Lunch	Dinner
Food	Estimated measure	Kind of ingredients
Breakfast		
Between Meals		
between neurs		
Lunch or Dinner		
Between Meals		
Dinner or Supper		
Rotucco Manla		
perween sears		

CALCULATION OF DAILY ENERGY REQUIREMENTS<sup>5</sup>

1. Energy expenditure for basal metabolic rate (BMR)

(A) (weight in kg) x l kcal = (B)

To find the daily expenditure for BMR, multiply (B) by 24 hours.

(B) x 24 hr = (c)

2. Energy expenditure for activity

Sedentary	20%	of	BMR
Light activity	30%	of	BMR
Moderate activity	40%	of	BMR
Heavy activity	50%	of	BMR

Select the appropriate activity leval and calculate the number of kcal you denote to activity

\_\_\_\_\_% of (C) \_\_\_\_ = (D) \_\_\_\_\_

3. Energy expenditure for specific dynamic action (SDA)

Estimation of SDA

10% of (C) \_\_\_\_\_+ (D) \_\_\_\_\_ = (E) \_\_\_\_\_

4. Total daily energy expenditure

(C) \_\_\_\_\_ + (D) \_\_\_\_\_ + (E) \_\_\_\_\_ = TOTAL \_\_\_\_\_

 $^{5}\!\text{Williams},$  S.R. Nutrition and Diet Therapy. (2nd ed.). St. Louis: C.V. Mosby Co., 1973, p. 484.

### 1.200 KILOCALORIE DAILY FOOD PLAN

6 meat exchanges 4 bread exchanges 2 milk exchanges 3 vegetable exchanges 3 fruit exchanges 3 fat exchanges

Breakfast:

Lunch:

Dinner:

Snack:

1 oz. toast 1 bread 1 tsp. margarine 1 fat 1 orange 1 fruit 1/3 C. cottage cheese 1 meat coffee

chicken sandwich: (2 slices bread 2 bread 2 oz. chicken 2 meat 1 tsp. mayonaise) 1 fat cucumber slices 1 vegetable 1 vegetable carrot sticks 3/4 C. plain yogurt 1 milk 1/2 C. strawberries 1/2 fruit ice tea

3 oz. brofled lean beef 3 meat baked potato 1 bread 1 T. sour cream 1 fat broccóli spears 1 vegetable apple sliced on 1 fruit lettuce leaf coffee

banana milk shake (1/4 banana 1 C. skim milk 1 milk artificial sweetner)

1/2 fruit

#### QUESTIONNAIRE EVALUATING SUBJECT COMPLIANCE TO AN ENERGY-RESTRICTED FOOD EXCHANGE DIET

- On a scale from 1 to 5, with 1 representing very good adherence and 5 representing very poor adherence, please indicate how closely you've been following your diet.
- When are you most likely to be tempted not to follow your diet? Please check the appropriate answer.

\_\_\_\_\_morning \_\_\_\_\_afternoon evening

- 3. Is it more difficult to follow your diet when you are alone or with others?
- Please list two of your favorite foods. Have you included them in your diet plan?

INDIVIDUAL SUBJECT VALUES

Sub ject	Group	Age	Height (in)	Weight (1b)	Percentage fat	Triglyceride <sup>a</sup>	Gholesterol <sup>a</sup>	HDL <sup>a</sup>	LDL <sup>a</sup>	HDL/cho1
9004	$^{1}$ p	42	58.5	176* 175**	41.8 40.2	36 30	150 146	53 52	90 88	.35 .36
2006	-	31	63 <b>.</b> 3	144 134	36.2 32.8	47 22	196 156	75 43	112 109	.38 .28
9008	1	47	65°0	156 163	38.0 40.6	86 68	170 162	45 54	108 94	•26
9010	1	34	64.0	167 165	37.8 37.0	193 88	238 155	50 40	125	.21 .26
9011	1	39	70.0	223 211	43.5 40.4	44 35	171 127	39 53	123 67	•23 •42
9013	1	34	65.4	142 139	30.4 29.9	98 54	179 180	44 42	115 127	• 25 • 23
9039	1	30	66.0	216	44 <b>.</b> 0		 166		106	.28
9043	1	31	0.69	231 222	42.8 41.7	89 84	260 253	47 48	195 188	.18 .19
9051	1	46	67.0	150 149	38.8 37.0	60 	191		139	.21
ase	erum lit	ofd m	- easurement und	lts = (mg/d1).	bl=behavior mo	odification plus	s exercise. *=	-pre ti	reatmer	ų

Sub ject	Group	Age	Height (in)	Weight (1b)	Percentage fat	Triglyceride <sup>a</sup>	Cholesterol <sup>a</sup>	HDL <sup>a</sup>	LDL <sup>a</sup>	HDL/chol
1006	<sup>2</sup> <sup>b</sup>	35	62.0	141* 142**	36.2 37.6	57 43	195 207	49 49	135 149	•25 •24
6006	2	22	63.0	128 121	38.3 35.1	30	192 208	63 65	123 136	•33 •31
9018	2	33	66 <b>.</b> 6	164 142	34 <b>.</b> 4 28.3	55 36 ·	163 198	46 31	106 160	•28 •16
9028	2	51	62.0	168 156	41.4 39.7	123	233	41 	167	.18
9034	2	20	61.5	140	35.6 	193	238	50	149	.21 
0706	2	42	63 <b>.</b> 9	151 151	46.2 45.3	119 77	194 193	75 79	95 98	.39
9047	2	31	64.0	135	29.5 28.8	45 39	177 169	40 45	128 116	• 23 • 27
9053	2	38	64.3	148 142	37.5	48 34	183 181	58 62	115 112	<b>.</b> 32 .34

<sup>a</sup> serum lipid measurement units = (mg/d1). <sup>b</sup> = behavior modification. <sup>A=</sup>pre treatment measurement. \*\*-post treatment measurement.

Sub ject	Group	Age	Height (in)	Weight (1b)	Percentage fat	Triglyceride <sup>a</sup>	Cholesterol <sup>a</sup>	HDL <sup>a</sup>	$LDL^{a}$	HDL/cho1
9026	3 b	43	60*0	132* 133**	 40,3	61 39	211 204	42 54	157 142	.20
9030	£	35	65.1	146 147	36.2 35.9	43 86	172 204	50 60	113 127	.29
9036	e	20	64.0	139	27.2 	31 54	141 153	39 45	96 97	.28 .29
9042	e	38	61.0	127 128	35.3 35.3	49 42	184 187	50 63	124 116	.27 .34
9048	e	35	66.5	142 144	33.7 35.4	85 	178	41	120	. 23
9052	e	21	63.0	142 		46 34	170 137	47 42	114 88	.28 .31
9054	en en	20	63 <b>.</b> 0			97 103	181 210	50 50	112 139	.28 .24

 $a^{a}_{Serum}$  lipid measurement units = (mg/dl),  $b_{3=diet}$  therapy plus exercise. \*= pre treatment measurement.

HDL/chol	.25	.22	.23	.19	.18 .19	.22 .24	.27 .26	.23	.19	
LDL <sup>a</sup>	115 148	152 147	157 121	166 187	169 184	147 133	133 124	93	113	
HDL <sup>a</sup>	43 43	46 48	54 55	43 48	43 50	48 50	54 47	35 32	31 36	
Cholesterol <sup>a</sup>	172 · 202	210 209	234 187	230 249	233 255	221 207	203 182	153 146	163 165	
Triglyceride <sup>a</sup>	72 55	62 68	116 54	103 69	103 104	128 122	78 55	126 93	94 92	
Percentage fat	ł, I	33 <b>.</b> 8 33 <b>.</b> 9		37.3 37.0	37.7 37.0	43.9 44.2	30.4 29.4	51.1	27.4	
Weight (1b)		144 146	180 177	144 144	166 161	154 154	162 157	206	132	
(in)										
Height	146	65.0	65.0	61.0	62.9	63.0	67.0	60.0	64.0	
Age	27	35	31	41	45	49	31	29	26	
Group	4 <sup>b</sup>	4	4	4	4	4	4	4	4	
Sub ject	9003	9006	9014	9023	9024	1406	9045	9046	9056	

\*\*=post <sup>a</sup>Serum lipid measurement units = (mg/d1). <sup>b</sup>4=diet therapy. <sup>\*=</sup>pre treatment measurement. treatment measurement.

THE EFFECTS OF DIET THERAPY, BEHAVIOR MODIFICATION, AND EXERCISE ON WEIGHT REDUCTION AND SERUM LIPIDS

Ъy

Joyce Petitjean Oldenburg

B. S., Kansas State University, 1977 M. S., Kansas State University, 1979

.

AN ABSTRACT OF A MASTER'S THESIS

submitted in partial fulfilment of the

requirements for the degree

MASTER OF SCIENCE

Department of Foods and Nutrition

KANSAS STATE UNIVERSITY Manhattan, Kansas

Thirty-three premenopausal, non-pregnant, obese (mean = 38% fat) women aged 20 to 52 years were participants in a 12-week study designed to investigate the effectiveness of two methods of weight reduction and the effects of diet therapy, behavior modification, and exercise on weight reduction, body composition, urinary creatinine excretion, blood pressure, serum lipids, and nutritional status. Women were assigned randomnly to one of two groups: 1) behavior modification(BM) or 2) diet therapy (DT). These two groups were subdivided into exercise and non-exercise groups. Subjects who were presently participating in some form of physical activity were assigned to one of the exercise groups. Women in the behavior modification groups were to lose weight by developing appropriate eating patterns rather than by following a special, restrictive diet. Women in the diet therapy groups were to lose approximately one pound, weekly, by following their individualized energy-restricted diets. Exercise consisted of increased physical activity (equivalent of about 250 kcal energy expenditure). Subjects were tested before and after the 12-week experimental period. A follow-up study was conducted 12 weeks later.

The attrition rate for subjects in the two behavior modification groups was slightly less than for subjects in the diet therapy groups (41% versus 45%). Only subjects in the EM group lost a significant ( $P\leq.05$ ) amount of weight (-6.9 pounds), total body fat (-4.9 pounds), and percentage fat (-1.9%) between pre and post treatment evaluation. Total body weight was correlated with total body fat (r=.96) and percentage fat (r=.89). No significant changes in lean body weight occurred for subjects in any of the four groups. A 12-week follow-up study indicated that body composition did not change significantly frum post treatment to follow-up investigation. Mean urinary creatining values for subjects in the EM+EX group decreased significantly ( $P\leq.05$ )

and the changes in urinary creatinine excretion (UCE) for members of this group were significantly (P<.05) different from the changes in UCE for subjects in the DT+EX group. Lean body weight was slightly correlated (rpre=.58; rpost=.53) with UCE. Subjects in the BM+EX group significantly (P4.05) decreased serum cholesterol levels by 17.7 mg/dl. A significant (P<.05) decrease in serum triglyceride (-19.0 mg/dl) was noted for subjects in the DT group. Low density lipoprotein and high density lipoprotein (HDL) were not significantly changed from pre to post treatment evaluation, but the HDL/cholesterol ratio was significatnly (P<.05) increased for subjects in the two exercise groups. No significant changes were found in systolic or diastolic blood pressure. Diastolic blood pressure was slightly correlated (r=.48) with total body weight. Mean energy intakes were not significantly different among subjects in the four groups, however, subjects in the BM group consumed the least amount of energy (1,165 kcal) and lost the most weight (-6.9 pounds). Subjects met two-thirds of the Recommended Dietary Allowances for all nutrients but iron. Serum blood lipids were not correlated with fat. carbohydrate, or dietary cholesterol intake. Subjects in the DT group indicated fair to poor adherence to their respective diets.

The results of the present study suggest that motivation is more easily developed in a behavior modification program than an energy-restricted dietary program. Increased daily physical activity (equivalent of 250 kcal energy expenditure) may reduce cholesterol levels and increase the HDL/cholesterol ratio.