



OPEN

Selection of Reference Genes for Expression Analysis in *Diuraphis noxia* (Hemiptera: Aphididae) Fed on Resistant and Susceptible Wheat Plants

Deepak K. Sinha & C. Michael Smith

Department of Entomology, Kansas State University, Manhattan, KS 66506-4004, USA.

The Russian wheat aphid (RWA), *Diuraphis noxia* Kurdjumov, is a major global pest of wheat and barley production that causes enormous economic damage. Few studies have been conducted to explore and decipher the molecular basis of RWA strategies to evade plant defense mechanisms. Gene expression studies of RWA in response to wheat genotypes carrying different RWA resistance genes have been initiated in our group; however, a secure and accurate understanding of RWA gene expression is dependent on identification of suitable reference genes. This study analyzed expression profiles of five potential reference genes selected and sequenced during RNA sequencing experiments. The expression of genes coding for actin and ribosomal protein L27 was comparatively less variable in RWA fed on different wheat hosts. Results of geNorm, NormFinder, and BestKeeper expression analyses support the use of actin and ribosomal protein L27 in RT-qPCR studies of RWA gene expression in studies involving RWA-wheat interactions.

The advent of next-generation sequencing technologies has resulted in a significant increase in transcriptomic data for various organisms¹. The generated transcriptomes have helped researchers not only to decipher expression pattern of genes and transcripts but also define the genetic architecture of many species^{2,3}. Validation of gene expression from such transcriptomic resources has become mandatory for reporting and reconfirming expression profiles⁴, and reverse transcription quantitative PCR (RT-qPCR) is rapidly replacing traditional methods such as Northern blotting and Ribonuclease Protection Assays (RPA)^{5,6}. RT-qPCR's speed, sensitivity, efficiency and reproducibility has made it the gold standard for rapid and accurate quantification of gene expression profiles⁷ from various next-generation sequencing (NGS) datasets. Commercially available instruments and consumables have further led to universal acceptance of RT-qPCR⁶.

Gene expression using RT-qPCR assay is based on the principle of quantifying target mRNA during the exponential phase of the PCR. During this phase, the target is doubled with each PCR cycle and the probability of PCR-bias due to limited reagents is nullified or decreased. In RT-qPCR, the amplification of product is detected on accumulation of fluorescent signal. The cycle at which the fluorescent signal exceeds background signals is referred to as the threshold cycle, or Ct (also referred to as the quantification cycle, or Cq). Analysis of Cq is used to estimate expression of the respective genes. Several factors, such as RNA quality and quantity, mastermix components used in the PCR and efficiency of PCR reaction, influence Cq values⁸. Absolute and relative quantification methods are generally used to estimate gene expression using RT-qPCR. The absolute quantification approach necessitates generating the standard curve of known copy numbers of each target, which in turn requires standard curves for multiple targets in a study and knowledge of copy numbers of each target, thus limiting its usage. The most widely adopted approach, relative quantification, is based on estimation of gene expression normalized to the expression of a control gene known as a reference; therefore, reliable determination of reference/references is the central factor in accuracy of this method.

A gene can be used as a reference when it is highly and stably expressed in all samples under investigation and is not co-regulated with a target gene^{9,10}. Reference genes traditionally have been housekeeping genes believed to possess inherent stable and constitutive expression irrespective of physiological conditions in different samples or treatments under investigation^{11,12}. The universal stability of housekeeping gene expression was disproven in recent years^{13,14}, negating their use¹⁵. According to the Minimum Information for Publication of Quantitative

SUBJECT AREAS:

MOLECULAR BIOLOGY

ZOOLOGY

GENE EXPRESSION ANALYSIS

Received
14 March 2014Accepted
7 May 2014Published
27 May 2014

Correspondence and requests for materials should be addressed to C.M.S. (cmsmith@ksu.edu)

Table 1 | Primers designed for amplification and sequencing of selected *D. noxia* genes

Primer	Maximum homology (E-value)	Sequence (5'-3')	Annealing temperature (°C)	Amplicon size in cDNA (bp)
Actin	<i>Acyrtosiphon pisum</i> actin-LOC100145822 (0)	F:TGCCTGATGGTCAAGTCATC R:TCAAGGAATGCTTTGAGCTG	55	553
RPL27	<i>Acyrtosiphon pisum</i> ribosomal protein L27 (0)	F:GAACCTTAACCTTCTGGAAGAACCA R:GCTCCGTCGTAACCTCAA	55	519
RPL9	<i>Acyrtosiphon pisum</i> 39S ribosomal protein L9-LOC100159528 (4e-114)	F:TCGAGGTAATCGATCTGCAA R:GCATGAATAAAGATGAGCCTTG	55	245
RPL5	<i>Acyrtosiphon pisum</i> 60S ribosomal protein L5-LOC100162441 (0)	F:AGCGTTGGTCAATTTCTTGG R:TGAAGGTCCAAATGCTTTCC	58	448
TEF2	<i>Acyrtosiphon pisum</i> elongation factor 2-LOC100168198 (2e-86)	F:CAAGGTATGAATTGATGTCTGGT R:TCGTTAAAGCATACTTGCTGT	58	220

Real-Time PCR Experiments (MIQE) guidelines¹⁶, reference gene(s) are now selected based on their specificity in interactions between a species or cell type/s subjected to different treatments or conditions.

The Russian wheat aphid (RWA), *Diuraphis noxia* Kurdjumov, originally from Central Asia, was introduced globally in the 1900s¹⁷ and has become a major destructive pest of wheat and barley and caused huge economic losses. Host-plant resistance is the most acceptable and ecologically beneficial aphid pest management strategy in wheat growing regions, yet host-plant resistance is transitory in many cases, because virulent populations develop to survive in once cultivated resistant varieties¹⁸. The short-term solution to this problem traditionally has been to identify new sources of resistance and breed new RWA-resistant wheat varieties, but new technologies such as NGS and RT-qPCR offer the possibility of understanding the molecular principles supporting RWA virulence and engineering resistant plants with greater longevity. Our research is focused on investigating the molecular mechanisms of RWA-plant interaction with the long-term goal of finding novel and durable solutions to RWA management.

The objective of this investigation was to identify a robust RWA reference gene for use in validation of gene expression studies in RWA-wheat interactions. We identified four previously unreported RWA sequences commonly used as reference controls in other biological systems and report the most suitable reference controls for RT-qPCR assays of RWA genes expressed in aphid-wheat interactions.

Results and Discussion

Analysis of RNA quality, sequencing and sequence analysis. Bioanalyzer analysis of RNA revealed a ribosomal shift due to denaturation of 28 s rRNA (see Supplementary Fig. 1 online), consistent with previously published examples of insect RNA¹⁹. Multiple peaks were absent, indicating no RNA degradation. Thus, screening total RNA using NanoDrop spectrophotometry, gel electrophoresis and capillary electrophoresis produced high-quality RNA for RNAseq library preparation. The libraries analyzed using

the Bioanalyzer revealed fragment sizes of >230 bp. The quality of the 18 libraries (6 treatments × 3 replications) is shown in Supplementary Fig. 2 online. Libraries were sequenced and filtered to remove low-quality sequences, providing >10 million reads for each library. Results of BLAST analysis and differential gene/transcript analysis will be reported separately.

Gene selection and RT-qPCR. The commonly used reference genes actin, ribosomal protein L9, ribosomal protein L27, transcription elongation factor 2 and ribosomal protein L5 were selected based on a literature survey. Actin has been used as a reference gene in many studies of model insects such as the fruit fly, *Drosophila melanogaster* Meigen²⁰. Transcription elongation factor genes were evaluated to check their suitability as reference genes in several insects, particularly Hemipterans such as the sweetpotato whitefly, *Bemisia tabaci* (Gennadius)²¹ and brown plant hopper, *Nilaparvata lugens* (Stål)²². Genes coding for ribosomal proteins have been reported as most reliable reference in insects such as *N. lugens*²². Many transcripts coded for these genes were identified from RNAseq data, revealing >93% homology to the respective genes in the pea aphid, *Acyrtosiphon pisum* (Harris) (Table 1). Representations of these genes observed in RNAseq data confirmed that there was little or no change in their expression. These genes were amplified from cDNAs, yielding amplicon sizes ranging from 220 bp to 553 bp (Table 1). These sequences have been deposited in DNA Databank of Japan with accession numbers AB914563-AB914566.

Amplicon size and melting temperature of RT-qPCR primers designed from these sequences are shown Table 2. Primer efficiency ranged from 91.50 to 102.65, with corresponding R² values from 0.9922 to 0.9995 (Table 2). Primers were redesigned and efficiencies rechecked in cases where the efficiency or R² values differed from the range of recommended values. Melt curve analysis of all genes showed no primer dimers or nonspecific product amplification.

Analysis using geNorm software. Cq values derived from RT-qPCR assay for expression stability were logarithmically transformed as

Table 2 | Primers designed for amplification and RT-qPCR of selected genes from *D. noxia* feeding on different wheat genotypes

Primer	Sequence (5'---3')	Tm (°C)	Amplicon size in cDNA (bp)	RT-qPCR efficiency (%)	Correlation coefficient (R ²)
RTActin	F:TGCCTGATGGTCAAGTCATC R:TCAAGGAATGCTTTGAGCTG	55	110	91.99	0.9992
RTRPL27	F:ACCAGCACGATTTTACCAGATTC R:CGTAGCCTGCCCTCGTGTA	55	90	91.50	0.9995
RTRPL9	F:CATTCCACTGGCACITTTTCTTT R:TGCCAGAACGCCCAATTAGT	55	100	92.60	0.9993
RTRPL5	F:AACCCAAAATGTGTTTACGATGAG R:AGCTGTCGATGGAGGTTTCAAC	55	110	102.65	0.9922
RTTEF2	F:CACATGGGTACCTGGGAATA R:TCGGATTCACTGCTGATTGC	55	96	93.07	0.9993




Figure 1 | The average expression stability values (M) derived using geNorm software for the candidate genes actin, ribosomal protein L5 ($RPL5$), ribosomal protein L9 ($RPL9$), translation elongation factor 2 ($TEF2$) and ribosomal protein L27 ($RPL27$). The least stable candidate gene is plotted on the left, and the most stable gene is on the right.

input for geNorm analysis. The principle behind this algorithm is that if two genes are stably expressed in a sample set then the ratio of their logarithmic transformed expression should be constant; thus, geNorm ranks genes based on their average expression stability (M), and the candidate gene possessing the lowest M value is the most stably expressed gene in that set. Figure 1 depicts the average expression stability of actin ($M = 0.210$) and ribosomal protein L27 ($M = 0.267$), the two most stable genes that should be used as references in RT-qPCR assays involving *D. noxia*-wheat interactions.

geNorm also predicts an optimal number of reference genes for accurate representation of gene expression based on calculation of pairwise variation (V_n/V_{n+1}) between sequential normalization factors (NF_n and NF_{n+1}). A cutoff value of 0.15 is considered for the ratio, below which there is no requirement of any other reference gene¹³. Large pairwise variation represents a significant effect in gene expression due to the addition of another gene or genes, reinforcing the need for the second gene to be included to derive reliable




Figure 2 | geNorm pairwise variation (V) analysis to determine optimal number of reference genes for normalization in RT-qPCR reaction.

normalization factors. Candidate genes are added based on the ranking derived using the M value. This analysis, depicted in Figure 2 for our experiments, clearly showed that adding a third gene does not increase the ratio by more than 0.15; therefore, actin and ribosomal protein L27 represent the required reference genes for accurate estimation of *D. noxia* gene expression in aphid-wheat interactions.

Analysis by NormFinder software. NormFinder analyzes candidate reference genes according to inter- and intra-group variation in expression. Similar results were obtained using NormFinder, which predicted stability value of 0.735 for ribosomal protein L27 followed by stability value of 0.738 for actin (Figure 3). Therefore, both geNorm and NormFinder outputs provide proof for stable ribosomal protein L27 gene that can be used as a reference in all the RT-qPCR assays that studies expression patterns of genes in




Figure 3 | NormFinder stability index for actin, ribosomal protein L5 ($RPL5$), ribosomal protein L9 ($RPL9$), translation elongation factor 2 ($TEF2$), and ribosomal protein L27 ($RPL27$). Genes with the least stable expression are on the left, and the most stable genes are on the right.



Table 3 | Descriptive statistics based on crossing point (CP) data and expression stability of five *D. noxia* reference genes calculated by BestKeeper Software

CP factor	Actin	Ribosomal protein			Transcription elongation factor 2
		L27	L9	L5	
Mean Geo CP	17.96	18.77	27.38	23.85	19.28
Mean Ar CP	17.96	18.78	27.65	24.66	19.33
min CP	17.28	18.07	23.06	17.66	17.18
max CP	18.87	19.54	33.30	32.95	21.82
CP \pm SD	0.51	0.47	3.72	6.24	0.97
CP CV (%)	2.85	2.48	13.47	25.29	5.00
CP CV (%) \pm SD	2.85 \pm 0.51	2.48 \pm 0.47	13.47 \pm 3.72	5 \pm 1.95	25.29 \pm 6.24

Abbreviations. CP – Crossing point, Geo – geometric, Ar – arithmetic, SD – standard deviation, CV – percentage coefficient of variation.
n = 18 samples for each gene.

the Russian wheat aphid feeding on different wheat varieties containing aphid resistance genes.

BestKeeper analysis software. BestKeeper considers the Cq values of all candidate reference genes, in order to calculate standard deviation (SD) and coefficient of variation (CV). The software excludes genes with Cq value SDs greater than 1. Results of geNorm and NormFinder analyses were further verified when Cq data were analysed using BestKeeper software, which predicts gene stability based on low CV and SD. Ribosomal protein L27 (CV = 2.48 \pm 0.47%) and actin (2.85 \pm 0.51%) were found to be the most stable (Table 3).

Validation of reference genes. Results of the current study indicate actin and ribosomal L27 to be the most appropriate reference genes in RT-qPCR assays involving RWA-wheat interactions. To validate their use, we assessed expression of tRNA-Leu, which we previously reported to be up-regulated in the gut transcriptome of RWA2 fed *Dn4* plants²³. Results of this experiment revealed significant ($p < 0.001$) over-expression of tRNA-Leu in RWA2/*Dn4* interactions in comparison to other interactions (Figure 4), and very stable CV- and M values (Mean CV = 0.168, Mean M = 0.48) for the actin - ribosomal L27 combination. In contrast, use of the actin - RPL5 combination (Mean CV = 0.97, Mean M = 4.2) or the RPL27 - RPL5 combination (Mean CV = 0.952, Mean M = 3.95) - resulted in huge changes in expression, and Mean CV- and M values above acceptable

stability values [homogenous (CV < 0.25; M < 0.5) and heterogeneous (CV < 0.5; M < 1)]. Thus, the results of the tRNA-Leu expression validation experiment confirm the use of actin and RPL27 for RWA expression analysis.

Conclusions. This study emphasizes the use of an appropriate reference gene(s) for RT-qPCR studies, reports sequences of five selected genes from the Russian wheat aphid, and identifies reference genes with stable expression in RWA feeding on wheat plants carrying different RWA resistance genes. Analyses of our data using three different statistical techniques indicated that RWA ribosomal protein L27 (*RPL27*) is the most stable reference gene, closely followed by actin. Ideally, both *RPL27* and actin genes should be used as references for normalizing expression profiles in RT-qPCR studies. This study is an important step in our ongoing RNAseq studies to define differential expression of RWA genes in RWA-wheat interactions. Results of these experiments will be of utmost importance to RWA genome sequencing efforts as well as the huge amount of expression data derived from current studies using NGS technologies.

Methods

Insect and plant material. RWA1 (biotype 1) collected from wheat fields near Hays, KS, and RWA2 (biotype 2) individuals collected from wheat fields near Briggsdale, CO (via the USDA-ARS Plant Science Research Laboratory at Stillwater, OK), were cultured in separate housing in the greenhouse at Kansas State University on plants of susceptible wheat cultivar 'Jagger' for use in all experiments. The identity of each




Figure 4 | Relative normalized expression of the t-RNA-Leu gene using reference gene combinations of: (a) actin and ribosomal protein L27 (RPL27), (b) actin and ribosomal protein L5 (RPL5) and (c) RPL27 and RPL5. R1D0 = RWA1 fed wheat plants carrying the *Dn0* (no resistance) gene; R1D4 = RWA1 fed wheat plants carrying the *Dn4* resistance gene; R1D7 = RWA1 fed wheat plants carrying the *Dn7* resistance gene; R2D0 = RWA2 fed wheat plants carrying the *Dn0* (no resistance) gene; R2D4 = RWA2 fed wheat plants carrying the *Dn4* resistance gene; R2D7 = RWA2 fed wheat plants carrying the *Dn7* resistance gene.



biotype was verified independently in diagnostic plant differential greenhouse assays at Stillwater and Manhattan. Voucher specimen no. 176 (RWA2) is deposited with the Kansas State University Museum of Entomological and Prairie Arthropod Research.

RWA1 and RWA2 adults were starved for 12 h, then fed on wheat plants containing RWA-resistant genes *Dn4* (Yumar), *Dn7* (94M370), and the susceptible control Yuma (*Dn0*) in a fine screen mesh cage in the greenhouse. All plants were grown in the greenhouse in 16.5-cm-diameter-plastic pots containing Pro-Mix-Bx potting mix (Premier ProMix, Lansing, MI). Environmental conditions were 24:20°C day/night and a photoperiod of 14:10 [L:D] h. Aphids were collected from plants at 24, 48, 72 and 96 h post-feeding, stored immediately in RNAlater solution (Qiagen, GmbH, Hilden, Germany), incubated overnight at 4°C and frozen to stabilize total RNA according to manufacturer's recommendations.

D. noxia RNA isolation and quality check. Total RNA was isolated from each sample (24, 48, 72 and 96 h post infestation) separately. Silica matrix-based RNA extraction was performed using the RNeasy Plus Kit (Qiagen). Tissues were homogenized using a micropipette according to the manufacturer's recommendations for isolation with some modifications. The gDNA eliminator column was not used for DNase digestion. Instead, an extra on-column-based digestion was performed using an RNase-free DNase set (Qiagen). Purity and integrity of isolated total RNA was assessed by three different methods, including absorbance at 260 and 280 nm in a NanoDrop spectrophotometer (ThermoFisher Scientific, Wilmington, DE), 1% agarose (RNase-free grade) gel electrophoresis using GelGreen staining (Biotium Inc., Hayward, CA) and capillary electrophoresis using an RNA Nano Lab-Chip (Agilent, Santa Clara, CA) and an Agilent 2100 Bioanalyzer system²⁴.

Total RNAs qualifying according to recommended NanoDrop spectrophotometry quality standards were subjected to agarose gel electrophoresis, and samples with intact bands were processed in the Agilent 2100 Bioanalyzer at the Kansas State University Integrated Genomics Facility in Manhattan, KS. The Bioanalyzer provided details about RNA integrity summarized as a RNA integrity number (RIN) (from 1–10; maximum-no degradation) and rRNA (28 s/18 s) ratio (~2 means very high quality RNA)²⁵. Total RNA collected at 24, 48, 72 and 96 h post-feeding by each biotype was pooled to form the treatments RWA1 fed *Dn0* (RWA1/*Dn0*), RWA1 fed *Dn4* (RWA1/*Dn4*), RWA1 fed *Dn7* (RWA1/*Dn7*), RWA2 fed *Dn0* (RWA2/*Dn0*), RWA2 fed *Dn4* (RWA2/*Dn4*), and RWA2 fed *Dn7* (RWA2/*Dn7*).

TrueSeq library construction, sequencing and bioinformatics analysis. mRNA was purified, using oligo-dT beads, from 1 µg of total RNA of each treatment-replication sample provided in TruSeq RNA Sample Preparation Kit v2 (Illumina Inc., San Diego, CA). After purification, first and second strand cDNA was synthesized, 3' ends were repaired and adenylated, and DNA fragments were enriched after adapter ligation and indexing. The generated library was validated and pooled based on the specific index or bar-code for sequencing. Samples were loaded in two lanes and sequenced in a HiSeq 2500 Illumina sequencer at the University of Kansas Medical Center in Kansas City, KS.

Raw sequences generated from sequencing were de-multiplexed to define their sample of origin. Fastq output files of each sample were assessed for quality control (QC) using FastQC (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>), and de novo assembly of sequences was conducted using Trinity software (<http://www.trinity-software.com>). Transcript and gene abundance were estimated and expression was evaluated using DESeq (differential gene expression analysis) and EdgeR (empirical analysis of digital gene expression), both from Bioconductor Open Source Software for Bioinformatics (<http://bioconductor.org/>).

cDNA synthesis and selection of candidate reference genes. cDNA used for PCR standardization and gene amplification was synthesized using Superscript III First-strand Synthesis Supermix (Life Technologies, NY) and oligodT primers according to manufacturer's protocol. For RT-qPCR, an equal amount of total RNA (500 ng) from each sample was used for cDNA synthesis, using the iScript Reverse Transcription Supermix for RT-qPCR (Bio-rad, Hercules, CA).

Candidate reference genes were selected based on literature survey and identification in the library. Primers were designed using Primer 3 (<http://primer3.ut.ee/>) (Table 1). The PCR profile used was 94°C for 30 s, 30 cycles of 94°C for 30 s followed by 55–58°C and 72°C, each for 30 s. The amplicons were sequenced in the Kansas State University Integrated Genomics Facility.

RT-qPCR primer design and test of amplification efficiency. Primers of sequenced transcripts considered for RT-qPCR were designed using Primer Express Software V.3.0.1 (Life Technologies) with slight modifications of the default parameters (Table 2) to avoid hairpin, self-dimer and cross-dimer secondary structures. Primers were selected based on low penalty scores, which indicated a good match to the set parameter settings.

Amplification efficiency of primers was estimated by SYBR Green chemistry RT-qPCR assay using 1, 10⁻¹, 10⁻², 10⁻³ and 10⁻⁴-fold dilutions of pooled cDNAs of three technical replicates for each gene. Each PCR mixture consisted of 1 µl of cDNA, 250 mM of primer and 12.0 µl of iTaq Universal SYBR Green Supermix (Bio-Rad Laboratories) in a total volume of 20 µl. The RT-qPCR profile used was 95°C for 3 min followed by 40 cycles of 95°C for 10 s, 55°C for 30 s and 72°C for 30 s. After RT-qPCR, melt curve analysis was also performed to check for unwanted products or contaminants during the PCR reaction. Three technical replicates were used for standard curve generation. The fluorescence signals were captured, quantification

cycle (C_q) values were tabulated and each mean C_q was plotted against the logarithm of the pooled cDNA dilution factor. The slope in a linear regression model was used to determine the efficiency ($E = [10^{1/(-\text{slope})} - 1] \times 100\%$) for each primer. Correlation coefficient (R²) values were calculated using the standard curve.

RT-qPCR for calculation of expression stability and statistical analysis. The expression stability of the selected genes was calculated in RT-qPCR assay with selected primers and PCR conditions stated above. Each PCR mixture constituted of 1 µl of cDNA from each treatment sample, 250 mM of each primer, and 12.0 µl of iTaq Universal SYBR Green Supermix (Bio-Rad Laboratories) in a total volume of 20 µl. The RT-qPCR profile was 95°C for 3 min followed by 40 cycles of 95°C for 10 s, 55°C for 30 s and 72°C for 30 s. Each reaction was performed for three technical replicates and three biological replicates in 96-well optical grade PCR plates sealed with optical sealing tape (Bio-Rad Laboratories). Melt curves were generated for each sample by heating the PCR amplicon from 65°C to 95°C with a gradual increase of 0.5°C per cycle of 0.5 s.

The Microsoft Excel-based software geNorm¹³, NormFinder²⁶ and BestKeeper²⁷ was used to rank reference genes for stability of expression across all experimental samples. For geNorm and NormFinder analyses, C_q values were transformed to relative quantities using the formula $2^{-\Delta C_q}$, where $\Delta C_q = C_q$ of the gene in selected sample - minimum C_q of corresponding gene in the experiment. The sample with minimum C_q or maximum expression was used as the calibrator with a set value of 1. No transformed C_q values are required for BestKeeper analysis. Other procedures were followed according to the instructions mentioned in the software manual.

- Hudson, M. E. Sequencing breakthroughs for genomic ecology and evolutionary biology. *Mol. Ecol. Resour.* **8**, 3–17 (2008).
- Adams, J. Transcriptome: connecting the genome to gene function. *Nature Edu.* **1**, 195 (2008).
- Stevenson, B. J., Iseli, C., Beutler, B. & Jongeneel, C. V. Use of transcriptome data to unravel the fine structure of genes involved in sepsis. *J. Infect. Dis.* **187**, S308–14 (2003).
- Paim, R. M. *et al.* Validation of reference genes for expression analysis in the salivary gland and the intestine of *Rhodnius prolixus* (Hemiptera, Reduviidae) under different experimental conditions by quantitative real-time PCR. *BMC Res. Notes.* **5**, 128. doi: 10.1186/1756-0500-5-128 (2012).
- Klie, M. & Debener, T. Identification of superior reference genes for data normalisation of expression studies via quantitative PCR in hybrid roses (*Rosa hybrida*). *BMC Res. Notes* **4**, 518 (2011).
- Kheirelseid, E. A., Chang, K. H., Newell, J., Kerinand, M. J. & Miller, N. Identification of endogenous control genes for normalisation of real-time quantitative PCR data in colorectal cancer. *BMC Mol. Biol.* **11**, 12. doi: 10.1186/1471-2199-11-12 (2010).
- Arya, M. *et al.* Basic principles of real-time quantitative PCR. *BioTechniques* **5**, 209–219 (2005).
- Mahoney, D. J. *et al.* Real-time RT-PCR analysis of housekeeping genes in human skeletal muscle following acute exercise. *Physiol. Genomics* **18**, 226–231 (2004).
- Thellin, O. *et al.* Housekeeping genes as internal standards: use and limits. *J. Biotechnol.* **75**, 291–295 (1999).
- Lee, P. D., Sladek, R., Greenwood, C. M. & Hudson, T. J. Control genes and variability: absence of ubiquitous reference transcripts in diverse mammalian expression studies. *Genome Res.* **12**, 292–297 (2002).
- Bustin, S. A. Developments in real-time PCR research and molecular diagnostics. *Expert Rev. Mol. Diagn.* **10**, 713–715 (2010).
- Butte, A. J., Dzauand, V. J. & Glueck, S. B. Further defining housekeeping, or “maintenance,” genes focus on “A compendium of gene expression in normal human tissues.” *Physiol. Genomics* **7**, 95–96 (2001).
- Vandesompele, J. *et al.* Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.* **3**, research0034.1-0034.11 (2002).
- Klie, M. & Debener, T. Identification of superior reference genes for data normalisation of expression studies via quantitative PCR in hybrid roses (*Rosa hybrida*). *BMC Res. Notes* **4**, 518 (2011).
- Gutierrez, L., Mauriat, M., Pelloux, J., Belliniand, C. & Van Wuytswinkel, O. Towards a systematic validation of references in Real-Time RT-PCR. *Plant Cell* **20**, 1734–1735 (2008).
- Nolan, T., Hands, R. E. & Bustin, S. A. Quantification of mRNA using real-time RTPCR. *Nat. Protoc.* **1**, 1559–1582 (2006).
- Morrison, W. P. & Peairs, F. B. [Response model concept and economic impact] *Response Model For An Introduced Pest-The Russian Wheat Aphid* [Quisenberry, S.S. & Peairs, F. B. (eds.)] [1–11] (Thomas Say Publications in Entomology, Lanham, MD, 1998).
- Marasas, C. Wheat management practices and adoption of the Russian wheat aphid integrated control programme in the study area. Socio-economic impact of the Russian wheat aphid integrated control programme. PhD Thesis, University of Pretoria, Pretoria, South Africa. (1999).
- Winnebeck, E. C., Millar, C. D. & Warman, G. R. Why does insect RNA look degraded? *J. Insect Sci.* **10**, 159 (2010).
- Ponton, F., Chapuis, M.-P., Pernice, M., Sword, G. A. & Simpson, S. J. Evaluation of potential reference genes for reverse transcription-qPCR studies of



- physiological responses in *Drosophila melanogaster*. *J. Insect Physiol.* **57**, 840–850 (2011).
21. Li, R. *et al.* Reference gene selection for qRT-PCR analysis in the sweet potato whitefly, *Bemisia tabaci* (Hemiptera: Aleyrodidae). *Plos One* **8**, e53006. doi:10.1371/journal.pone.0053006 (2013).
 22. Yuan, M. *et al.* Selection and evaluation of potential reference genes for gene expression analysis in the brown planthopper, *Nilaparvata lugens* (Hemiptera: Delphacidae) using reverse-transcription quantitative PCR. *PLoS One* **9**, e86503. doi:10.1371/journal.pone.0086503 (2014).
 23. Anathakrishnan, R. *et al.* Comparative gut transcriptome analysis reveals differences between virulent and avirulent Russian wheat aphids, *Diuraphis noxia*. *Arthropod Plant Interact* **8**, 79–88 (2014).
 24. Van de Goor, T. A. (2003) The principle and promise of labchip technology. *Pharma Genomics* **3**, 16–18 (2003).
 25. Schroeder, A. *et al.* The RIN: an RNA integrity number for assigning integrity values to RNA measurements. *BMC Mol. Biol.* **7**, 3 (2006).
 26. Andersen, C. L., Jensen, J. L. & Orntoft, T. F. Normalization of real-time quantitative reverse transcription-PCR data: a model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Res.* **64**, 5245–5250 (2004).
 27. Pfaffl, M. W., Tichopad, A., Prgomet, C. & Neuvians, T. P. Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper-Excel-based tool using pair-wise correlations. *Biotechnol. Lett.* **26**, 509–515 (2004).

Acknowledgments

The authors thank Drs. Kan Yan Zhu and M-S Chen for their critical reviews of the manuscript. This research was supported by the Kansas Wheat Commission, the Kansas Wheat Alliance and Kansas State University. This is contribution 14-240-J from the Kansas Agricultural Experiment Station.

Author contributions

D.K.S. and C.M.S. scoped and designed the study; D.K.S. performed the experiments and analyzed data; D.K.S. and C.M.S. interpreted results and wrote the manuscript.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Sinha, D.K. & Smith, C.M. Selection of Reference Genes for Expression Analysis in *Diuraphis noxia* (Hemiptera: Aphididae) Fed on Resistant and Susceptible Wheat Plants. *Sci. Rep.* **4**, 5059; DOI:10.1038/srep05059 (2014).



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. The images in this article are included in the article's Creative Commons license, unless indicated otherwise in the image credit; if the image is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the image. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>