

Effectiveness of a Web- and Mobile Phone-Based Intervention to Promote Physical Activity and Healthy Eating in Middle-Aged Males: Randomized Controlled Trial of the ManUp Study

Mitch Duncan; Corneel Vandelanotte; Gregory S. Kolt; Richard R. Rosenkranz; Cristina M. Caperchione; Emma S. George; Hang Ding; Cindy Hooker; Mohan Karunanithi; Anthony J. Maeder; Manny Noakes; Rhys Tague; Pennie Taylor; Pierre Viljoen; W. Kerry Mummery

Published Version Information

Citation: Duncan, M., Vandelanotte, C., Kolt, G. S., Rosenkranz, R. R., Caperchione, C. M., George, E. S., et al. (2014). Effectiveness of a web- and mobile phone-based intervention to promote physical activity and healthy eating in middle-aged males: Randomized controlled trial of the manup study. *Journal of Medical Internet Research*, 16(6):e136.

Digital Object Identifier (DOI): 10.2196/jmir.3107

Publisher's Link: <http://www.jmir.org/2014/6/e136/>

This item was retrieved from the K-State Research Exchange (K-REx), the institutional repository of Kansas State University. K-REx is available at <http://krex.ksu.edu>

Type of Paper: Original Paper

Title: A Randomised Controlled Trial examining the effectiveness of a web- and mobile phone-based intervention to promote physical activity and healthy eating in middle-aged males: Outcomes of the ManUp Study

Authors:

Mitch J Duncan,

Corneel Vandelanotte,

Gregory S Kolt,

Richard R Rosenkranz,

Cristina M Caperchione,

Emma S George,

Hang Ding,

Cindy Hooker,

Mohan Karunanithi,

Anthony Maeder,

Manny Noakes,

Rhys Tague,

Pennie Taylor,

Pierre Viljoen,

W Kerry Mummery.

Author Affiliations and contact details:

Mitch J Duncan (PhD): Central Queensland University, Institute for Health and Social Science Research, Centre for Physical Activity Studies Rockhampton, Australia.

Corneel Vandelanotte (PhD): Central Queensland University, Institute for Health and Social Science Research, Centre for Physical Activity Studies, Rockhampton, Australia.

Gregory S Kolt (PhD): University of Western Sydney, School of Science and Health, Sydney, Australia.

Richard R. Rosenkranz (PhD): Kansas State University, Department of Human Nutrition, Manhattan, U.S.A.

Cristina M Caperchione (PhD): University of British Columbia, School of Health and Exercise Sciences, Kelowna, Canada

Emma S George BHSc (Hons): University of Western Sydney, School of Science and Health, Sydney, Australia.

Hang Ding (PhD): CSIRO, The Australian eHealth Research Centre, ICT Centre, Herston, Australia.

Cindy Hooker BHMSc: Central Queensland University, Institute for Health and Social Science Research, Centre for Physical Activity Studies, Rockhampton, Australia.

Mohan Karunanithi (PhD): CSIRO, The Australian eHealth Research Centre, ICT Centre, Herston, Australia.

Anthony J Maeder (PhD): University of Western Sydney, School of Computing,
Engineering and Mathematics, Sydney, Australia

Manny Noakes (PhD): CSIRO Animal, Food and Health Sciences, Adelaide,
Australia

Rhys Tague: University of Western Sydney, School of Computing, Engineering and
Mathematics, Sydney, Australia

Pennie Taylor (PhD): CSIRO Animal, Food and Health Sciences, Adelaide, Australia

Pierre Viljoen (PhD): Central Queensland University, Mackay, Australia

W Kerry Mummery (PhD): University of Alberta, Faculty of Physical Education,
Edmonton, Canada.

Corresponding Author:

Mitch J Duncan: Central Queensland University, Institute for Health and Social
Science Research, Rockhampton, Australia. Phone: +61 7 4930 6977; Fax: +61 7
4930 6402, Email: m.duncan@cqu.edu.au

Abstract

Background: The high number of adult males engaging in low levels of physical activity and poor dietary practices, and the health risks posed by these behaviours, necessitate broad-reaching intervention strategies. IT-based (web and mobile phone) interventions can be accessed by large numbers of people, yet there are few reported IT-based interventions targeting males' physical activity and dietary practices.

Objective: This study examines the effectiveness of a 9-month IT-based intervention to improve the physical activity, dietary behaviours and health literacy in middle-aged males compared to a print-based intervention.

Methods: Participants, recruited offline (e.g. newspaper ads), were randomized into either an IT-based or print-based intervention arm on a 2:1 basis in favour of the fully automated IT-based arm. Participants were adult males aged 35-54 years living in two regional cities in Queensland Australia who could access the internet, owned a mobile phone and were able to increase their activity level. The intervention, ManUp, was informed by social cognitive and self regulation theories and was specifically designed to target males. Educational materials were provided and self-monitoring of physical activity and nutrition behaviours was promoted. Intervention content was the same in both intervention arms, only the delivery mode differed, and content could be accessed throughout the 9-month study period. Participants' physical activity, dietary behaviours, and health literacy were measured using online surveys at baseline, 3 months and 9 months.

Results: A total of 301 participants completed baseline assessments, 205 in the IT-based arm and 96 in the print-based arm. A total of 124 participants completed all three assessments. There were no significant between group differences in physical

activity and dietary behaviours ($p \geq 0.05$). Participants reported an increased number of minutes and sessions of physical activity at 3 months ($b(\text{exp})=1.45$, 95% CI=1.09-1.95; $b(\text{exp})=1.61$, 95% CI=1.17-2.22) and 9 months ($b(\text{exp})=1.55$, 95% CI=1.14-2.10; $b(\text{exp})=1.51$, 95% CI=1.15-2.00). Overall dietary behaviours improved at 3 months ($b(\text{exp})=1.07$, 95% CI=1.03-1.11) and 9 months ($b(\text{exp})=1.10$, 95% CI=1.05-1.13). The proportion of participants in both groups eating higher-fibre bread and low-fat milk increased at 3 months ($b(\text{exp}) = 2.25$, 95% CI = 1.29-3.92; $b(\text{exp})=1.65$, 95% CI = 1.07-2.55). Participants in the IT-based arm were less likely to report that 30 minutes of physical activity per day improves health ($b(\text{exp})=0.48$, 95% CI=0.26-0.90) and more likely to report that vigorous intensity physical activity 3 times per week is essential ($b(\text{exp})=1.70$, 95% CI=1.02-2.82). The average number of logins to the IT-platform at 3 and 9 months was 6.99 (SE=0.86) and 9.22 (SE=1.47), respectively. The average number of self-monitoring entries at 3 and 9 months was 16.69 (SE=2.38) and 22.51 (SE=3.79), respectively.

Conclusions: The ManUp intervention was effective in improving physical activity and dietary behaviours in middle aged males with no significant differences between IT- and print-based delivery modes.

Trial Registration: Australian New Zealand Clinical Trials Registry (ANZCTR):
ACTRN12611000081910

Keywords: physical activity; dietary behaviours; mobile phone; web-based; RCT

Introduction

Regular physical activity and healthy eating are key health behaviours that contribute to reducing the risk of chronic disease [1, 2]. These behaviours, and their impact on health, are particularly relevant for Australian males, as the majority of Australian males are physically inactive and have poor dietary behaviours [3]. For example, approximately 48% of males are not sufficiently physically active and the majority of males do not meet the recommended intake levels of fruit (54%), vegetables (85%), consumption of low-fat dairy (63%) and consumption of foods containing high levels of saturated fat and sugar (70%) [3-5]. Males are also less likely to participate in behavioural and IT-based interventions compared to females [6-8]. In addition, many males have low levels of health literacy, which is the ability to understand and process health information and use this to assist in changing behaviours [9-11]. Health literacy is an important determinant of health and higher levels of health literacy is associated with engagement in various health promoting behaviours including physical activity and fruit and vegetable consumption [12]. Improving the health literacy has been recognised as key to assisting in improving the overall health of Australian males [13]. The high prevalence of poor health behaviours and low levels of health literacy in males combined with the low levels of engagement by males in many behavioural interventions, particularly IT-based interventions, highlights the high need for broad-reaching effective interventions specifically developed for this population.

It is widely acknowledged that web-based- and/or mobile phone-based interventions (IT-based) provide a delivery method that can be conveniently accessed by a large number of individuals thereby increasing the potential reach relative to other commonly used intervention modes such as print-based materials [14-16]. IT-based

interventions have been used to effectively change physical activity and healthy eating behaviours and are viewed positively by males as an intervention delivery mode [7, 17-19]. A broad range of features and components can be implemented in these interventions, including education materials, social interaction/support tools, self-monitoring and goal-setting features, all of which have been associated with increased behaviour change [16, 20]. Increased participant use of and engagement with the intervention platform is also associated with greater behaviour change [21-24]. Mobile phones and smartphones offer participants greater convenience to access intervention materials, and intervention delivery via smartphone and a website is associated with greater levels of self-monitoring and behaviour change when compared to self-monitoring via website only [23]. Therefore delivery of IT-based interventions using a combination of website and mobile devices may be an effective way to increase participant engagement with the intervention and promote greater behaviour change. Despite the potential of IT-based interventions to change physical activity and dietary behaviours, there have been few IT-based interventions that have been specifically developed for and targeted towards males [7, 8]. Therefore, the purpose of this study is to examine the effectiveness of a 9-month web- and mobile phone-based (IT-based) intervention to improve the physical activity and dietary behaviours compared to a print-based intervention [25]. A secondary objective is to compare changes in health literacy between the IT- and print-based intervention groups. It was hypothesised that the IT-based intervention would be more effective in improving outcomes compared to the print-based intervention.

Methods

Design

The rationale, design and methods for the ManUp Study, including outline of the intervention, are described in detail elsewhere and only summarized in brief here [25]. The ManUp Study was a two-arm randomised controlled trial (RCT) with participants randomly allocated to either the IT-based (website and mobile device) intervention arm or a print-based intervention arm. A print-based intervention was selected as the comparison group as they are effective in improving health behaviours [26, 27]. All participants received written and verbal explanation of the project requirements prior to providing consent, provided informed consent prior to participation in the study, and the Central Queensland University (H10/07-131) and the University of Western Sydney Human Research Ethics Committee approved the study (H8605). The study was registered with the Australian New Zealand Clinical Trials Registry (ACTRN12611000081910).

Participants

Males aged 35 to 54 years old who; (1) owned a mobile telephone, (2) had access to the internet, (3) did not have a mobility impairment, (4) resided in the cities of Gladstone or Rockhampton (Queensland, Australia), and (5) were classified as low risk to increase physical activity according to established guidelines were eligible to participate in the study [28]. A combination of online forms and phone contact was used to screen participants for eligibility criteria. To recruit participants, advertisements via local newspapers, trading magazines, face-to-face information sessions with local businesses, and distribution of leaflets and posters to local businesses, medical clinics, and offices of allied health professionals were used. No

participant incentives were provided in the study. Participants (n = 317) were recruited from October 2010 to September 2011, and following initial screening for inclusion criteria, participants were randomly allocated to one of the two intervention arms. As IT-based interventions are less frequently examined in male populations [7, 8], the number of participants allocated to the intervention arm was maximised in a 2:1 ratio in favour of the IT-based intervention arm. Randomization lists were generated by one of the authors (MJD) using freely available software (www.randomization.com). Participants were advised of their group allocation via phone, participants in the IT-based intervention were emailed details to access the intervention platform (website URL, username and password), and participants in the print-based group provided their mailing address to receive the print-based intervention materials. Participants were blinded to group allocation until after baseline assessments were completed. Given participants completed the assessment of outcome measures via online survey, non-blinding of researchers to participant group allocation was unlikely to bias outcomes.

ManUp intervention

The ManUp Study was informed by our reviews of published physical activity and dietary interventions for males, our formative research concerning barriers to physical activity and healthy eating behaviours, and our research regarding males' preferences for IT-based interventions [7, 8, 17, 18]. Both intervention arms provided participants with the same intervention materials and capacity to self-monitor physical activity and dietary behaviours. The IT-based intervention, however, provided participants with the additional ability to receive automated feedback on their progress towards completing their physical activity and dietary behaviour goals (ManUp challenge), as

well as the ability to interact with other participants on the website [25]. Additionally, specific components of the IT-based intervention (My Profiles; My Mates; My Groups) were intended to foster social support between participants via commenting on and viewing the progress of others in line with social cognitive theory. There was little to no interaction between project staff and participants in either intervention arm.

Both interventions arms were provided with educational materials that were specifically designed to present information on the benefits of physical activity and healthy eating and on the volume and type of activity needed to achieve health benefits. Materials provided to participants allowed daily self-monitoring of physical activity and dietary behaviours and highlighted the importance of self-monitoring as a way to change behaviour and keep track of the changes made. Participants could record physical activity and dietary behaviours using any ‘metric’ specified in Table 1. Informed by social cognitive theory and self regulation theory, ManUp “challenges” were developed to change target behaviours by providing a goal for behaviour change, having participants engage in goal setting and self-monitoring behaviours and also build confidence to make positive changes to behaviours [25, 29, 30]. An overview of these theories and the role of self-monitoring in changing behaviour can be found elsewhere [29-31].

ManUp challenges

The ManUp challenges consisted of six physical activity and a multi-component healthy eating challenge. Each challenge had three different “strengths” (light, mid, full), which varied the duration and the amount of activity or healthy eating that males were asked to achieve in order to complete the challenge. In order to complete a

challenge participants had to record the required number of minutes/distance/steps for activities or the number of healthy eating goals before the end of the challenge period, failure to do this meant the challenge was not completed. The variation between challenge strengths was intended to provide participants with an appropriate target relative to their current level of physical activity or healthy eating, or to provide a progression towards engaging in higher levels of physical activity or healthy eating. Although the challenges could be completed in any order preferred by participants and there was no requirement to complete all of the different strength challenges or different physical activity challenges. The different activities selected for inclusion were based on those activities frequently performed by Australian males [32]. The ManUp healthy eating challenges were based on achieving a maximum of ten daily healthy eating goals. These goals were informed by the dietary guidelines for Australian adults that promote dietary diversity and encourage the reduction of the intake of saturated fat, salt, alcohol and foods that contain added sugars [33]. Although the challenges were informed by the sorts of physical activities frequently participated in by males, and the need to promote dietary diversity, they were not intended to promote adherence to public health guidelines for either physical activity or dietary behaviours. Rather, they were designed to increase overall engagement in physical activity and healthy eating. Further details on the types of different physical activities and dietary behaviours targeted, the requirements for each challenge, and supporting educational materials are provided in Table 1 and elsewhere [25].

Table 1. Description of the ManUp physical activity and healthy eating challenges

| Activity | Light Strength (3 weeks) | Mid Strength (6 weeks) | Full strength (12 weeks) |
|-----------------------------|---|--|---|
| Walking | 1.5 hrs/week or 7500 steps/day | 2.5 hrs/week or 10000 steps/day | 3.5 hrs/week or 12000 steps/day |
| Cycling | 1 hr/week or 25 km/week | 2 hrs/week or 50 km/week | 4 hrs/week or 100 km/week |
| Swimming | 0.5 hr/week or 1 km/week | 1 hr/week or 2 km/week | 1.5 hrs/week or 3 km/week |
| Running | 0.5 hr/week or 5 km/week | 1 hr/week or 10 km/week | 2.0 hrs/week or 20 km/week |
| Sport & Recreation | 0.5 hr/week | 1 hr/week | 1.5 hrs/week |
| Strengthening | Set 8 exercises 1 x set (8–10 reps) 2 x/week | Set 8 exercises 2 x set (8–10 reps) 2 x/week | Set 8 exercises 3 x set (8–10 reps) 2 x/week |
| Healthy Eating ^a | ≥3 healthy eating goals/day | ≥5 healthy eating goals/day | ≥7 healthy eating goals/day |

Notes. a. The ManUp healthy eating goals were: (1) eat two serves of fruit, (2) eat five serves of vegetables, (3) eat a serve of fish, (4) choose whole-grain bread instead of white bread, (5) choose low-fat dairy products, (6) have a soft drink- (soda-) free day, (7) have an alcohol-free day, (8) have an red-meat-free day, (9) have an unhealthy snack-free day, and (10) have a fast-food-free day

Intervention arms

IT-based intervention arm

Upon completing the baseline assessment participants in the IT-based intervention arm received access to the password-protected ManUp website, which had six main sections [25]. The six sections were: 1) *My Profile* - where participants could review their current challenges, record their progress towards any current challenges, post personal updates to their profile, schedule future activities, and view information on the groups they were a member of and the list of their 'mates' (online friends on the website); 2) *My Progress* - where participants could review their progress towards their current challenges; 3) *My Mates* - where participants could search for online friends and view their mates progress. Online friends were limited only to participants allocated to the IT-based intervention, participants could not view an online list of other participants **nor were they informed by project staff who else was enrolled in the study** due to privacy concerns but participants could search for other users by entering a name or part of a name into the search tool provided on the website; 4) *My Groups* - where participants could create a group and view the progress of groups they were part of; 5) *My Weight* - which provided participants with information on the benefits of achieving a healthy weight, and allowed them to record their height, weight and waist circumference. This section did not allow participants to track these metrics

over time; rather it provided immediate feedback on what category (BMI category, or waist circumference category) the respective measure was classified as in comparison to established categories for BMI and waist circumference [34]; 6) *Information Centre* - which provided educational materials related to physical activity and healthy eating, and the challenges [25].

As a form of online social support, participants could comment on their mates' My Profile page. In addition participants could also 'challenge' their mates to complete a physical activity or healthy eating challenge either in a one-on-one basis, or as part of a larger group. A mobile phone web application was developed as an additional tool to facilitate quick and convenient recording of progress towards the ManUp challenges. The mobile phone web application only allowed users to self-monitor behaviour, body weight, and to review progress towards challenge completion. Any participant in the IT-based intervention arm who owned a mobile phone capable of accessing the internet had access to the mobile phone web application. Figure 1 shows the My Profile section of the website and Figure 2a and 2b, displays the app data entry screen for healthy eating, the screen showing feedback information on the level of activity needed to complete a particular challenge. Graphical and text based feedback in terms of progress towards the completion of a challenge was automatically provided by both the website and app, and was updated based on participant's self-monitoring entries of physical activity and health eating. Participants were not provided with any detailed instruction on how to use the IT-platform or the frequency it with which it should be used. The initial email providing login details suggested they should visit the intervention platform and initiate a ManUp challenge. A short video presenting the main features and functionality of the

website was available for viewing on the front page of the website without having to login to the website. This video was viewed 243 times in total throughout the intervention period.

Figure 1. Screenshot of the My Profile section of the website.





Figure 2a and 2b. Screenshots of the ManUp app healthy eating data entry screen (2a) and challenge progress feedback screen (2b).

Figure 2a.



Figure 2b.



Print-based intervention arm

Participants in the print-based group received a hard-copy booklet that provided the same educational materials (including content from the My Weight section) and

ManUp challenges as those provided to participants in the IT-based intervention.

Participants in the print-based group were provided with information on using the provided log sheets and could self-monitor progress and/or successful completion of the ManUp physical activity or healthy eating challenges using the log sheets.

Participants in the print-based group were not provided with information regarding their peers who were also part of this group. The hard copy booklet was not collected from participants and no information on the challenges completed **or self-monitoring** was obtained. Hard copy booklets were not collected due to both logistical reasons and also to allow participants to keep a record of their progress to assist in behaviour change.

Outcome measures

Participants completed online surveys at baseline (0 months), 3 months, and 9 months to assess socio-demographic, behavioural, and health literacy outcomes. Measures of satisfaction with the intervention were also obtained at the 9 months assessment point.

All participants received up to 3 phone calls or emails at each assessment point to remind them to complete their assessments.

Physical activity

Physical activity was assessed using the Active Australia Questionnaire, a valid and reliable instrument that is also sensitive to change in physical activity [35-38]. This questionnaire asks participants to report the duration of recreational and transport walking, moderate and vigorous intensity physical activity in the previous week, and the number of times (sessions) they engaged in these activities. Standard scoring

protocols were applied to provide two outcomes; total minutes of physical activity, and the total number of sessions of physical activity [35].

Dietary behaviours

Dietary behaviours were assessed using 19 items adapted from existing instruments used to monitor dietary habits of the Australian population [39, 40]. These items have sound psychometric properties [39]. Two separate items assessed the daily number of servings of fruit and vegetables consumed in the last week, response options ranged from zero servings (don't eat this food) to ten or more servings. The frequency that red meat, fish, meat products (sausages, salami, meat pies, etc.), cooked cereals, soft drink, chips, takeaway foods, and sweet or savoury foods were consumed in the last week was assessed using response options from rarely/never (don't eat this food) to more than ten times. The type of milk (whole milk (i.e. full cream)), reduced fat, soy milk, condensed milk, don't drink milk) and bread (white, wholemeal, multigrain, rye, sour dough, other, don't eat bread) usually consumed was also assessed. Three dietary outcomes were created; type of milk consumed (reduced fat vs. whole milk), type of bread consumed ((higher fibre) wholemeal, multigrain, white with high fibre, sour dough, rye vs. white) and an overall index of other dietary behaviours – the dietary score. The dietary score was created by summing the number of servings and number of times the following foods were consumed: fruit, vegetables, red meat, fish, meat products, soft drink, chips, takeaway (take-out) foods, sweet and savoury foods. Several items were reversed score so that higher dietary scores (a better diet) reflected more frequent consumption of healthy food and less frequent consumption of less healthy foods. The dietary score reflected the fact that the ManUp healthy eating

challenges focussed on maximising consumption of healthful foods and minimising consumption less healthy foods.

Health literacy

Health literacy in relation to physical activity was assessed using the five awareness items from the Active Australia Questionnaire [35]. Using a five-point Likert type scale from strongly agree to strongly disagree, the items assess awareness of the benefits associated with physical activity participation, the intensity and duration required to receive health benefits. Dietary behaviour literacy was assessed using the Nutritional Literacy Survey, a valid and reliable 28-item instrument that assesses participants' understanding of the type of foods that promote heart health, and the fat and cholesterol content of different foods and portion sizes [41].

Satisfaction

Participant satisfaction with the intervention platform and challenge concept were assessed using 4 items. Using a 5 point scale ranging from *Strongly Agree* to *Strongly Disagree*, participants indicated if they would like to continue to use the IT- or print-based materials, if the materials (print booklet, or IT-based platform) were easy to use, and if they liked the overall concept of the physical activity and healthy eating challenges.

IT Platform usage

Usage of the IT-based platform was measured using inbuilt tracking software measuring the number of times a participant logged into the web- and mobile-based

platform, made a self-monitoring entry, the type and number of challenges they initiated and completed.

Sample Size

Using established methods to estimate sample size [42], the study was powered to detect a 60-minute change in moderate-to-vigorous intensity physical activity per week from baseline to 9 months using an alpha level of 0.05 and a power level of 90%. Based on this calculation, it was estimated that 197 participants would be required. This number was increased, however, to account for the 2:1 allocation of participants in favour of the IT-based intervention arm and the expected drop-out rate of participants (45%) [21, 43]. A higher drop-out rate was used in the current study given the acknowledged difficulty in engaging and retaining males in interventions [7, 8]. As a result, the estimated total sample size was 321; 107 to be allocated to the print-based group and 214 to be allocated to the IT-based group [25].

Analysis


Comparisons between groups at baseline were conducted using generalized linear models and chi-square tests. Comparisons between those participants completing all three assessment points (completers) and those completing less than three assessment points (non-completers) were made on age, education, physical activity, dietary behaviours and health literacy, using t-tests (where parametric assumptions were met) or Mann-Whitney U for continuous variables, and chi-square tests for categorical variables. Generalized linear mixed models use all available data at each time point allowing participants with missing data at follow-up time points to be retained in the

analysis. Therefore, generalized linear mixed models with an unstructured covariance matrix were used to examine change over time and differences between intervention arms in physical activity, dietary behaviours, and health literacy outcomes. All analyses were adjusted for baseline age, occupation, and education as these variables are likely to impact upon the physical activity and dietary behaviours of males [44]. Outcomes of the generalized linear mixed model analyses are reported as exponentiated coefficients ($b(\exp)$). To explore the impact of missing data, a sensitivity analysis using baseline observation carried forward (BOCF) for participants with missing data at follow-up time points was performed for physical activity, dietary behaviours and health literacy outcomes; this analysis also adjusted for baseline age, occupation, and education. Comparison of change in physical activity, dietary behaviours and health literacy with and without BOCF revealed only small differences in the magnitude of these outcomes with the exception of consumption of higher fibre bread and low fat milk consumption. For both of these outcomes, the significant time effects present at 3 months in the analysis without BOCF were in the same direction although not statistically significant in the analysis with BOCF. Given these minor differences, only the results from the analyses without BOCF are reported.

Analyses examining the relationship between usage of the IT-platform and change in behaviour within the IT-based intervention arm were conducted using generalized linear models adjusted for age, occupation, education and the baseline level of the outcome examined. The specific model type, link function used for analyses and the total number of observations included are listed in the footnotes of Tables 3, 4 and 5.

All analyses were conducted with SPSS (version 20), followed intention-to-treat principles, and used an alpha level of 0.05.

Figure 3. Flow of participants through the RCT.



Results

Participants

The flow of participants through the study, including the number of participants completing each assessment, is provided in Figure 3. A total of 327 males expressed an interest in participation in the study; 10 males were excluded as they did not satisfy eligibility criteria or were no longer interested in participating; 317 males were randomised to an intervention arm of which 16 withdrew from the study prior to completing a baseline assessment (Figure 3). A total of 301 participants completed the baseline assessment, 159 completed the three-month assessment, 148 completed the nine-month assessment and 125 completed all three assessment points. No significant differences were observed between those who completed three assessment points (completers) compared to those completing less than three assessment points (non-completers) on baseline minutes and sessions of physical activity, dietary score and type of bread consumed (data not shown). Completers were significantly different from non-completers in terms of age ($M = 45.11$, $SE = 0.51$ vs. $M = 43.32$, $SE = 0.44$; $P = 0.01$), the proportion working in professional level occupations (66.4% vs. 49.4%; $p = 0.026$), the proportion reporting a university-level education (62.6% vs. 41.5%; $P = 0.002$) the proportion reporting to consume low-fat milk (66.4% vs. 49.7%; $P = 0.01$), the proportion reporting that at least 20 minutes of vigorous intensity physical activity 3 times per week is essential to improve health (35.8% vs. 64.2%; $P = 0.01$), and nutritional literacy ($M = 25.53$, $SE = 0.16$ vs. $M = 24.98$, $SE = 0.16$, $P = 0.02$).

There were no significant baseline differences between the print-based arm and the IT-based arm for any demographic behavioural, and health literacy variable, with the exception that there were fewer participants who agreed that 30 minutes of physical

activity is enough to improve health in the IT-based arm compared to the print-based arm (Table 2). In line with the eligibility criteria, all participants owned a mobile phone, and 151 (73.2%) of the IT-based intervention owned a phone that could access the internet.

Table 2. Socio-demographic and anthropometric characteristics of participants at baseline.

| | Print-based | | IT-based | | <i>P</i> |
|--------------------------|-------------|---|----------|---|----------|
| | n | %, M (SE), median (1 st Quartile, 3 rd Quartile) | n | %, M (SE), median (1 st Quartile, 3 rd Quartile) | |
| Age | 96 | 43.84 (0.59) | 205 | 44.17 (0.41) | 0.66 |
| Occupation | | | | | 0.64 |
| Professional | 52 | 54.2 | 118 | 57.6 | |
| White Collar | 8 | 8.3 | 16 | 7.8 | |
| Blue Collar | 23 | 24.0 | 37 | 18.0 | |
| Other | 13 | 13.5 | 34 | 16.6 | |
| Education Level | | | | | 0.72 |
| Secondary School or Less | 20 | 20.8 | 45 | 22.0 | |

| | | | | | |
|---|----|--------------------|-----|-------------------|------|
| TAFE | 25 | 26.0 | 61 | 29.8 | |
| University | 51 | 53.1 | 99 | 48.3 | |
| Self-Reported BMI | | | | | 0.46 |
| Healthy Weight | 13 | 13.5 | 19 | 9.3 | |
| Overweight | 41 | 42.7 | 85 | 41.5 | |
| Obese | 42 | 43.8 | 101 | 49.3 | |
| Self-reported minutes of physical activity per week | 96 | 277.94 (29.15) | 205 | 286.12 (24.72) | 0.83 |
| Self-reported sessions of physical activity per week | 96 | 4.0 (1.0, 8.0) | 205 | 4.0 (1.0, 7.0) | 0.95 |
| Dietary score | 96 | 52.0 (46.0, 56.75) | 205 | 52.0 (47.0, 57.0) | 0.33 |
| Higher Fibre bread | 93 | 57.0 | 195 | 68.2 | 0.06 |
| Low fat milk | 87 | 57.5 | 182 | 56.0 | 0.83 |
| Physical Activity Literacy (% agree) | | | | | |
| ≥ 30 minutes/day improves health | 79 | 82.3% | 144 | 70.2% | 0.03 |

| | | | | | |
|---|----|-------------|-----|-------------|-------|
| 30 minutes brisk walking improves health | 79 | 82.3% | 153 | 74.6% | 0.14 |
| 20 minutes of vigorous activity 3 times a week is essential | 54 | 56.3% | 139 | 67.8% | 0.051 |
| 10 minute blocks of activity are okay | 52 | 54.2% | 106 | 51.7% | 0.69 |
| Moderate activity can improve health | 87 | 90.6% | 177 | 86.3% | 0.29 |
| Nutritional Literacy | 96 | 25 (24, 26) | 205 | 26 (24, 27) | 0.66 |

Notes. TAFE is a provider of vocational non-bachelor education up to level of advanced diploma.

Change in physical activity and dietary behaviours

There were no significant between-group differences or group-by-time interaction effects in any of the physical activity and dietary behaviours examined however significant main effects for time were observed (Table 3). Self-reported minutes (3 months: $b(\text{exp})=1.45$, 95% CI=1.09-1.95; 9 months: $b(\text{exp})=1.55$, 95% CI=1.14-2.10) and sessions of physical activity (3 months: $b(\text{exp})=1.61$, 95% CI=1.17-2.22; 9 months: $b(\text{exp})=1.51$, 95% CI=1.15-2.00) were significantly higher at 3 and 9 months

compared to baseline in both groups. Dietary scores were significantly higher (improved) at both 3 and 9 months compared to baseline (3 months: $b(\text{exp})=1.07$, 95% CI=1.03-1.11 9 months: $b(\text{exp})=1.10$, 95% CI=1.05-1.13) in both groups. A significantly higher proportion of participants reported consuming higher-fibre bread ($b(\text{exp})=2.25$, 95% CI=1.29-3.92); and low-fat milk ($b(\text{exp})=1.65$, 95% CI=1.07-2.55) at three months compared to baseline in both groups; consumption of higher-fibre bread and low-fat milk were not significantly higher at nine months compared to baseline in both groups.

Table 3. Comparison of self-reported measured health behaviours between intervention groups over the intervention period.

| | | <i>Model Effects</i> | | |
|--|------------------------|----------------------|----------|-----------------|
| | | | <i>P</i> | |
| | <i>b (exp) 95% C.I</i> | Group | Time | Group * Time |
| Self-report minutes of physical activity per week^a | | 0.60 | <0.001 | 0.66 |
| <i>IT-based vs. print-based^b</i> | 1.03 (0.78-1.36) | | | |
| <i>3 mths vs. 0 mths^b</i> | 1.45 (1.09-1.95) | | | |
| <i>9 mths vs. 0 mths^b</i> | 1.55 (1.14-2.10) | | | |

| | | | | |
|---|------------------|------|--------|------|
| Self-report sessions of physical activity per week^a | | 0.32 | <0.001 | 0.55 |
| <i>IT-based vs. print-based^b</i> | 0.97 (0.75-1.25) | | | |
| <i>3 mths vs. 0 mths^b</i> | 1.61 (1.17-2.22) | | | |
| <i>9 mths vs. 0 mths^b</i> | 1.51 (1.15-2.00) | | | |
| Dietary score^c | | 0.68 | <0.001 | 0.09 |
| <i>IT-based vs. print-based^b</i> | 1.02 (0.98-1.06) | | | |
| <i>3 mths vs. 0 mths^b</i> | 1.07 (1.03-1.11) | | | |
| <i>9 mths vs. 0 mths^b</i> | 1.10 (1.05-1.13) | | | |
| Higher-fibre bread^d | | 0.05 | <0.001 | 0.92 |
| <i>IT-based vs. print-based^b</i> | 1.60 (0.94-2.71) | | | |
| <i>3 mths vs. 0 mths^b</i> | 2.25 (1.29-3.92) | | | |
| <i>9 mths vs. 0 mths^b</i> | 1.89 (0.99-3.60) | | | |
| Low fat milk^e | | 0.54 | 0.002 | 0.90 |
| <i>IT-based vs. print-based^b</i> | 0.88 (0.52-1.49) | | | |
| <i>3 mths vs. 0 mths^b</i> | 1.65 (1.07-2.55) | | | |
| <i>9 mths vs. 0 mths^b</i> | 1.41 (0.92-2.17) | | | |

Notes.

- a. Model (negative binomial with log link) included age, education level, and occupational classification as covariates. Number of observations = 616
- b. Reference category for comparison.
- c. Model (negative binomial with log link) included age, education level, and occupational classification as covariates. Number of observations = 608. This outcome was examined as the change in the total number of times the food was consumed and the servings of a food.
- d. Model (binomial with logit link) included age, education level, and occupational classification as covariates. Number of observations = 587. This outcome was examined as the change in the proportion of participants consuming higher-fibre bread.
- e. Model (binomial with logit link) included age, education level, and occupational classification as covariates. Number of observations = 542. This outcome was examined as the change in the proportion of participants consuming low fat milk.

Table 4. Comparison of health literacy outcomes between intervention groups over the intervention period^a.

| | | <i>Model Effects</i> | | |
|--|-------------------------|----------------------|----------|-----------------|
| | | | <i>P</i> | |
| | <i>b</i> (exp) 95% C.I. | Group | Time | Group * Time |
| ≥ 30 minutes/day improves health^b | | 0.17 | 0.11 | 0.28 |
| <i>IT-based vs print-based^c</i> | 0.48 (0.26-0.90) | | | |
| <i>3 mths v.s. 0 mths^c</i> | 1.02 (0.50-2.09) | | | |
| <i>9 mths v.s. 0 mths^c</i> | 1.37 (0.65-2.89) | | | |
| 30 minutes brisk walking improves health^b | | 0.91 | 0.01 | 0.13 |
| <i>IT-based vs print-based^c</i> | 0.63 (0.34-1.16) | | | |
| <i>3 mths v.s. 0 mths^c</i> | 1.33 (0.58-3.06) | | | |
| <i>9 mths v.s. 0 mths^c</i> | 1.51 (0.60-3.81) | | | |
| 20 minutes of vigorous activity 3 times a week is essential^b | | 0.01 | 0.99 | 0.88 |

| | | | | |
|--|------------------|------|-------|------|
| <i>IT-based vs print-based^c</i> | 1.70 (1.02-2.82) | | | |
| <i>3 mths v.s. 0 mths^c</i> | 0.96 (0.49-1.87) | | | |
| <i>9 mths v.s. 0 mths^c</i> | 1.08 (0.57-2.04) | | | |
| 10 minute blocks of activity are okay^b | | 0.33 | 0.001 | 0.58 |
| <i>IT-based vs print-based^c</i> | 0.89 (0.54-1.45) | | | |
| <i>3 mths v.s. 0 mths^c</i> | 1.51 (0.83-2.72) | | | |
| <i>9 mths v.s. 0 mths^c</i> | 2.52 (1.28-4.94) | | | |
| Nutrition Literacy^d | | 0.78 | 0.44 | 0.51 |
| <i>IT-based vs print-based^c</i> | 1.01 (0.99-1.03) | | | |
| <i>3 mths v.s. 0 mths^c</i> | 1.01 (0.99-1.03) | | | |
| <i>9 mths v.s. 0 mths^c</i> | 1.01 (0.97-1.05) | | | |

Notes.

a. Analysis of change in the physical activity literacy outcome of “moderate physical can improve health” is not reported as there was insufficient variation in the outcome to allow the model to be accurately estimated. The proportion of participants agreeing with this statement at each time point in each group is: Baseline: IT-based = 86.3%, print-based = 90.6%; Three months: IT-based = 88.9%, print-based = 86.7%; Nine months: IT-based = 100.0%, print-based = 95.8%.

b. Model (binomial with logit link) included age, educational level, and occupational classification as covariates. Number of observations = 608

c. Reference category for comparison.

d. Model (negative binomial with log link) included age, educational level, and occupational classification as covariates. Number of observations = 608

Change in health literacy

A significantly lower proportion of participants in the IT-based intervention arm reported agreeing with that 30 minutes of physical activity per day improves health compared to the print-based arm ($b(\text{exp})=0.48$, 95% CI=0.26-0.90); there were no significant time or group-by-time interaction effects for this outcome (Table 4). A significantly higher proportion of participants in the IT-based intervention arm reported agreeing that 20 minutes of vigorous intensity physical activity performed 3 times per week is essential to improve health ($b(\text{exp})=1.70$, 95% CI=1.02-2.82).

There were no significant group or group-by-time interaction effects for the proportion of participants reporting that blocks of a minimum of 10 minutes physical activity are acceptable to acquire health benefits; however, a significantly higher proportion of participants reporting agreeing with this statement at 9 months compared to baseline in both groups ($b(\text{exp})=2.52$, 95% CI=1.28-4.94). No other statistically significant differences were observed in physical activity and nutrition literacy (Table 4).

ManUp challenges

Due to the difficulty in obtaining records in log book usage and challenge completion in the print-based intervention arm, data on the usage of ManUp challenges is only reported for the IT-based intervention arm. Figure 4 demonstrates the number of participants in the IT-based intervention arm who started and completed light, mid, or full strength physical activity and healthy eating challenges over the 9-month intervention period. A higher number of participants initiated physical activity challenges compared to the healthy eating challenges, and no participants completed a healthy eating challenge - this was due to participants not completing the required number of healthy eating goals in the specified time period (3, 6 or 12 week challenge duration depending on challenge strength selected) and thus, they 'did not complete the challenge'. When examining the number of challenges initiated and completed over the 9-month period by all IT-group participants, light strength physical activity challenges were the most frequently selected ($n = 147$) and completed ($n = 137$), followed by mid strength physical activity challenges (initiated: $n = 80$; completed: $n = 68$), and full strength physical activity challenges (initiated: $n = 69$; completed: $n = 53$). Healthy eating challenges were initiated at a lower frequency compared to physical activity challenges but followed a similar pattern; light strength healthy eating challenges were the most frequently selected (initiated: $n = 60$; completed: $n = 0$), followed by mid strength healthy eating challenges (initiated: $n = 28$; completed: $n = 0$), and full strength healthy eating challenges (initiated: $n = 14$; completed: $n = 0$). Information on the type of physical activity challenges selected by participants is provided in Figure 5. Walking was the most frequently selected challenge type, followed by cycling and strengthening activities. Figure 6 shows the number of

different health eating goals selected; Fastfood, Softdrink and Alcohol Free days and eating two pieces of fruit were the most frequently selected goals across all challenges.

Figure 4. Number of Participants in IT-based intervention who started and completed ManUp Challenges




Figure 5. Number of Physical Activity Challenge Type Completed by Challenge Strength.





Figure 6. Number of Number of Healthy Eating Goals Selected by Challenge Strength.



Note. This data is based on goals logged via the website only.

IT-platform usage

The median number of logins to the IT-platform per week at 3 months and 9 months was 2.00 (IQR = 6.00) and 2.00 (IQR = 6.50), respectively; the average number of logins to the IT-platform at these same time periods was 6.99 (SE = 0.86) and 9.22 (SE = 1.47). Median number of self-monitoring entries per week at 3 months and 9 months was 1.00 (IQR = 20.0) and 1.00 (IQR = 21.5.0) respectively; the average number of self-monitoring entries at 3 months and 9 months was 16.69 (SE = 2.38) and 22.51 (SE = 3.79), respectively. Participants who logged in 2 or more times in the first three months of the intervention made significantly more self-monitoring entries (Median = 18.00, IQR = 38.00) compared to participants logging in less than 2 times (Median = 0.00, IQR = 0.00) ($U = 1195.50$, $P = <0.001$). Figure 7 shows the number of users logging in at least once and making at least one self-monitoring entry each week over the intervention period. Following the initial reduction in usage between week 1 and week 3, usage continued to decline throughout the intervention period. No measure of usage of the IT-based platform was associated with any of the physical activity or dietary behaviours examined (Table 5). Use of the ‘mate’ feature and posts to update of their progress was low; 21 participants used the ‘mate’ feature and no participants using this feature had more than one mate on the platform; 36 participants used the post feature (minimum = 1 post, maximum = 26 posts). Minimal reports of bugs or errors (<10) on the IT-based platform were received during the intervention period via the “report a bug” feature. Of the 33 participants who completed survey items regarding satisfaction of the print materials, 33.3% agreed or strongly agreed that they would like to continue to use the ManUp booklet in the future, and 75.8% agreed or strongly agreed that it was easy to use. Of the 60 participants who completed survey items regarding satisfaction of the website, 25.0% of participants

agreed or strongly agreed that they would like to continue to use the website and 85.0% reported it was easy to use. Of the 139 participants who completed items regarding satisfaction with the concept of the physical activity and healthy eating challenges 60.4% and 52.7% of participants in print and IT-based groups, respectively, reported satisfaction with the physical activity challenge with no significant differences between groups ($\chi^2 = .748, p = .387$). There were no differences in the proportion of participants in the print (60.4%) and IT-based groups (48.4%) who reported being satisfied with the healthy eating challenge ($\chi^2 = 1.834, p = .176$)

Figure 7. Participant usage of the IT-based intervention platform each week over the intervention period.




Table 5. Associations between IT-platform usage and self-reported physical activity and dietary behaviours.

| | | | <i>Model Effects</i> | |
|---|-------------------------|--|-------------------------|--|
| | | | <i>Number of logins</i> | <i>Number of self-monitoring entries</i> |
| | <i>Number of logins</i> | <i>Number of self-monitoring entries</i> | <i>Number of logins</i> | <i>Number of self-monitoring entries</i> |
| | <i>b (exp) 95% C.I.</i> | <i>b (exp) 95% C.I.</i> | <i>P</i> | <i>P</i> |
| Self-report minutes of physical activity per week | | | | |
| <i>3 mths^a</i> | 1.00 (0.98-1.01) | 1.00 (0.997-1.01) | .43 | .38 |
| <i>9 mths^b</i> | 1.00 (0.99-1.00) | 1.00 (1.00-1.01) | .25 | .10 |
| Self-report sessions of physical activity per week | | | | |
| <i>3 mths^c</i> | 0.99 (0.98-1.01) | 1.01 (1.00-1.01) | .19 | .05 |
| <i>9 mths^d</i> | 1.00 (0.99-1.00) | 1.00 (1.00-1.01) | .41 | .16 |
| Dietary score | | | | |
| <i>3 mths^e</i> | 1.00 (0.998-1.00) | 1.00 (0.99-1.00) | .65 | .76 |

| | | | | |
|---------------------------|-------------------|-------------------|-----|-----|
| <i>9 mths^f</i> | 1.00 (0.998-1.00) | 1.00 (0.999-1.00) | .11 | .63 |
| Higher Fibre bread | | | | |
| <i>3 mths^g</i> | 1.03 (0.93-1.13) | 1.02 (0.99-1.05) | .59 | .25 |
| <i>9 mths^h</i> | - | - | - | - |
| Low fat milk | | | | |
| <i>3 mthsⁱ</i> | 0.97 (0.90-1.04) | 1.01 (0.98-1.04) | .33 | .71 |
| <i>9 mths^j</i> | 1.00 (0.98-1.03) | 1.00 (0.99-1.01) | .71 | .54 |

Notes:

a. Model (tweedie with log link) included age, educational level, occupational classification, access to the mobile platform, and baseline minutes of physical activity as covariates. Number of participants = 101

b. Model (tweedie with log link) included age, educational level, occupational classification, access to the mobile platform, and baseline minutes of physical activity as covariates. Number of participants = 100

c. Model (negative binomial with log link) included age, educational level, occupational classification, access to the mobile platform, and baseline sessions of physical activity as covariates. Number of participants = 101

d. Model (negative binomial with log link) included age, educational level, occupational classification, access to the mobile platform, and baseline sessions of physical activity as covariates. Number of participants = 100

e. Model (tweedie with log link) included age, educational level, occupational classification, access to the mobile platform, and baseline dietary score as covariates.

Number of participants = 99

f. Model (tweedie with log link) included age, educational level, occupational classification, access to the mobile platform, and baseline dietary score as covariates.

Number of participants = 96

g. Model (binomial with logit link) included age, educational level, occupational classification, access to the mobile platform, and baseline bread consumption as covariates. Number of participants = 93

h. Results are not reported for this time point as the model had partial or complete separation and parameters could not be reliably estimated.

i. Model (binomial with logit link) included age, educational level, occupational classification, access to the mobile platform, and baseline milk consumption as covariates. Number of participants = 82

j. Model (binomial with logit link) included age, educational level, occupational classification, access to the mobile platform, and baseline milk consumption as covariates. Number of participants = 80

Discussion

This study examined the relative effectiveness of the ManUp intervention materials delivered by an IT-based intervention platform compared to a print-based intervention to improve middle aged males' physical activity and dietary behaviours, and health literacy of these behaviours. Analyses revealed significant improvements over time in self-reported minutes and sessions of physical activity and self-reported overall dietary behaviours in both groups. These changes did not significantly differ between participants receiving access to the IT - or print-based intervention materials. Three components of physical activity literacy changed during the intervention period. First, a lower proportion of participants in the IT-based intervention arm reported agreeing that 30 minutes of physical activity per day is enough to improve health. Second, a higher proportion of participants in the IT-based intervention arm reported agreeing that 20 minutes of vigorous intensity physical activity 3 times per week is necessary to improve health. Finally, a higher proportion of participants from both intervention arms reported agreeing that accumulating physical activity in blocks of minimum 10 minutes are acceptable to improve health at the nine month assessment point compared to baseline. Nutrition literacy did not change over time, nor between intervention arms.

IT- and print-based interventions have been shown to be effective in improving physical activity, dietary behaviours, or both behaviours in male populations in earlier research [19, 45-48]. These studies demonstrated positive intervention effects on target behaviours for time periods ranging between 14 weeks to 12 months [19, 47]. Consistent with this literature, the current study demonstrated positive changes in both physical activity and dietary behaviours over a nine-month period.

Notwithstanding the limitations of the self-report data in the current study, the sustained nature of the changes is encouraging given that poor physical activity and dietary behaviours are prevalent and significant contributors to chronic disease risk in male populations [3-5, 13]. Most previous research has targeted only one of these behaviours and comparatively fewer interventions aimed at males have targeted both behaviours simultaneously [7, 8]. As such outcomes of the ManUp study contribute to the evidence that it is feasible to target and significantly improve multiple health behaviours simultaneously in various populations [49-51]. The ability of the intervention platform to improve physical activity and dietary behaviours and the lack of differences between the delivery modes suggests that IT-based approaches are useful to improve males' engagement in these behaviours. This observation is useful given the large number of males engaging in poor levels of physical activity and dietary behaviours and the potential reach of IT-based interventions relative to other delivery modes. However as others have noted the challenge lies in attracting and engaging individuals to the platform [15, 52], particularly males who are an acknowledged difficult group to engage in these types of interventions [7, 8].

An advantage of IT-based platforms is that participant engagement and usage can be monitored throughout the intervention period. Figure 7 shows a modest level of engagement and usage in the initial weeks of the intervention period followed by a steady decline over time. The modest level of initial usage may be related to the minimal instruction provided to participants on usage of the platform, however, this was intended to reflect the level of instruction provided in other publicly available websites. Platform usage rates may also be related to the levels of participant satisfaction reported which were not as high as expected given the materials and

intervention platform were developed specifically for this population. However, given the limited number of participants who completed participant satisfaction survey items these results should be interpreted cautiously. Analysis of the frequency of user login and self-monitoring entries revealed no statistically significant association with behaviour at either the three- or nine-month assessment point. While the pattern of declining participant usage over time is similar to that of previous studies [44, 53] the absence of an association with behaviour change is not [22-24, 44, 54]. The lack of association could be due to a lack of statistical power or the measures of engagement and usage applied in the current study failed to capture real participant engagement with the platform. Donkin et al., suggested that more in-depth analysis of participant engagement and platform usage is required to understand the relationship between engagement and behaviour change [24]; the findings of this study support the assertion that simplistic measures of usage may not be the strongest predictors of behaviour change. Alternatively, it may be that participants received an adequate exposure to intervention materials to promote behaviour change during their initial use of the platform. The suggestion that individuals behaviour can be changed following a single exposure to an intervention is supported by previous studies [49, 55, 56]. Several interventions now seek to maintain and increase greater engagement through various IT-based strategies to foster greater or more sustained behaviour change, and evaluation of these strategies to prolong user engagement are in their infancy [44, 57]. Yet given the consistently reported declines in platform usage and engagement over time [15, 44], and evidence of effectiveness of repeated interventions delivered in a 'booster' style [58, 59], comparing the effectiveness of these two intervention approaches to change behaviour may contribute to understanding the most effective way to change behaviour. Potential reasons for low

engagement, usage and satisfaction could be a mismatch between participants' expectations of the intervention and intervention reality. For example, process evaluation of participants in this trial revealed that many wanted prescriptive and personalized information and feedback on their progress [60]. Participants also expressed a desire to transfer the print-based intervention to online platforms to increase accessibility, and for the ManUp app to be usable without having to have an internet connection [60]. As such, it appears that participants desired an increased accessibility to content above that provided by the current intervention platform. These issues may have adversely impacted engagement and satisfaction. Managing the user expectations by providing a flexible intervention platform that is highly accessible may improve platform usage above that observed in the current trial [60].

One strategy intended to promote prolonged engagement is social interaction among participants [57]. In the current study, 10% of participants had an online friend or a 'mate' and website statistics showed that all of these participants had no more than one mate. The number of participants with online friends is slightly higher than that reported in a sub-sample of users of the 10,000 Steps website (4.3%) yet use of online friends appears to be low by participants of health behaviour change interventions [22]. Low use of this feature in the current study could be due to several factors including: a lack of awareness of this feature on the intervention platform; the fact the platform required users to search by name for a mate without knowing who else was on the platform; a reluctance to befriend individual's online when they are not 'real life' friends; and the limited number of participants on the website (n = 205) compared to other online social networking sites (e.g.. Twitter, Facebook). **The study did not assess if participants knew the identity of other participants also enrolled in**

the study. These issues should be considered in the design of future interventions seeking to implement a social support / interaction feature. Some of these restrictions were imposed to preserve the integrity of the RCT design which poses interesting design issues for future studies seeking to evaluate the effectiveness of social interaction within RCT designs. These issues include how to foster online social interaction between individuals who do not know each other in real life, or allowing study participant's real life friends to use the platform and maintain the integrity of the trial.

The ManUp challenges allowed participants to select from a range of different challenges that varied in the length of challenge and the amount of the behaviour to be performed, and analysis of the challenges selected by participants revealed some interesting results. Figure 4 shows that light strength challenges were the most frequently initiated challenges for both physical activity and dietary behaviours. This may be a function of participants seeking to try a new activity or dietary change to build confidence in their ability to change prior to committing to longer term change [25]. This aligns with behaviour change theory which indicates that successfully completing a task is useful in building confidence to complete subsequent tasks [29]. In regard to the physical activity challenges, participant data indicated that some users selected a variety of different challenge activities which may be one way users sought to introduce variety to their physical activity regime in order to maintain interest over a longer time period. The most frequently completed challenge was walking which is interesting as it has been suggested that walking doesn't appeal to males in this age range [61]. Yet this does not seem to be the case in the current study, nor does it appear to be mirror broader data for Australian males that indicates walking is the

most popular recreational activity engaged in [32]. Figure 6, shows the number of each healthy eating goal completed by challenge strength. The higher number of full strength goals completed reflects the longer duration and higher number of goals required for this challenge strength. Fastfood, alcohol and softdrink free days and consuming two pieces of fruit were the four most frequently completed goals overall which partly aligns with the objective of the healthy eating challenges to incorporate positive changes to dietary habits to improve overall dietary quality. That no healthy eating challenges were completed indicates either poor compliance with implementing changes, or that the way in which a challenge needed to be completed (specific number of goals over a specified time period) could be improved. ManUp challenges were designed to provide 'ready made' targets for males, yet some males expressed a desire to have greater flexibility in setting their own goals and recording progress in their preferred metric [60]. Adopting this approach may have increased the completion rate of the challenges.

Approximately three-quarters of participants in the IT-based arm owned a mobile phone that allowed them to access the internet and therefore the mobile component of the intervention. This is higher than previous reports in Australia, and is likely to continue to increase as ownership of internet capable mobile devices continues to increase [62]. Yet, only 21.8% of participants in the IT-based intervention who had access, actually logged in to the mobile phone platform. The low level usage of the mobile phone component may be related to its limited functionality. Limiting functionality was a conscious design decision in order to maximise potential use across a wide variety of mobile phones including some older mobile phones that did not allow the navigation and functionality of newer smartphone devices. Interviews

with intervention participants indicated that needing to be connected to the internet to use the mobile app was a limiting factor for some users and this may have also contributed to the low usage [60]. Recent growth in and expected growth of smartphone sales and ownership, self-monitoring devices (i.e. Fitbit, Jawbone) and willingness of some males to use smartphone technology [17, 62, 63] will allow future interventions to take advantage of the greater functionality offered by these devices including the ability to perform self-monitoring via installed apps without the device/app requiring an internet connection.

Health literacy allows individuals to use and apply knowledge to process information and inform decisions concerning their health and is identified as a priority for males [13]. ManUp educational materials were designed to provide clear and concise information on the rationale for changing physical activity and dietary behaviours and how to achieve these changes consistent with formative research and behaviour change theory [18, 25, 29]. Study outcomes show mixed support for the effectiveness of the materials to improve physical activity and nutritional literacy. Nutrition literacy remained unchanged throughout the study and may have been impacted by the moderately high level of nutritional literacy at baseline [41]. Dietary educational materials provided information on the amount and frequency that particular food groups should be eaten consistent with national guidelines and the benefits of consuming the particular food group and an overall healthy diet [25, 33]. Whilst the nutritional literacy scale assessed some components addressed by educational materials (e.g. required servings of fruit and vegetables, high-sugar foods contain a high number of kilojoules) other components were much broader (e.g. how to prevent food poisoning from eggs, cost and weed control methods of organic farms). As such,

the discrepancy between the educational materials provided in the intervention and the instrument used to assess nutritional literacy may have limited the ability to detect change in this construct. Alternatively, participants may not have read these educational materials or preferred a more prescriptive approach to the dietary information provided [60]. Baseline physical activity literacy was lower than previously reported, particularly regarding knowledge that blocks of a minimum of 10 minutes of activity are acceptable to improve health [64]. This was the only measure to significantly change over time with a significantly higher number of participants agreeing with this at 9 months (68.9%) compared to baseline (52.5%). Changes in the proportion of participants agreeing that moderate intensity physical activity can improve health were not statistically examined as over 95% participants agreed with this statement at nine months which resulted in insufficient variability for the analysis to take place (Table 4). Although not statistically examined, this may be viewed as an improvement in one component of physical activity literacy. The significant between-group differences in the proportion of participants agreeing that vigorous intensity physical activity performed three times per week is essential for health and that 30 minutes of moderate intensity physical activity per day can improve health are likely due to differing levels between groups at baseline. The reasons for the baseline differences between intervention groups on the proportions who agreed that 30 minutes of moderate intensity physical activity per day can improve health are unknown, particularly as qualitative research indicates that males have a good knowledge of the volumes and frequencies of physical activity needed to improve health [18, 65]. Based on these observations the ManUp education materials may have been more effective at improving physical activity literacy of participants compared to nutritional literacy.

Males are acknowledged as a hard-to-reach population in the health behaviour intervention literature and this is reflected in the low recruitment rate in this study (approximately 27 participants per month of recruitment). IT-based interventions frequently report low participant retention rates and in this study the overall retention rate at 9 months was 49.2% with a lower retention rate in the IT-based group (46.8%) compared to the print-based group (54.2%) ($P = 0.24$). Low retention rates in IT-based studies are an acknowledged issue and the level of retention in this study is comparable previous intervention studies [66-69], as such, the low retention rate may not be an issue specific to the target population. The low retention rate limited the number of observations available for analysis over time however analytic methods using all available data were employed to minimise the impact of this potential limitation. Consistent with study objectives to improve regular engagement in physical activity and healthy eating, the ManUp study did not focus on weight loss during recruitment which previous studies have promoted in their recruitment [46, 47]. This may have contributed to lower than expected recruitment rates given the prevalence of overweight and obesity in males. Due to logistical constraints it was not possible to assess usage of the print-based materials and self-monitoring behaviour of participants in this intervention arm. The lack of this usage data prohibits between group comparisons of usage and behaviour change which may contribute to better understanding the relationship between platform usage and behaviour change; this is a limitation of the study. Other limitations of the study include a reliance on self-report measures. Although a subsample of participants ($n = 91$) were provided with accelerometers to objectively measure of physical activity [25]; poor participant compliance with measurement protocols resulted in too few participants providing

valid data for meaningful analysis and this data is not reported in this paper. The usage rate of the mobile component and use and functionality aspects of the 'mate feature' are also limitations of the current study.

This study evaluated the effectiveness of an intervention delivered by IT- and print-based materials to promote self-monitoring of physical activity and dietary behaviours and health literacy of these behaviours. While study outcomes show mixed support for the intervention to change health literacy, IT- and print-based modes were effective in improving physical activity and dietary behaviours in middle aged males with no differences between delivery modes. This may suggest both may be useful intervention delivery modes.

Acknowledgements.

Queensland Health provided funding to conduct this project and to develop all intervention materials. This manuscript was partially supported by the CQUniversity Health CRN.

References

1. Artinian, N.T., et al., Interventions to promote physical activity and dietary lifestyle changes for cardiovascular risk factor reduction in adults: a scientific statement from the American Heart Association. *Circulation*, 2010. **122**(4): p. 406-41.
2. Australian Institute of Health and Welfare, Cardiovascular disease: Australian facts 2011., in Cardiovascular disease series. . 2011, AIHW: Canberra.
3. Australian Bureau of Statistics, 4364.0.55.001 - Australian Health Survey: First Results. 2012, Australian Bureau of Statistics,,: Canberra.
4. Vandelanotte, C., et al., Physical activity trends in Queensland (2002 to 2008): are women becoming more active than men? *Aust N Z J Public Health*, 2010. **34**(3): p. 248-54.
5. McNaughton, S.A., et al., An index of diet and eating patterns is a valid measure of diet quality in an Australian population. *J Nutr*, 2008. **138**(1): p. 86-93.
6. Waters, L.A., et al., Who participates in physical activity intervention trials? *J Phys Act Health*, 2011. **8**(1): p. 85-103.
7. George, E.S., et al., A review of the effectiveness of physical activity interventions for adult males. *Sports Med*, 2012. **42**(4): p. 281-300.
8. Taylor, P.J., et al., A review of the nature and effectiveness of nutrition interventions in adult males--a guide for intervention strategies. *Int J Behav Nutr Phys Act*, 2013. **10**: p. 13.
9. WHO, Achieving health equity: From root causes to fair outcomes, C.o.t.S.D.o. Health, Editor. 2007, World Health Organisation: Geneva.
10. Nutbeam, D., The evolving concept of health literacy. *Soc Sci Med*, 2008. **67**(12): p. 2072-8.
11. Australian Bureau of Statistics, Health Literacy, Australia. 4233.0. 2006, ABS: Canberra.
12. Adams, R.J., et al., Functional health literacy mediates the relationship between socio-economic status, perceptions and lifestyle behaviors related to cancer risk in an Australian population. *Patient education and counseling*, 2012.
13. Australian Government, National Male Health Policy: Building on the strengths of Australian males - healthy routines, D.o.H.a. Ageing, Editor. 2010: Canberra.
14. Norman, G.J., et al., A review of eHealth interventions for physical activity and dietary behavior change. *Am J Prev Med*, 2007. **33**(4): p. 336-345.
15. Vandelanotte, C., et al., Website-delivered physical activity interventions a review of the literature. *Am J Prev Med*, 2007. **33**(1): p. 54-64.
16. Kohl, L.F., R. Crutzen, and N.K. de Vries, Online prevention aimed at lifestyle behaviors: a systematic review of reviews. *J Med Internet Res*, 2013. **15**(7): p. e146.
17. Vandelanotte, C., et al., What Kinds of Website and Mobile Phone-Delivered Physical Activity and Nutrition Interventions Do Middle-Aged Men Want? *J Health Commun*, 2013.
18. Caperchione, C.M., et al., What a man wants: understanding the challenges and motivations to physical activity participation and healthy eating in middle-aged Australian men. *Am J Mens Health*, 2012. **6**(6): p. 453-61.
19. Morgan, P.J., et al., 12-month outcomes and process evaluation of the SHED-IT RCT: an internet-based weight loss program targeting men. *Obesity (Silver Spring)*, 2011. **19**(1): p. 142-51.
20. Webb, T.L., et al., Using the internet to promote health behavior change: a systematic review and meta-analysis of the impact of theoretical basis, use of behavior change techniques, and mode of delivery on efficacy. *J Med Internet Res*, 2010. **12**(1): p. e4.

21. Neve, M., et al., Effectiveness of web - based interventions in achieving weight loss and weight loss maintenance in overweight and obese adults: a systematic review with meta - analysis. *Obesity reviews*, 2010. **11**(4): p. 306-321.
22. Davies, C., et al., Prospective associations between intervention components and website engagement in a publicly available physical activity website: the case of 10,000 Steps Australia. *J Med Internet Res*, 2012. **14**(1): p. e4.
23. Kirwan, M., et al., Using Smartphone Technology to Monitor Physical Activity in the 10,000 Steps Program: A Matched Case Controlled Trial. *Journal of Medical Internet Research*, 2012.
24. Donkin, L., et al., Rethinking the dose-response relationship between usage and outcome in an online intervention for depression: randomized controlled trial. *J Med Internet Res*, 2013. **15**(10): p. e231.
25. Duncan, M.J., et al., Effectiveness of a website and mobile phone based physical activity and nutrition intervention for middle-aged males: trial protocol and baseline findings of the ManUp Study. *BMC Public Health*, 2012. **12**: p. 656.
26. Marcus, B.H., et al., A comparison of Internet and print-based physical activity interventions. *Arch Intern Med*, 2007. **167**(9): p. 944-9.
27. Short, C.E., et al., Efficacy of tailored-print interventions to promote physical activity: a systematic review of randomised trials. *Int J Behav Nutr Phys Act*, 2011. **8**: p. 113.
28. Sports Medicine Australia *The Sports Medicine Australia pre-exercise screening system*. http://sma.org.au/wp-content/uploads/2009/05/new_pre_screening.pdf. 2005.
29. Bandura, A., Social Cognitive Theory of self-regulation. *Organi Behav Hum Decis*, 1991. **50**: p. 248-287.
30. Maes, S. and P. Karoly, Self-Regulation assessment and intervention in physical health and illness: A review. *Applied Psychology: An international review*, 2005. **54**(2): p. 267-299.
31. Bandura, A., The Primacy of Self-Regulation in Health Promotion. *Applied Psychology*, 2005. **54**(2): p. 245-254.
32. Australian Bureau of Statistics, Participation in Sport and Physical Recreation, Australia 4177.0. 2010, Australian Bureau of Statistics: Canberra.
33. National Health & Medical Research Council, Dietary Guidelines for Australian Adults. 2003, National Health & Medical Research Council: Canberra.
34. Obesity: preventing and managing the global epidemic. Report of a WHO consultation. *World Health Organ Tech Rep Ser*, 2000. **894**: p. i-xii, 1-253.
35. Australian Institute of Health and Welfare, The active Australia survey: A guide and manual for implementation, analysis and reporting. 2003, AIHW: Canberra.
36. Brown, W., et al., Comparison of surveys used to measure physical activity. *Aust N Z J Public Health*, 2004. **28**(2): p. 128-34.
37. Brown, W.J., et al., Test-retest reliability of four physical activity measures used in population surveys. *Journal of Science and Medicine in Sport*, 2004. **7**(2): p. 205-215.
38. Timperio, A., et al., Validation of adult physical activity questions for use in Australia population surveys. 2002, Department of Health and Ageing. unpublished report: Canberra.
39. Marks, G.C., et al., Monitoring food habits in the Australian population using short questions, in National Food and Nutrition Monitoring and Surveillance Project. 2001, Commonwealth Department of Health and Aged Care.
40. Australian Bureau of Statistics, National Health Survey Questionnaire 2005. 2005, Australian Bureau of Statistics: Canberra.
41. Diamond, J.J., Development of a reliable and construct valid measure of nutritional literacy in adults. *Nutr J*, 2007. **6**: p. 5.

42. Wittes, J., Sample size calculations for randomized controlled trials. *Epidemiol Rev*, 2002. **24**(1): p. 39-53.
43. Eysenbach, G., The law of attrition. *Journal of Medical Internet Research*, 2005. **7**(1): p. e11.
44. Brindal, E., et al., Features predicting weight loss in overweight or obese participants in a web-based intervention: randomized trial. *J Med Internet Res*, 2012. **14**(6): p. e173.
45. Leahey, T.M., et al., Effect of teammates on changes in physical activity in a statewide campaign. *Prev Med*, 2010. **51**(1): p. 45-9.
46. Morgan, P.J., et al., The SHED-IT randomized controlled trial: evaluation of an Internet-based weight-loss program for men. *Obesity (Silver Spring)*, 2009. **17**(11): p. 2025-32.
47. Morgan, P.J., et al., Efficacy of a workplace-based weight loss program for overweight male shift workers: The Workplace POWER (Preventing Obesity Without Eating like a Rabbit) randomized controlled trial. *Preventive medicine*, 2011. **52**(5): p. 317-325.
48. Maruyama, C., et al., Effect of a worksite-based intervention program on metabolic parameters in middle-aged male white-collar workers: a randomized controlled trial. *Prev Med*, 2010. **51**(1): p. 11-7.
49. Parekh, S., et al., Improving diet, physical activity and other lifestyle behaviours using computer-tailored advice in general practice: a randomised controlled trial. *Int J Behav Nutr Phys Act*, 2012. **9**: p. 108.
50. King, A.C., et al., Behavioral Impacts of Sequentially versus Simultaneously Delivered Dietary Plus Physical Activity Interventions: the CALM Trial. *Ann Behav Med*, 2013. **46**(2): p. 157-68.
51. Hyman, D.J., et al., Simultaneous vs sequential counseling for multiple behavior change. *Arch Intern Med*, 2007. **167**(11): p. 1152-8.
52. Davies, C.A., et al., Meta-analysis of internet-delivered interventions to increase physical activity levels. *Int J Behav Nutr Phys Act*, 2012. **9**: p. 52.
53. Glasgow, R.E., et al., Engagement in a diabetes self-management website: usage patterns and generalizability of program use. *Journal of Medical Internet Research*, 2011. **13**(1): p. e9.
54. Funk, K.L., et al., Associations of internet website use with weight change in a long-term weight loss maintenance program. *J Med Internet Res*, 2010. **12**(3): p. e29.
55. Vandelanotte, C., I. De Bourdeaudhuij, and J. Brug, Two-year follow-up of sequential and simultaneous interactive computer-tailored interventions for increasing physical activity and decreasing fat intake. *Ann Behav Med*, 2007. **33**(2): p. 213-9.
56. Spittaels, H., I. De Bourdeaudhuij, and C. Vandelanotte, Evaluation of a website-delivered computer-tailored intervention for increasing physical activity in the general population. *Prev Med*, 2007. **44**(3): p. 209-17.
57. Kolt, G.S., et al., WALK 2.0 - Using Web 2.0 applications to promote health-related physical activity: A randomised controlled trial protocol. *BMC Public Health*, 2013. **13**(1).
58. Grandes, G., et al., Two-year longitudinal analysis of a cluster randomized trial of physical activity promotion by general practitioners. *PLoS One*, 2011. **6**(3): p. e18363.
59. Scholz, U., S. Ochsner, and A. Luszczynska, Comparing different boosters of planning interventions on changes in fat consumption in overweight and obese individuals: a randomized controlled trial. *Int J Psychol*, 2013. **48**(4): p. 604-15.
60. Short, C.E., et al., Examining participant engagement in an information technology-based physical activity and nutrition intervention for men: the manup randomized controlled trial. *JMIR Res Protoc*, 2014. **3**(1): p. e2.

61. Burton, N.W., A. Walsh, and W.J. Brown, It just doesn't speak to me: mid-aged men's reactions to '10,000 Steps a Day'. *Health Promot J Austr*, 2008. **19**(1): p. 52-9.
62. Australian Communications and Media Authority, Report 3-Smartphones and tablets Take-up and use in Australia, in Communications report 2011–12 series. 2013, Australian Communications and Media Authority,; Canberra.
63. Walker, S., *Wearable Technology - World - 2013*. 2013, IHS Electronics & Media.
64. Armstrong, T., A. Bauman, and J. Davies, Physical activity patterns of Australian adults. Results of the 1999 National Physical Activity Survey. 2000, Australian Institute of Health and Welfare: Canberra.
65. George, E.S., et al., Physical Activity and Sedentary Time: Male Perceptions in a University Work Environment. *Am J Mens Health*, 2013.
66. Buis, L., et al., Evaluating Active U: an Internet-mediated physical activity program. *BMC Public Health*, 2009. **9**(1): p. 331.
67. Kosma, M., B.J. Cardinal, and J.A. Mccubbin, A pilot study of a web-based physical activity motivational program for adults with physical disabilities. *Disability & Rehabilitation*, 2005. **27**(23): p. 1435-1442.
68. Carr, L.J., et al., Internet-delivered behavior change program increases physical activity and improves cardiometabolic disease risk factors in sedentary adults: results of a randomized controlled trial. *Preventive medicine*, 2008. **46**(5): p. 431-438.
69. McConnon, Á., et al., The Internet for weight control in an obese sample: results of a randomised controlled trial. *BMC health services research*, 2007. **7**(1): p. 206.