

Home Search Collections Journals About Contact us My IOPscience

Ab-initio validation of a simple heuristic expression for the sequential-double-ionization contribution to the double ionization of helium by ultrashort XUV pulses

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys.: Conf. Ser. 635 092039

(http://iopscience.iop.org/1742-6596/635/9/092039)

View the table of contents for this issue, or go to the journal homepage for more

Download details:

IP Address: 129.130.37.147

This content was downloaded on 24/03/2016 at 15:00

Please note that terms and conditions apply.

## Ab-initio validation of a simple heuristic expression for the sequential-double-ionization contribution to the double ionization of helium by ultrashort XUV pulses

## Aihua Liu <sup>1</sup> and Uwe Thumm <sup>2</sup>

Department of Physics, Kansas State University, Manhattan, Kansas 66506, USA

Synopsis We study two-photon double ionization of helium by short XUV pulses by numerically solving the time-dependent Schrödinger equation in full dimensionality within a finite-element discrete-variable-representation scheme. Based on the emission asymmetries in joint photoelectron angular distributions, we identify sequential and non-sequential contributions to two-photon double ionization for ultrashort pulses whose spectrum overlaps the sequential ( $\hbar\omega > 54.4 \text{ eV}$ ) and non-sequential ( $39.5 \text{ eV} < \hbar\omega < 54.4 \text{ eV}$ ) double-ionization regimes.

For sufficiently long extreme ultraviolet (XUV) pulses, sequential and non-sequential two-photon double ionization of helium atoms are distinguishable by the central XUV photon energy. If this energy is above the second ionization potential (54.4 eV), emission proceeds sequentially, otherwise non-sequential correlated double ionization occurs [1]. For ultrashort XUV pulses, this distinction breaks down if their spectrum overlaps the sequential and non-sequential regimes [2].

Solving the time-dependent Schrödinger equation (TDSE) ab initio, we investigate joint photoelectron angular distributions (JADs) for two-photon double ionization of helium atoms in the sequential and non-sequential regimes. We find that strong electronic correlation results in similar angular distributions over a large range of energy sharings between the two photoelectrons [3]. Based on the emission asymmetry in JADs (ranging from -1 for emission in opposite to 1 for emission into the same hemisphere), we distinguish "sequential" and "non-sequential" contributions to double ionization by linearly polarized ultrashort XUV pulses.

Figure 1 displays the "effective sequential double-ionization (SDI) yield" and double-ionization emission-asymmetry dependence on the XUV-pulse duration. We define the effective sequential yield as the product of the effective interaction time (pulse duration) and the integrated XUV-pulse-spectral intensity for  $\hbar\omega > 54.4$  eV (blue area in the inset of Fig. 1). For XUV pulses with a central photon energy of 50 eV, the SDI contribution reaches a maximum value at a pulse duration of 650 as, which we ex-

plain as being due to the competition between increasing temporal and decreasing spectral pulse widths.




Figure 1. Effective SDI yield for helium in a 10<sup>14</sup> W/cm<sup>2</sup> peak intensity XUV pulse with a central energy of 50 eV as a function of the XUV pulse duration. The stars are results of our *ab initio* TDSE calculations [1,3] for the double-emission asymmetry at extremely unequal energy sharing of the emitted electrons. The inset displays a typical XUV-spectral intensity, the blue area indicating the spectrum above the second ionization potential of helium.

This work is supported by the U.S. NSF and the Division of Chemical Sciences, Office of Basic Energy Sciences, Office of Energy Research, U.S. DoE.

## References

- [1] A. Liu and U. Thumm 2014 Phys. Rev. A 89 063423.
- [2] J. Feist et al. 2009 Phys. Rev. Lett. 103 063002.
- [3] A. Liu and U. Thumm 2015, *Phys. Rev. A*, submitted for publication.

<sup>1</sup>E-mail: iwaa@phys.ksu.edu <sup>2</sup>E-mail: thumm@phys.ksu.edu

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.