Sorghum Grain As the Only Protein Source In All-Concentrate Heifer Finishing Rations: Two Levels of Urea In An All-Concentrate Ration, (Project 253-6) 1968

E. F. Smith, D. Richardson, C. L. Drake and B. E. Brent.

Tests reporting on sorghum grain as the only protein source in a diet for finishing cattle are reported in Kansas Agr. Expt. Station Bulletins 493 and 507 and on page 24 in this bulletin.

The three rations fed are shown in table 10. The sorghum grain was obtained as needed at a local elevator, mixed with other ration ingredients at the Animal Husbandry elevator and delivered, usually in 2000 lb. loads to the self feeders when needed.

The heifers were purchased by an order buyer through sale barns and were mostly Angus and Angus-Hereford crossbreds. They were started on feed with a mixture of 60% dehydrated alfalfa crumbles and 40% ground sorghum grain fed in a self-feeder with prairie hay free choice. During two weeks the alfalfa crumbles were gradually reduced. A premix was included when alfalfa crumbles were less than 25 percent of the ration. After the heifers were on all concentrate, the prairie hay was gradually removed from the ration during one week. At the end of three weeks the heifers were on all concentrate ration. Then the following three rations were compared:

1. All sorghum grain without added protein, 2. Sorghum grain plus one percent urea, and 3. Sorghum grain plus 2 percent urea.

Results

The diet containing 2% urea reduced rate of gain and increased costs. It seemed to reduce feed intake in one of the two lots, table 11. Least feed per pound of gain was from the ration containing 1% urea. The heifers fed sorghum grain as the only protein source required about a half pound more feed to produce a pound of gain than those feed 1% urea.

No differences in carcasses could be explained by ration treatments. The reason for the low dressing percentage (58) for lot 1 is unknown. Carcass grades averaged low choice to choice for the different lots.

This test substantiates previous tests that excellent results may be obtained with all concentrate rations where the source of protein is sorghum grain with 1 percent urea added. Nearly as good results may be obtained by omitting the urea.

Table 10
Composition of Rations, 1968

	Sorghum	grain and	Sorghum grain and	
	grain	1% urea	2% urea	
In	gredients per ton			
l arain	1945	1925	1905	
round sorghum grain remix	50	50	50	
alt	5	5	5	
rea (45% nitrogen)	0	20	40	
Total, lb.:	2000	2000	2000	
round limestone race mineral premix ^l	20) L.0		
tilbestrol premix (2 grams				
stilbestrol per lb.)		0.5		
itamin A premix (10,000				
IU per gram)		0.3 (140 gms.)		
Chlortetracycline premix		0.8 (380 gms.)		
(10 grams per lb.)		0.8 (300 gms.)		
Ground sorghum grain				
Jedana SorBumm Samm				
(enough to make the premix up to 50 lbs.)	2	7.4		

Percentages of indicated elements in trace mineral premix: Manganese, 4.4; iron, 6.6; copper, 1.32; cobalt, 0.23; iodine, 0.30; zinc, 5; Magnesium, 20; sulfur, 2.70.

Sorghum Grain With No Added Protein In a Cattle Finishing Ration; Two Levels of Urea In An All Concentrate Ration, October 23, 1967, to January 25, 1968 -

Table 11

Lot number	Sorghum grain		Sorghum grain and 1% urea		Sorghum grain and 2% urea	
	1 .	2	3	4	5	6
No. of heifers	,					
per lot	11	11	11	11	11	11
Av. initial wt.						
lbs.	632	628	628	625	629	(00
Av. daily gain			3.20	023	029	620
IDS.	2.60	2.61	2.62	2.66	2.36	2.08
Av. daily feed						
intake, lbs.	21.0	20.4	19.6	19.6	19.6	17.6
Feed required per lb. of					27.0	17.0
gain, 1bs.	8.1	8.0	7.5	7 /		
Feed cost per		0.0	7.5	7.4	8.3	8.5
cwt. of gain, 1 \$	14.84	14.33	14.11	13.79	15.86	16.22
Carcass data:		t .				
Av. Carcass wt.	508	518	524	519	507	502
Dressing Percent	58.0	59.3	59.9	59.3	59 . 5	503 59.7
Carcass grade ² Rib eye area,	19.5	19.6	19.6	19.6	20.0	19.5
sq. in.	10.3	10 /				
Fat thickness	10.3	10.4	10.3	10.2	10.7	9.73
inches	.43	.45	•47	.49	.53	.48
Yield grade ³	2.8	3.1	3.0	2.7	3.0	3.1
ercent protein in						
concentrate mix-						
ture (88% dry mat-						
ter basis).	9.74	9.95	12.60	12.54	14.82	14.99
ost of concentrate						
mixture/ton1	\$36.79	\$36.79	\$37.53	\$37.53	\$38.27	

¹ Feed prices are on inside back cover.
2 Carcass grade score: Low choice, 19; Average choice, 20.
3 Yield grade: Ranges from 1 to 5, 1 is most desirable.