

**Open Access** 

# Weight Control, Endocrine Hormones, and Cancer Prevention

### Weigun Wang\*

Department of Human Nutrition, Kansas State University, Manhattan, KS 66506, USA

Obesity is well recognized to increase the risk of several types of cancer [1]. It has been suggested that those of 25% over normal weight have a 33% greater cancer risk than those who maintain a normal body weight [2]. There is ample evidence that weight control via decreasing calorie intake and/or increasing physical activity reduces cancer risk. For almost a century, Dietary Calorie Restriction (DCR) has been shown to inhibit the development of almost all types of cancer in animal models [3]. Alternatively, exercise has also been shown to reduce some types of cancer [4]. When compared with DCR, however, the impact of exercise may demonstrate a modest but intricate result for both weight maintenance and cancer prevention. Our previous studies conducted in an animal model, for example, showed that treadmill exercise alone with ad libitum feeding was not sufficient to decrease body weight due to, at least in part, the corresponding increase in dietary intake [5]. If the food intake of the exercised mice was limited by iso-caloric intake with sedentary counterpart, then body weight and body fat were significantly reduced [6]. Indeed, a negative calorie homeostasis appears to be the most potent for preventing cancer, which may be even extrapolated from the greatest of weight control diets to many dietary cancer preventive factors such as whole grain, fiber, and resistant starch, as well as vegetables and/or fruits, etc.

The mechanism by which weight control may reduce cancer risk is not well defined. Recently the National Cancer Institute called for research proposals regarding 24 Provocative Questions (http:// provocativequestions.nci.nih.gov). The number one question is how obesity contributes to cancer risk. Previous studies by us and others have found that weight loss is associated with a reduction of many plasma levels of hormones such as IGF-1, leptin, insulin, and thyroid hormones [7,8]. It is possible that the reduction of these hormones accounts for cancer prevention by weight loss due to a mitogenic role, especially by IGF-1 and leptin, in promoting cellular growth and metabolic complications [9].

As a mitogen, IGF-1 is well known to stimulate cell proliferation, inhibit apoptosis, and enhance angiogenesis. Chronically high levels of IGF-1 observed in obese subjects have been linked with an increased risk of developing several types of cancer in case-control and prospective cohort studies [10,11]. It has been suggested that the role of IGF-1 in mitogenesis is triggered through cellular signaling cascades leading to the activation of both IGF-1 dependent MAPKproliferation and PI3K-anti-apoptosis pathways for cancer promotion [12]. Furthermore, the role of adipocytes on malignance has been suggested via tumor-stroma interaction required during metastases or early micrometastases [13]. Recent data also suggest that adipocytes not only respond to hormonal signals, but also produce hormone-like factors such as pro-inflammatory leptin that may enhance oncological risk [14]. Although it is not clear how leptin interacts in IGF-1 mitogenic signaling, our previous studies conducted in TPA-promoted mouse skin cancer model have demonstrated that the reduction of both IGF-1 and leptin levels in response to weight loss corresponded to a concomitant inhibition of IGF-1-dependent mitogenic cascades and thus diminution of TPA-promoted signaling [5,6,15].

In addition to fundamentally advancing in the fields of obesity

and cancer research, better understanding of the relationship between weight control, endocrine hormones, and hormone-dependent signal pathways may lead to a novel approach for cancer prevention.

#### References

- 1. Bianchini F, Kaaks R, Vainio H (2002) Overweight, obesity, and cancer risk. Lancet Oncol 3: 565-574.
- 2. International Agency for Research on Cancer (2002) IARC handbooks of
- 3. Hursting SD, Smith SM, Lashinger LM, Harvey AE, Perkins SN (2010) Calories and carcinogenesis: lessons learned from 30 years of calorie restriction research. Carcinogenesis 31: 83-89.
- 4. Anzuini F, Battistella A, Izzotti A (2011) Physical activity and cancer prevention: a review of current evidence and biological mechanisms. J Prev Med Hyg 52:
- 5. Xie L, Jiang Y, Ouyang P, Chen J, Doan H, et al. (2007) Effects of Dietary Calorie Restriction or Exercise on the PI3K and Ras Signaling Pathways in the Skin of Mice. J Biol Chem 282: 28025-28035.
- Ouyang P, Jiang Y, Doan HM, Xie L, Vasquez D, et al. (2010) Weight Loss via Exercise with Controlled Dietary Intake May Affect Phospholipid Profile for Cancer Prevention in Murine Skin Tissues. Cancer Prev Res 3: 466-477.
- Pardina E, Ferrer R, Baena-Fustegueras JA, Lecube A, Fort JM, et al. (2010) The relationships between IGF-1 and CRP, NO, leptin, and adiponectin during weight loss in the morbidly obese. Obes Surg 20: 623-632.
- 8. Fouladiun M, Körner U, Bosaeus I, Daneryd P, Hyltander A, et al. (2005) Body composition and time course changes in regional distribution of fat and lean tissue in unselected cancer patients on palliative care--correlations with food intake, metabolism, exercise capacity, and hormones. Cancer 103: 2189-2198.
- Jiang Y, Wang W (2008) Potential mechanisms of cancer prevention by weight control, Biophy Rev Lett 3: 421-437.
- 10. Toriola AT, Surcel HM, Lundin E, Schock H, Grankvist K, et al. (2011) Insulinlike growth factor-I and C-reactive protein during pregnancy and maternal risk of non-epithelial ovarian cancer: a nested case-control study. Cancer Causes Control 22: 1607-1611.
- 11. Rohrmann S, Linseisen J, Becker S, Allen N, Schlehofer B, et al. (2011) Concentrations of IGF-I and IGFBP-3 and brain tumor risk in the European Prospective Investigation into Cancer and Nutrition. Cancer Epidemiol Biomarkers Prev 20: 2174-2182.
- 12. Pollak M (2012) The insulin and insulin-like growth factor receptor family in neoplasia: an update. Nat Rev Cancer 12: 159-169.
- 13. Gnacińska M, Małgorzewicz S, Stojek M, Łysiak-Szydłowska W, Sworczak K (2009) Role of adipokines in complications related to obesity: a review. Adv Med Sci 54: 150-157.

\*Corresponding author: Weiqun Wang, Department of Human Nutrition, Kansas State University, Manhattan, KS 66506, USA, Tel: 1-785-532-0153; E-mail:

Received April 30, 2012; Accepted May 03, 2012; Published May 07, 2012

Citation: Wang W (2012) Weight Control, Endocrine Hormones, and Cancer Prevention. J Nutr Food Sci 2:e107. doi:10.4172/2155-9600.1000e107

Copyright: © 2012 Wang W. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

- 14. Wrann CD, Laue T, Hübner L, Kuhlmann S, Jacobs R, et al. (2012) Shortterm and long-term leptin exposure differentially affect human natural killer cell immune functions. Am J Physiol Endocrinol Metab 302: E108-E116.
- 15. Lu J, Xie L, Sylvester J, Wang J, Bai J, et al. (2007) Different gene expression of skin tissues between mice with weight controlled by either calorie restriction or physical exercise. Exp Biol Med (Maywood) 232: 473-480.

## Submit your next manuscript and get advantages of OMICS **Group submissions**

### Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper Digital articles to share and explore

### Special features:

- 200 Open Access Journals 15,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing Indexing at PubMed (partial), Scopus, DOAJ, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
  Authors, Reviewers and Editors rewarded with online Scientific Credits
  Better discount for your subsequent articles

Submit your manuscript at: http://www.editorialmanager.com/lifesciences