

2006

Kansas Performance Tests with Alfalfa Varieties

Report of Progress 971




Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service

K-STATE AGRONOMY

Centennial

1906

2006



● dryland

★ irrigated

TABLE OF CONTENTS

2006 Performance Tests

Objectives and Procedures.....	1
Variety Characterization.....	1
Northeast, Topeka, Shawnee County, Seeded 2003	Table 1.....2
Southeast, Mound Valley, Labette County, Seeded 2005	Table 2.....3
North Central, Belleville, Republic County, Seeded 2004	Table 3.....4
South Central, Hutchinson, Reno County, Seeded 2004	Table 4.....5
Northwest Irrigated, Colby, Thomas County, Seeded 2003	Table 5.....6
Southwest Irrigated, Garden City, Finney County, Seeded 2002	Table 6.....7
2006 Entries With Disease and Insect Resistance Ratings for Released Varieties	Table 7.....8
Electronic Access and University Research Policy.....	back cover

Entrants in 2006 Kansas Alfalfa Performance Tests.

ABI Alfalfa (ABI) Ames, IA 515-292-2432 abialfalfa.com	Dairyland Research (Dairyland) West Bend, WI 608-676-2237 dairylandseed.com	J.R. Simplot Company (Simplot) Boise, ID 208-672-2732 simplot.com	Power Seeds, Inc. (Power) Fraserville, Ontario Can 705-944-5600
AgriPro Seed (AgriPro) Slater, IA 877-247-4776 agripro.com	Forage Genetics Inc (Croplan/NK) Boone, IA 515-432-9115	Johnston Seed Co (Johnston) Enid, OK 580-233-5800	Sharp Bros. Seed Company (Sharp) Healy, KS 800-462-8483 sharpseed.com
Allied Seed (Allied) Macon, MO 660-385-6690 alliedseed.com	Foundation Seed Division (NE AES & USDA) Lincoln, NE 877-229-1363 402-624-8038	KSU - Foundation Seed (KS AES & USDA) Manhattan, KS 785-532-6115	Star Seed, Inc. (Star) Osborne, KS 785-346-5447 gostarseed.com
Bio-Plant Research (Bio-Plant) Camp Point, IL 800-593-7708	Garst Seed Co. (Garst) Greensburg, KS 620-723-2454 garstseed.com	Monsanto Seed (Monsanto) St. Louis, MO 800-335-2676	Syngenta Seeds, Inc. (NK) Golden Valley, MN 763-593-7324 nk-us.com
Cal/West Seeds (Cal/West) West Salem, WI 608-786-1554	Great Plains Research Co. (Cimarron USA) Apex, NC 800-874-7945 CimarronUSA.com	Indianapolis, IN 317-337-7568	Taylor Seed Farms, Inc. (Taylor) White Cloud, KS 800-742-7473 taylorseedfarms.com
Channel Bio Corp. (Midwest Seed) Kentland, IN 800-369-8218 channelbio.com	Hytest Seeds (Hytest) Fort Dodge, IA 717-737-4529	NC+ Hybrids (NC+) Lincoln, NE 800-279-7999 nc-plus.com	W-L Research, Inc. (W-L) Madison, WI 608-240-0630
CroPlan Genetics (CroPlan Genetics) St. Paul, MN 800-851-8810	J.C. Robinson Seed Co. (Golden Harvest) Waterloo, NE 800-228-9906 goldenharvestseeds.com	PGI Alfalfa, Inc. (PGI) Woodland, CA 866-744-5710	Pioneer Hi-Bred, Intl., Inc. (Pioneer Brand) Johnston, IA 8800-247-6803

2006 PERFORMANCE TESTS

Objectives and Procedures

The Kansas Agricultural Experiment Station established an official alfalfa testing program in 1980 to provide Kansas growers with unbiased performance comparisons of alfalfa varieties marketed in the state. Each year, private companies are asked to enter varieties voluntarily at the locations slated for establishment that year. Announcements and entry forms are mailed to private companies in June for entry in fall-seeded tests. Companies enter varieties of their choice and pay entry fees to cover part of the costs of conducting the tests. Most tests are planted in mid-August or September, but the Southeast Kansas test usually is planted in the spring. Individual tests are conducted for a minimum of 3 years. New tests typically are established during the final production year of the previous test, or more frequently if there is enough interest.

Descriptive information is presented with the results for each test. This information, including soil type, establishment methods, fertilization, pest control, irrigation, harvest dates, and growing conditions unique to that location, can help explain test and/or variety performance.

Forage yields were estimated by harvesting four replications of each variety with a plot harvester. The amount of forage produced from a specific area (35-80 ft²) was weighed, and a subsample was taken to determine moisture content. This information was used to convert the plot weights to tons of dry matter per acre for each cutting, the season total, and the total for each previous season, as presented in Tables 1 through 8. The forage yield over the lifetime of a particular test is presented as the total tons of dry matter produced per acre, as the total tons of 15% moisture hay, and as a percentage of the test average.

Each table is separated into three sections. The first lists released cultivars that are generally available on the seed market or soon will be. The second section includes experimental cultivars that were entered in the test before being released for sale. These experimental lines often represent an earlier generation of seed than that used for the released cultivars. The third section includes summary statistics unique to that test.

At the bottom of each column, the Least Significant Difference (LSD) is listed at the 0.05 and 0.20 levels. These values indicate how large a difference is needed to be confident that one variety is superior to another. Differences between varieties that are equal to or greater than the 0.05 LSD have only a 1 in 20 chance of being due to chance or error. Differences equal to or greater than the 0.20 LSD have a 1 in 5 chance of being caused by chance or error.

The Coefficient of Variability (CV) provides an estimate of the consistency of the results of a particular test. In these tests, CVs less than 10% generally indicate reliable, uniform data, whereas CVs of 10 to 15% are not uncommon and generally indicate that the data are acceptable for rough comparisons. Tests with CVs greater than 15% may still be useful, but variety comparisons lack precision.

The Mean Coefficient of Variability (MCV) is similar to the CV in that it serves as an indicator of test precision. The MCV is calculated by dividing the 0.05 LSD by the test mean (average) and multiplying by 100. The MCV reveals the percentage difference required to detect differences between varieties with 95% confidence. Many alfalfa breeders and testers agree that tests with MCV values greater than 10% are of little benefit.

Variety Characterization

For variety selection, producers should consider the performance of a variety in each of the current tests in which it appears, its performance over time and locations relative to familiar or check varieties, and the disease and insect resistance characteristics that are potentially important in their situation.

Tables 1 through 6 contain updated yield data from individual tests currently in progress. First-season yields for a spring-planted test are often more variable than yields in subsequent years. Season totals are important, but yield distribution during the season may differ among varieties. Examine yields from individual cuttings to determine if differences in yield distribution exist. Yield totals over many years provide the best measure of variety performance over time.

Table 7 provides winter survival, disease and insect-resistance, multi-foliolate expression, and continuous grazing tolerance ratings for released varieties. These ratings were obtained primarily from the annual "Winter Survival, Fall Dormancy & Pest Resistance Ratings for Alfalfa Varieties" pamphlet published by the National Alfalfa Alliance. That report summarizes information submitted by developers of alfalfa varieties as part of the variety registration process. The Association of Official Seed Certifying Agencies (AOSCA) National Alfalfa Variety Review Board (NAVRB) reviewed the ratings before they were published. Companies submitting varieties for the tests provided ratings for some unregistered varieties. Experimental varieties are also listed in Table 7 for brand identification.

Table 1. Northeast Kansas, Topeka Alfalfa Performance Test, Seeded September 9, 2003.

Kansas River Valley Experiment Field, Eudora silt loam
 18 lb. seed/acre
 Plots 3'x20'; 3'x20' harvested
 36-92-216 lb/a of N-P-K before planting

Some yield reduction because of gopher infestations. Very little regrowth between 2nd and 3rd cuttings from the dry weather.

NAME	Forage Yield tons/acre								Total, 15% Moist.	Total, % of Mean		
	Dry Matter				2006	2005	2004	Total				
	5-17	6-20	7-25	9-28								
RELEASED CULTIVARS												
Notice II	1.91	1.57	1.23	1.39		6.11	6.77	7.18	20.06	23.60	104	
Perry	1.89	1.46	1.07	1.30		5.73	6.91	6.98	19.62	23.08	102	
6400HT	1.88	1.30	1.15	1.38		5.70	6.70	7.10	19.50	22.94	102	
Reward II	1.91	1.52	1.22	1.28		5.93	6.75	6.75	19.43	22.85	101	
HybriForce-420/wet	2.25	1.48	1.20	0.94		5.87	6.68	6.70	19.25	22.64	100	
Kanza	1.73	1.43	1.21	1.39		5.77	6.70	6.67	19.14	22.51	100	
Phirst	1.87	1.16	1.11	1.41		5.55	6.67	6.86	19.08	22.45	99	
Journey 204 Hybrid Alfalfa	1.81	1.48	1.08	1.18		5.54	6.83	6.70	19.07	22.44	99	
6530	1.83	1.48	1.00	1.27		5.58	6.63	6.76	18.98	22.33	99	
Power 4.2	1.77	1.23	1.17	0.89		5.07	6.69	7.21	18.97	22.31	99	
WL 357 HQ	2.02	1.43	1.06	1.31		5.82	6.41	6.71	18.94	22.28	99	
4A421	1.63	1.36	1.26	0.85		5.11	6.66	6.90	18.67	21.96	97	
SUMMARY STATISTICS												
Average	1.88	1.41	1.15	1.22		5.65	6.70	6.85	19.20	22.59	100	
LSD (0.05)	0.45	0.58	0.42	0.51		0.99	0.39	0.38	1.13	1.33	6	
LSD(0.20)	0.29	0.37	0.27	0.33		0.63	0.25	0.24	0.73	0.85	4	
CV (%)	16.68	28.55	25.54	29.04		12.14	4.09	3.86	4.09	4.09	4	
MCV(%)	24.00	41.07	36.75	41.78		17.47	5.89	5.55	5.88	5.88	6	

Table 2. Southeast Kansas, Mound Valley Alfalfa Performance Test, Seeded April 14, 2005.

Southeast Ag. Research Center, Parsons silty clay loam
 15 lb. seed/acre
 Plots 5'x20'; 3'x20' harvested
 20-50-200 lb/a of N-P-K before planting

Dry, dry, dry. Very little weevil damage observed.

NAME	Forage Yield tons/acre						Total, 15% Moist.	Total, % of Mean		
	Dry Matter			2006	2005	Total				
	2006	2005	Total							
RELEASED CULTIVARS										
FSG505	1.20	0.64	0.29		2.13	4.51	6.64	7.81	108	
Good as Gold II	1.20	0.59	0.30		2.08	4.46	6.54	7.69	107	
Kanza	1.10	0.51	0.27		1.89	4.46	6.35	7.47	104	
WL 357 HQ	1.11	0.50	0.29		1.89	4.28	6.18	7.27	101	
FSG408DP	1.13	0.50	0.24		1.87	4.27	6.14	7.23	100	
Perry	1.22	0.51	0.22		1.95	4.13	6.08	7.16	99	
6420	1.15	0.53	0.27		1.95	4.09	6.04	7.11	99	
6530	1.14	0.53	0.21		1.88	4.14	6.02	7.08	98	
Cimarron VL400	1.16	0.47	0.19		1.83	4.19	6.01	7.07	98	
Integrity	1.15	0.45	0.22		1.82	3.89	5.71	6.71	93	
EXPERIMENTAL STRAINS										
AA112E	1.17	0.55	0.23		1.95	4.21	6.16	7.25	101	
CW 15030	1.13	0.59	0.27		2.00	4.14	6.13	7.21	100	
AA108E	1.11	0.42	0.21		1.73	3.88	5.62	6.61	92	
SUMMARY STATISTICS										
Average	1.15	0.52	0.25		1.92	4.20	6.12	7.21	100	
LSD (0.05)	0.12	0.15	0.06		0.20	0.33	0.39	0.46	6	
LSD(0.20)	0.08	0.09	0.04		0.13	0.21	0.25	0.29	4	
CV (%)	7.14	19.46	15.89		7.11	5.53	4.40	4.40	4	
MCV(%)	10.25	27.91	22.78		10.20	7.93	6.32	6.32	6	

Table 3. North Central Kansas, Belleville Alfalfa Performance Test, Seeded September 1, 2004.

North Central Kansas Exp. Field, Crete silt loam
 20 lb. seed/acre
 Plots 5'x15'; 3'x15' harvested
 11-50-0 lb/a of N-P-K applied in February and after first cutting

Very little regrowth between 2nd and 3rd cuttings because of dry weather, but yields improved after late August rains.

NAME	Forage Yield tons/acre							
	Dry Matter				2006	2005	Total	Total, 15% Moist.
	2006	5-11	6-12	7-28	9-12			
RELEASED CULTIVARS								
Good as Gold II	2.01	1.01	0.61	1.30		4.92	6.83	11.75
DKA42-15	2.00	0.98	0.54	1.21		4.73	6.73	11.46
WL 335 HQ	1.95	1.05	0.56	1.14		4.70	6.67	11.37
Reward II	2.03	1.00	0.45	1.22		4.71	6.66	11.36
6415	1.89	1.10	0.49	1.06		4.54	6.75	11.28
Pioneer 54V46	1.92	1.06	0.52	1.02		4.51	6.68	11.19
HybriForce-420/wet	1.97	1.01	0.42	1.01		4.41	6.67	11.08
Genoa	1.85	1.14	0.49	1.04		4.53	6.46	10.99
6400HT	1.96	0.96	0.40	1.00		4.31	6.65	10.96
Kanza	1.80	1.07	0.53	1.12		4.52	6.24	10.76
DKA50-18	1.82	1.10	0.45	1.00		4.37	6.22	10.59
WL 357 HQ	1.73	0.96	0.40	1.03		4.11	6.12	10.23
EXPERIMENTAL STRAINS								
DS362HY	1.98	0.98	0.45	1.10		4.50	6.47	10.97
DS361HY	2.06	0.96	0.43	1.11		4.56	6.34	10.90
DS416	2.04	0.88	0.35	0.97		4.23	6.58	10.81
DS415	1.91	0.76	0.43	1.16		4.26	6.52	10.78
SUMMARY STATISTICS								
Average	1.93	1.00	0.47	1.09		4.49	6.54	11.03
LSD (0.05)	0.14	0.18	0.12	0.18		0.31	0.30	0.43
LSD(0.20)	0.09	0.12	1.69	0.11		0.20	0.19	0.28
CV (%)	5.11	12.57	17.85	11.27		4.85	3.23	2.75
MCV(%)	7.28	17.90	25.43	16.05		6.91	4.60	3.92

Table 4. South Central Kansas, Hutchinson Alfalfa Performance Test, Seeded September 1, 2004.

South Central Experiment Field, Ost silt loam
 10 lb. seed/acre
 Plots 5'x24', 3'x18' harvested
 75-40-0 lb/a of N-P-K before planting

Extended hot and dry weather greatly reduced yields.

NAME	Forage Yield tons/acre						Total, 15% Moist.	Total, % of Mean		
	Dry Matter		2006	2005	Total					
	5-12	6-8								
RELEASED CULTIVARS										
Good as Gold II	0.51	0.55		1.06	4.53	5.59	6.57	115		
WL 335 HQ	0.41	0.65		1.06	4.16	5.22	6.15	107		
FSG408DP	0.43	0.55		0.98	4.21	5.19	6.11	107		
6400HT	0.36	0.48		0.84	4.26	5.09	5.99	105		
Jade III	0.37	0.41		0.78	4.17	4.95	5.83	102		
6420	0.42	0.44		0.86	4.08	4.94	5.81	101		
Perry	0.38	0.53		0.91	4.00	4.91	5.78	101		
DKA50-18	0.34	0.50		0.84	3.96	4.80	5.65	98		
FSG406	0.39	0.41		0.80	3.99	4.79	5.63	98		
Kanza	0.40	0.35		0.75	4.00	4.75	5.59	97		
WL 357 HQ	0.26	0.39		0.65	4.06	4.72	5.55	97		
DKA42-15	0.35	0.47		0.83	3.87	4.70	5.53	96		
Genoa	0.34	0.37		0.71	3.92	4.64	5.45	95		
HybriForce-420/wet	0.34	0.33		0.67	3.91	4.58	5.39	94		
FSG505	0.32	0.41		0.74	3.80	4.53	5.33	93		
FSG351	0.30	0.38		0.68	3.73	4.41	5.19	90		
EXPERIMENTAL STRAINS										
405	0.49	0.57		1.05	4.45	5.50	6.47	113		
CW 15030	0.38	0.61		0.99	3.98	4.97	5.84	102		
404	0.39	0.32		0.71	4.17	4.88	5.74	100		
407	0.39	0.42		0.81	3.82	4.63	5.44	95		
406	0.39	0.45		0.84	3.73	4.58	5.38	94		
SUMMARY STATISTICS										
Average	0.38	0.46		0.84	4.04	4.87	5.73	100		
LSD (0.05)	0.13	0.24		0.27	0.37	0.46	0.54	9		
LSD(0.20)	0.08	0.16		0.18	0.24	0.30	0.35	6		
CV (%)	24.10	37.62		23.25	6.52	6.71	6.71	7		
MCV(%)	34.08	53.21		32.89	9.21	9.49	9.49	9		

Table 5. Northwest Kansas, Colby Alfalfa Performance Test, Seeded August 29, 2003.

Northwest Research-Extension Center, Keith silt loam
 18 lb. seed/acre
 Plots 3'x20'; 3'x17' harvested
 16-55-0 lb/a of N-P-K before planting

Hot, dry weather from the end of May until the end of August slowed growth and reduced yields.

NAME	Forage Yield tons/acre								Total, 15% Moist.	Total, % of Mean		
	Dry Matter				2006	2005	2004	Total				
	2006	5-26	6-26	7-24								
RELEASED CULTIVARS												
Pioneer 54V46	3.20	1.43	1.54	1.15		7.32	9.64	8.17	25.13	29.56	105	
Kanza	3.07	1.55	1.66	1.32		7.61	9.47	7.77	24.85	29.24	104	
FSG505	2.83	1.52	1.56	1.35		7.27	9.69	7.84	24.80	29.17	103	
Phirst	2.75	1.25	1.70	1.06		6.75	9.60	8.22	24.57	28.90	102	
WL 357 HQ	2.82	1.44	1.74	1.22		7.22	9.41	7.93	24.55	28.89	102	
Expedition	3.27	1.38	1.56	1.34		7.54	9.11	7.88	24.53	28.86	102	
Jade III	3.34	1.29	1.52	1.08		7.23	9.38	7.69	24.30	28.59	101	
Pioneer 54Q25	2.86	1.34	1.60	1.04		6.84	9.29	8.17	24.30	28.59	101	
FSG406	3.08	1.31	1.64	1.08		7.10	9.41	7.77	24.28	28.57	101	
631	3.00	1.25	1.49	1.12		6.86	9.96	7.43	24.26	28.54	101	
Arapaho	2.72	1.29	1.54	1.05		6.60	9.41	8.00	24.01	28.24	100	
HybriForce-420/wet	3.26	1.23	1.47	1.13		7.08	9.08	7.65	23.81	28.01	99	
Abundance	3.00	1.28	1.47	1.10		6.85	9.28	7.65	23.78	27.97	99	
FSG351	2.62	1.32	1.50	1.09		6.54	9.25	7.96	23.74	27.93	99	
Journey 204 Hybrid Alfalfa	3.07	1.19	1.47	1.01		6.73	9.18	7.83	23.74	27.93	99	
Notice II	2.69	1.40	1.57	1.14		6.79	8.88	7.96	23.63	27.80	99	
Regal	2.62	1.21	1.44	1.02		6.28	9.30	7.82	23.40	27.53	98	
6400HT	2.53	1.21	1.41	0.98		6.13	9.44	7.69	23.26	27.37	97	
Maximizer	2.70	1.25	1.48	1.09		6.53	9.12	7.48	23.13	27.21	96	
Perry	2.78	1.23	1.38	1.04		6.43	9.16	7.44	23.04	27.10	96	
Evermore	2.49	1.13	1.52	1.06		6.19	9.25	7.44	22.89	26.93	95	
EXPERIMENTAL STRAINS												
CL2000	2.84	1.32	1.39	1.09		6.65	9.10	7.62	23.36	27.49	97	
SUMMARY STATISTICS												
Average	2.89	1.31	1.53	1.12		6.84	9.34	7.79	23.97	28.20	100	
LSD (0.05)	0.70	0.22	0.23	0.23		0.81	0.60	0.77	1.27	1.49	5	
LSD(0.20)	0.46	0.14	0.15	0.15		0.52	0.39	0.84	0.82	0.97	3	
CV (%)	17.26	12.09	10.73	14.26		8.34	4.56	6.99	3.74	3.74	4	
MCV(%)	24.39	17.08	15.17	20.16		11.79	6.44	9.87	5.28	5.28	5	

Table 6. Southwest Kansas, Garden City Alfalfa Performance Test, Seeded September 3, 2002.

Southwest Research-Extension Center, Keith silt loam
 20 lb. seed/acre
 Plots 3'x20'; 3'x20' harvested
 22-104-0 lb/a of N-P-K before planting

Very dry winter from January through late March.
 Optimum growth for the fifth cutting was delayed by cooler-than-normal temperatures. Some gopher damage reported.

NAME	Forage Yield tons/acre										
	Dry Matter					Total, 15% Moist.			Total, % of Mean		
	2006					2006	2005	2004	2003	Total	
	5-25	6-29	7-28	8-29	10-24						
RELEASED CULTIVARS											
WL 327	2.77	2.73	1.82	1.44	0.96	9.72	11.90	10.11	9.35	41.08	48.33
HybriForce-400	2.89	2.71	1.99	1.51	0.88	9.97	12.01	9.43	9.11	40.52	47.67
4A421	2.61	2.60	1.97	1.50	0.95	9.63	12.35	9.49	9.00	40.46	47.60
Abundance	2.78	2.63	1.86	1.42	0.87	9.55	11.97	9.58	9.00	40.11	47.19
Hytest 410	2.46	2.51	1.95	1.50	0.99	9.41	12.05	9.60	9.04	40.10	47.17
Hytest 520	2.57	2.72	2.09	1.60	0.94	9.93	12.06	9.39	8.68	40.06	47.12
WL 342	2.38	2.40	1.87	1.44	0.90	8.98	12.11	9.51	9.14	39.74	46.75
Pioneer 54V54	2.72	2.66	1.89	1.47	1.02	9.76	11.72	9.36	8.61	39.45	46.41
Dagger+EV	2.55	2.54	1.79	1.42	1.03	9.32	11.80	9.52	8.70	39.34	46.28
Journey 204 Hybrid Alfalfa	2.50	2.49	1.80	1.44	0.91	9.13	11.67	9.67	8.75	39.22	46.14
Reward II	1.87	2.55	1.85	1.42	0.89	8.57	11.75	9.72	9.17	39.21	46.13
GH 750	2.56	1.98	1.83	1.11	0.87	8.35	11.99	9.79	9.03	39.16	46.07
Magna 601	2.35	2.62	1.93	1.52	1.00	9.42	12.04	9.27	8.28	39.01	45.89
5-Star	2.39	2.45	1.89	1.38	0.94	9.04	11.67	9.42	8.73	38.86	45.71
Key	2.62	2.44	1.76	1.37	0.87	9.05	11.13	9.68	8.82	38.68	45.51
WL 319 HQ	2.32	2.39	1.79	1.38	0.83	8.71	11.43	9.35	9.11	38.60	45.41
Masterpiece	2.49	1.98	1.89	1.46	0.95	8.77	11.70	9.24	8.69	38.39	45.17
Feast+EV	2.44	2.42	1.81	0.97	0.86	8.50	11.26	9.24	8.56	37.57	44.19
Kanza	2.34	2.43	1.78	1.38	0.84	8.77	11.17	8.33	6.95	35.22	41.43
Perry	2.22	2.12	1.47	1.17	0.72	7.69	--	--	--	7.69	9.05
EXPERIMENTAL STRAINS											
DS201HYB	3.07	2.84	2.16	1.72	1.06	10.84	12.62	9.98	8.55	42.00	49.41
DS218HYB	2.73	2.82	2.15	1.64	1.11	10.45	12.78	10.04	8.57	41.84	49.22
DS106HYB	2.84	2.79	2.04	1.57	0.96	10.19	12.49	9.46	9.05	41.18	48.45
DS108HYB	2.21	2.86	2.14	1.70	0.92	9.83	12.11	9.96	8.79	40.69	47.87
DS9809HYB	2.10	2.87	2.11	1.53	0.95	9.55	12.13	9.81	8.88	40.37	47.50
DS107HYB	2.84	2.62	1.91	1.10	0.92	9.40	11.86	9.46	9.13	39.85	46.88
Pioneer 55V05	2.67	2.65	1.95	1.52	0.93	9.72	12.15	9.45	8.36	39.68	46.68
CW 04030	2.78	2.73	2.12	1.56	1.03	10.21	11.90	9.08	8.49	39.67	46.67
CW 04022	2.54	2.01	2.01	1.54	0.96	9.05	12.27	9.45	8.80	39.57	46.55
ZC9953A	2.41	2.63	1.91	1.43	0.92	9.29	11.52	9.73	8.95	39.49	46.46
GPVL0144	3.25	2.59	1.91	1.44	0.88	10.06	11.52	9.52	8.30	39.40	46.35
CW 04027	2.46	2.57	2.02	1.45	0.99	9.49	12.04	9.20	8.46	39.19	46.10
CW 14026	2.58	2.63	1.98	1.48	0.70	9.37	12.03	9.32	8.36	39.08	45.97
CW 05009	2.35	2.57	2.02	1.54	1.02	9.50	12.20	9.20	8.12	39.02	45.90
CW 65086	2.60	2.61	1.86	1.51	0.98	9.55	11.85	9.12	8.43	38.94	45.82
FG 40M159A	2.23	2.39	1.84	1.36	0.89	8.72	11.49	8.93	8.53	37.67	44.31
Exp 80I	2.59	2.32	1.75	1.29	0.75	8.69	11.03	8.93	8.45	37.09	43.64
CW 65085	2.30	2.33	1.77	1.36	0.87	8.63	11.52	8.69	8.12	36.97	43.50
CW 94023	2.02	2.38	1.80	1.36	0.94	8.49	11.34	8.85	8.23	36.91	43.43
CW 94025	1.99	2.23	1.71	1.28	0.86	8.07	10.99	9.06	8.65	36.77	43.26
SUMMARY STATISTICS											
Average	2.51	2.52	1.90	1.43	0.92	9.28	11.84	9.37	8.66	39.15	46.06
LSD (0.05)	0.78	0.61	0.26	0.33	0.16	1.09	0.51	0.62	0.30	1.39	1.64
LSD(0.20)	0.51	0.39	0.17	0.21	0.10	0.71	0.57	0.40	0.23	0.91	1.07
CV (%)	22.22	17.20	9.93	16.45	12.49	8.36	3.06	4.70	2.92	2.54	2.54
MCV(%)	31.12	24.09	13.91	23.04	17.50	11.71	4.28	6.59	3.46	3.56	4

Table 7. 2006 Performance Test entries, with disease and insect resistance ratings for released varieties.*

Brand Name	A A S N												Brand Name	A A S N																			
	W	B	V	F	A	R	A	P	A	S	H	K	K	P	L	G	W	B	V	F	A	R	A	P	A	S	H	K	K	P	L	G	
	S	W	W	W	N	R	A	A	A	N	1	2	N	N	L	E	S	W	W	W	N	R	A	A	A	N	1	2	N	N	L	E	
ABI																																	
AA108E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	631	2	H	R	H	R	H	R	H	M	R	M	-	-	-	-		
AA112E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6400HT	2	H	H	H	H	-	H	-	-	H	-	-	-	-	-	Y	
Allied																6415	1	H	H	H	H	R	R	-	H	-	-	-	H	-	-		
CW 15030	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6420	2	H	R	H	R	H	R	R	-	R	R	-	H	-	-	-	
Evermore	2	H	H	H	H	H	H	R	-	R	H	-	M	-	-	6530	-	H	H	H	H	-	H	-	R	H	M	-	-	-	-	-	
FSG351	2	H	R	H	R	H	R	R	R	-	H	-	-	-	-	Dagger+EV	-	H	H	H	H	M	H	M	R	H	-	L	-	-	-	-	
FSG406	1	H	H	H	H	-	R	-	R	H	-	R	-	H	-	Feast+EV	2	H	H	H	R	H	-	M	-	H	-	-	-	-	-	-	
FSG408DP	2	H	R	H	H	H	-	R	R	R	-	H	-	-	-	ZC9953A	-																
FSG505	2	H	H	H	H	H	R	R	-	R	H	-	R	-	-	Golden Harvest																	
BioPlant																GH 750	-	H	H	H	H	R	R	-	M	H	-	-	-	-	-	-	
Phirst	2	H	R	H	H	H	R	R	-	R	R	-	H	-	-	Great Plains																	
Cal/West																CL2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CW 04022	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	GPVL0144	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CW 04027	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Key	-	H	H	H	H	H	M	M	M	M	-	-	-	-	-	-	
CW 04030	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Regal	-	H	R	H	R	H	-	H	M	-	M	-	-	-	-	-	
CW 05009	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Hytest																	
CW 14026	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Hytest 410	-	H	H	H	H	H	M	-	R	H	-	-	-	-	-	-	
CW 65085	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Hytest 520	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CW 65086	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Johnston																	
CW 94023	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Good as Gold	-	H	R	H	R	H	-	R	-	M	M	-	H	-	-	-	
CW 94025	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	II																	
Channel																KS AES & USDA																	
Notice II	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Kanza	-	R	-	-	-	R	R	-	-	-	-	-	-	-	-	-	
Cimarron USA																Monsanto																	
Cimarron	-	R	R	H	H	H	H	H	R	R	R	-	S	-	-	DKA42-15	1	H	H	H	H	H	R	H	-	R	H	-	-	H	-	-	
VL400	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	DKA50-18	2	H	H	H	H	R	R	-	R	H	-	-	H	-	-	-	
CroPlan Genetics																msSUNSTRA/Dairyland																	
5-Star	3	R	R	H	R	R	R	R	R	R	R	R	-	-	-	404	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Dairyland																405	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Arapaho	2	H	R	H	R	H	-	M	-	R	R	-	H	-	-	406	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
DS106HYB	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	407	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
DS107HYB	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mycogen																	
DS108HYB	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4A421	-	H	H	H	H	H	H	-	H	-	M	-	-	-	-	-	
DS201HYB	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	NC+																	
DS218HYB	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jade III	2	H	R	H	H	R	R	R	R	-	H	-	-	-	-	-	
DS361HY	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	NE AES & USDA																	
DS362HY	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Perry	-	R	-	L	-	M	R	-	-	-	M	-	-	-	-	-	
DS415	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	NK																	
DS416	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Expedition	3	R	H	H	H	R	-	R	H	-	R	-	-	-	-	-	
DS9809HYB	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Genoa	1	H	H	H	H	-	R	-	R	H	-	-	-	-	-	-	
HybriForce-400	2	H	R	H	R	H	H	R	M	R	M	-	H	-	-	PGI																	
HybriForce-420/wet	2	H	R	H	R	H	R	R	-	H	R	-	H	-	-	Integrity	-	H	H	H	H	-	-	H	R	-	-	Y	-	-	-	-	
Magna 601	3	R	M	H	R	H	H	R	-	R	M	-	R	R	-	Reward II	2	H	R	H	R	R	R	R	R	-	H	-	-	-	-	-	
Forage Genetics																																	
FG 40M159A	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-																		

*WS = Winter survival, 1 = superior
 BW = Bacterial wilt
 VW = Verticillium wilt
 FW = Fusarium wilt
 AN = Anthracnose race 1
 PRR = Phytophthora root rot
 SAA = Spotted alfalfa aphid
 PA = Pea aphid
 Disease and insect resistance ratings are from the National Alfalfa Alliance, NAAIC descriptions, or from developers of the varieties.

BAA = Blue alfalfa aphid
 SN = Stem nematode
 APH1 = Aphanomyces root rot race 1
 APH2 = Aphanomyces root rot race 2
 SRKN = Southern root knot nematode
 NRKN = Northern root knot nematode
 PL = Potato leafhopper
 MLE = Multi-foliate expression

GT = Continuous grazing tolerance, Y/N

Pest resistance ratings:		
Code	Resistance class	% Resistant plants
S	Susceptible	0-5%
L	Low Resistance	6-14%
M	Moderate Resistance	15-30%
R	Resistance	31-50%
H	High Resistance	>50%
-	Not adequately tested	

Table 7. 2006 Performance Test entries, with disease and insect resistance ratings for released varieties.*

Brand Name	A A S N												A A S N																			
	W	B	V	F	A	R	A	P	A	S	H	K	K	P	L	G	W	B	V	F	A	R	A	P	A	S	H	K	K	P	L	G
	S	W	W	W	N	R	A	A	A	N	1	2	N	N	L	E	S	W	W	W	N	R	A	A	A	N	1	2	N	N	L	E
Pioneer																																
54Q25	-	H	H	H	H	H	R	R	-	H	R	-	-	H	-	-																
54V46	-	R	H	H	H	H	R	R	L	M	H	R	-	H	-	-																
54V54	-	H	H	H	H	H	R	-	-	L	M	-	-	-	-	-																
55V05	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-																
Power																																
Power 4.2	-	H	R	H	H	H	R	R	-	H	H	-	-	R	-	H	-															
Sharp																																
Abundance	2	H	R	H	R	H	R	R	M	R	R	-	-	H	-	-																
Journey 204	-	H	R	H	H	H	R	R	-	R	R	-	-	H	-	-																
Hybrid Alfalfa																																
Simplot																																
Masterpiece	3	H	R	H	H	H	R	-	R	H	R	-	-	R	-	M	-															
Star																																
Exp 80I	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-																
Taylor																																
Maximizer	1	H	H	H	H	H	R	-	-	R	H	-	-	-	-	H	-															
W-L Research																																
WL 319 HQ	1	H	H	H	H	H	R	H	-	M	H	-	-	-	-	H	-															
WL 327	-	H	R	H	H	H	R	R	R	H	R	H	-	-	-	-	-															
WL 335 HQ	1	H	H	H	H	H	R	H	-	M	H	-	-	-	-	H	-															
WL 342	1	H	H	H	H	H	H	H	-	R	H	-	-	-	-	H	-															
WL 357 HQ	2	H	H	H	H	H	-	H	-	H	-	-	-	-	-	-	-															

*WS = Winter survival, 1 = superior
BW = Bacterial wilt

VW = Verticillium wilt

FW = Fusarium wilt

AN = Anthracnose race 1

PRR = Phytophthora root rot

SAA = Spotted alfalfa aphid

PA = Pea aphid

Disease and insect resistance ratings are from the National Alfalfa Alliance, NAAIC descriptions, or from developers of the varieties.

BAA = Blue alfalfa aphid

SN = Stem nematode

APH1 = Aphanomyces root rot race 1

APH2 = Aphanomyces root rot race 2

SRKN = Southern root knot nematode

NRKN = Northern root knot nematode

PL = Potato leafhopper

MLE = Multi-foliate expression

GT = Continuous grazing tolerance, Y/N

Pest resistance ratings:

Code	Resistance class	% Resistant plants
S	Susceptible	0-5%
L	Low Resistance	6-14%
M	Moderate Resistance	15-30%
R	Resistance	31-50%
H	High Resistance	>50%
-	Not adequately tested	

For those interested in accessing crop performance testing information electronically, visit our World Wide Web site. Most of the information contained in this publication is available for viewing or downloading.

The URL is <http://kscroptests.agron.ksu.edu>

Excerpts from the
University Research Policy Agreement with Cooperating Seed Companies

Permission is hereby given to Kansas State University to test varieties and/or hybrids designated on the attached entry forms in the manner indicated in the test announcements. I certify that seed submitted for testing is a true sample of the seed being offered for sale.

I understand that all results from Kansas Crop Performance Tests belong to the University and the public and shall be controlled by the University so as to produce the greatest benefit to the public. Performance data may be used in the following ways: 1) Tables may be reproduced in their entirety provided the source is referenced and data are not manipulated or reinterpreted; 2) Advertising statements by an individual company about the performance of its entries may be made as long as they are accurate statements about the data as published, with no reference to other companies' names or cultivars. In both cases, the following must be included with the reprint or ad citing the appropriate publication number and title: "See the official Kansas State University Agricultural Experiment Station and Cooperative Extension Service Report of Progress 971 '2006 Kansas Performance Tests with Alfalfa Varieties,' or the Kansas Crop Performance Test Web site, <http://kscroptests.agron.ksu.edu>, for details. Endorsement or recommendation by Kansas State University is not implied."

These materials may be freely reproduced for educational purposes. All other rights reserved. In each case, give credit to the author(s), name of work, Kansas State University, and the date the work was published.

Contributors

Main Station, Manhattan

Jane Lingenfelser, Assistant Agronomist (Senior Author)
Kraig Roozeboom, Assistant Professor

Research Centers

Pat Evans, Colby
Joseph Moyer, Mound Valley
Monty Spangler, Garden City

Experiment Fields

Barney Gordon, Belleville
William Heer, Hutchinson
Larry Maddux, Topeka

*NOTE: Trade names are used to identify products.
No endorsement is intended, nor is any criticism implied of similar products not named.*

**This Report of Progress was edited, designed, and printed
by the Department of Communications at Kansas State University**