


# 2002

KANSAS PERFORMANCE TESTS WITH  
**ALFALFA  
VARIETIES**

REPORT OF PROGRESS 904

Kansas State University  
Agricultural Experiment Station  
and Cooperative Extension Service



## TABLE OF CONTENTS

### **2002 Performance Tests**

Objectives and Procedures.....	1
Variety Characterization .....	1
Northeast, Powhattan, Brown County                          Table 1 .....	2
Southeast, Mound Valley, Labette County                  Table 2 .....	3
North Central, Belleville, Republic County                Table 3 .....	4
South Central, Hutchinson, Reno County                 Table 4 .....	5
Northwest Irr., Colby, Thomas County                    Table 5 .....	6
Southwest Irr., Garden City, Finney County            Table 6 .....	7
APPENDIX: Fall dormancy and pest resistance ratings of released varieties .....	9
Electronic Access and University Research Policy.....	back cover

### **Entrants in 2002 Kansas Alfalfa Performance Tests.**

Allied Seed Cooperative Angola IN 800-813-5025	Garst/AgriPro Seed Slater IA 877-247-4776 garstseed.com	Monsanto Seed DeKalb IL 815-758-9323 farmsource.com	Sharp Bros Seed Company Healy KS 620-398-2231 sharpseed.com
America's Alfalfa Princeton IL 815-875-6426 americasalfalfa.com	Great Plains Research Apex NC 919-362-1583 greatplainsresearch.com	Mycogen Seed Wamego KS 785-456-2724 mycogen.com	Star Seed Co. Beloit KS 800-782-7611 starseed1.com
CroPlan Genetics Arden Hills MN 800-851-8110 croplangenetics.com	JC Robinson Seed Co Waterloo NE 402-289-0265 goldenharvestseeds.com	NC+ Hybrids Lincoln NE 402-467-2517 nc-plus.com	Syngenta Seeds, Inc. Ames IA 800-258-0498 syngenta.com
Dairyland Research Clinton WI 608-676-2237 dairylandseed.com	MBS Inc Story City IA 800-247-3967	Pioneer Hi-Bred Intl Inc Johnston IA 515-334-6645 pioneer.com	United Suppliers Inc Eldora IA 877-714-4503 uniteds.com
Donley Seed Co. New Albany IN 812-941-9822	Midwest Seed Genetics Americus KS 620-443-5353	Producers Hybrids Battle Creek NE 402-675-2975 producershybrids.com	W-L Research Inc Evansville WI 608-882-4100 wlresearch.com

# 2002 PERFORMANCE TESTS

## Objectives and Procedures

The Kansas Agricultural Experiment Station established an official alfalfa performance-testing program in 1980 to provide Kansas growers with unbiased performance comparisons of alfalfa varieties marketed in the state. Each year, private companies are asked to enter varieties voluntarily at the locations slated for establishment that year. Announcements and entry forms are mailed to private companies in June for entry in fall-seeded tests. Companies enter varieties of their choice and pay entry fees to cover part of the costs of conducting the tests. Most tests are planted in mid-August or September; however, the Southeast Kansas test usually is planted in the spring. Individual tests are conducted for a minimum of 3 years. New tests typically are established during the final production year of the previous test.

Descriptive information is presented with the results for each test. This information, including soil type, establishment methods, fertilization, pest control, irrigation, harvest dates, and growing conditions unique to that location, can help explain test and/or variety performance.

Forage yields were estimated by harvesting four replications of each variety with a plot harvester. The amount of forage produced from a specific area ( $35\text{-}80 \text{ ft}^2$ ) was weighed, and a subsample was taken to determine moisture content. This information was used to convert the plot weights to tons of dry matter per acre for each cutting, the season total, and the total for each previous season as presented in Tables 1-6. The forage yield over the lifetime of a particular test is presented as the total tons of dry matter produced per acre, as the total tons of 15% moisture hay, and as a percentage of the test average.

Each table is separated into three sections. The first lists released cultivars that are generally available on the seed market or soon will be. The second section includes experimental cultivars that were entered in the test before being released for sale. These experimental lines often represent an earlier generation of seed than that used for the released cultivars. The third section includes summary statistics unique to that test.

At the bottom of each column, the Least Significant Difference (LSD) is listed at the 0.05 and 0.20 levels. These values indicate how large a difference is needed to be confident that one variety is superior to another. Differences between varieties that are equal to or greater than the 0.05 LSD have only a 1 in 20 chance of being due to chance or error. Differences equal to or greater than the 0.20 LSD have a 1 in 5 chance of being caused by chance or error.

The Coefficient of Variability (CV) provides an estimate of the consistency of the results of a particular test. In these tests, CV's below 10% generally indicate reliable, uniform data, whereas CV's of 10-15% are not uncommon and generally indicate that the data are acceptable for rough comparisons. Tests with CV's over 15% may still be useful, but variety comparisons lack precision.

The Mean Coefficient of Variability (MCV) is similar to the CV in that it serves as an indicator of test precision. The MCV is calculated by dividing the 0.05 LSD by the test mean (average) and multiplying by 100. The MCV reveals the percent difference required to detect differences between varieties with 95% confidence. Many alfalfa breeders and testers agree that tests with MCV values greater than 10% are of little benefit.

## Variety Characterization

For variety selection, producers should consider the performance of a variety in each of the current tests where it appears, its performance over time and locations relative to familiar or check varieties, and the disease and insect resistance characteristics that are potentially important in their situation.

Tables 1-6 contain updated yield data from individual tests currently in progress. First-season yields for a spring-planted test are often more variable than yields in subsequent years. Season totals are important, but yield distribution during the season may vary among varieties. Examine yields from individual cuttings to determine if differences in yield distribution exist. Yield totals over many years provide the best measure of variety performance over time.

The appendix provides additional descriptive information such as fall dormancy, disease resistance, and insect resistance ratings. These ratings were obtained primarily from the annual 'Fall Dormancy & Pest Resistance Ratings for Alfalfa Varieties' pamphlet published by the Alfalfa Council. That report summarizes information submitted by developers of alfalfa varieties as part of the variety registration process. The Association of Official Seed Certifying Agencies (AOSCA) National Alfalfa Variety Review Board (NAVRB) reviewed the ratings before they were published. Companies submitting varieties for the tests provided ratings for some unregistered varieties.

Fall dormancy values are based on the fall canopy height measured in Minnesota. Dormancy values generally are related to the speed of regrowth. The rapid regrowth types have higher values, and the slower regrowth types have lower values.

**Table 1. Northeast Kansas, Powhattan Alfalfa Performance Test, Seeded August 1998.**

NAME	Forage Yield							
	tons/acre			Dry Matter		Total, 00-02	Total, % of Mean	
	2002	2001	2000	Total	Total			
5-30	6-26	Total	Total	Total	Total	15% Moist.		
<b>RELEASED CULTIVARS</b>								
Pioneer 53V08	3.64	2.01	5.65	6.73	3.84	16.22	19.08	104
DK 142	3.52	2.25	5.77	6.49	3.85	16.11	18.95	103
WL 325 HQ	3.47	1.90	5.37	6.67	3.94	15.98	18.80	102
Amerigraze 401+Z	3.63	1.98	5.60	6.66	3.69	15.95	18.76	102
Dagger+EV	3.49	1.83	5.32	6.55	4.05	15.92	18.73	102
WL 232 HQ	3.51	1.74	5.25	6.68	3.98	15.91	18.72	102
DK 141	3.25	1.67	4.91	6.85	4.04	15.80	18.59	101
Geneva	3.52	1.78	5.30	6.44	4.03	15.77	18.55	101
Pioneer 54H55	3.52	1.80	5.32	6.42	4.01	15.75	18.53	101
Magnum V	3.26	1.91	5.17	6.79	3.74	15.70	18.47	100
ABT350	3.52	1.73	5.24	6.74	3.67	15.65	18.41	100
Yielder	3.17	2.10	5.27	6.58	3.78	15.63	18.39	100
WL 324	3.55	1.80	5.34	6.39	3.86	15.59	18.34	100
Gold Plus	3.52	1.67	5.19	6.37	3.96	15.52	18.26	99
Emperor	3.47	1.84	5.31	6.37	3.78	15.46	18.19	99
Ace	3.43	1.82	5.25	6.30	3.91	15.46	18.19	99
Cimarron 3i	3.47	1.67	5.14	6.53	3.76	15.43	18.15	99
Spur	3.55	1.60	5.15	6.31	3.97	15.43	18.15	99
Depend+EV	3.37	1.71	5.08	6.44	3.87	15.39	18.11	98
ProGro	3.14	1.87	5.00	6.45	3.86	15.31	18.01	98
Affinity+Z	3.53	1.87	5.39	6.29	3.62	15.30	18.00	98
TMF 4464	3.06	1.69	4.75	6.77	3.75	15.27	17.96	98
Perry	3.43	2.01	5.44	6.28	3.55	15.27	17.96	98
Kanza	3.19	1.74	4.94	6.08	3.72	14.74	17.34	94
<b>EXPERIMENTAL STRAINS</b>								
C304	3.64	1.94	5.57	6.59	4.10	16.26	19.13	104
ZC9751A	3.37	2.10	5.48	6.64	4.09	16.21	19.07	104
C230	3.49	2.04	5.53	6.75	3.88	16.16	19.01	103
4G70	3.64	1.58	5.22	6.49	4.11	15.82	18.61	101
ZH9731H	3.62	1.82	5.44	6.54	3.67	15.65	18.41	100
ZC9741A	3.75	1.71	5.46	6.33	3.79	15.58	18.33	99
ZC9740A	3.51	1.78	5.29	6.30	3.76	15.35	18.06	98
<b>SUMMARY STATISTICS</b>								
Average	3.46	1.84	5.29	6.51	3.86	15.66	18.42	100
LSD(0.05)	0.34	0.34	0.48	0.39	0.23	1.15	1.35	7
LSD(0.20)	0.22	0.22	0.31	0.39	0.12	0.75	0.88	5
CV(%)	7.03	13.24	6.44	4.32	5.14	3.12	3.12	3
MCV(%)	9.83	18.48	9.07	4.32	5.96	7.34	7.34	7

<b>LOCATION:</b> Northeast Kansas Site: Cornbelt Experiment Field County: Brown Town: Powhattan Soil: Grundy silty clay loam	<b>2002 FERTILIZATION:</b> None in 2002. <b>2002 PEST CONTROL:</b> None needed.	<b>2002 CONDITIONS:</b> Below-normal rainfall from July to the end of the season limited regrowth, preventing more than 2 cuttings.
<b>ESTABLISHMENT:</b> 9/2/98; RCBD, 4 reps Plots 5'x20'; 4'x20' harvested 15 lb. seed/acre		

**Table 2. Southeast Kansas, Mound Valley Alfalfa Performance Test, Seeded April 2001.**

NAME	Forage Yield									
	tons/acre									
	Dry Matter					2001 Total	01-02 Total	Total, 15% Moist.	01-02 Total, % of Mean	
	5-7	6-18	7-17	8-22	Total					
<b>RELEASED CULTIVARS</b>										
HybriForce-400	1.97	1.84	0.93	0.68	5.42	1.34	6.76	7.95	111	
6420	1.74	1.91	1.01	0.71	5.36	1.21	6.57	7.73	108	
Dagger+EV	1.69	1.88	0.84	0.67	5.08	1.29	6.37	7.49	104	
Pawnee	1.60	1.79	0.85	0.74	4.98	1.13	6.11	7.19	100	
Perry	1.71	1.70	0.93	0.69	5.03	1.07	6.09	7.16	100	
5-Star	1.57	1.74	0.96	0.62	4.89	1.17	6.06	7.13	99	
Pioneer 54V54	1.72	1.79	0.92	0.62	5.04	1.01	6.05	7.12	99	
400SCL	1.66	1.79	0.91	0.57	4.94	1.04	5.98	7.04	98	
350	1.63	1.68	0.90	0.61	4.82	1.12	5.94	6.99	97	
Kanza	1.52	1.72	0.88	0.59	4.70	1.25	5.94	6.99	97	
WL 342	1.63	1.79	0.72	0.64	4.77	1.08	5.85	6.88	96	
Rebound 4.2	1.58	1.72	0.95	0.55	4.80	1.02	5.82	6.85	95	
WL 327	1.39	1.69	0.96	0.66	4.70	1.12	5.82	6.85	95	
<b>SUMMARY STATISTICS</b>										
Average	1.65	1.77	0.90	0.64	4.96	1.14	6.10	7.18	100	
LSD(0.05)	0.22	NS	NS	NS	NS	0.15	0.53	0.62	9	
LSD(0.20)	0.14	NS	NS	NS	0.32	0.09	0.34	0.40	6	
CV(%)	9.28	8.08	14.13	20.28	6.94	8.96	6.08	6.08	6	
MCV(%)	13.31	NS	NS	NS	NS	12.85	8.71	8.71	9	

<b>LOCATION:</b> Southeast Kansas Site: Southeast Ag. Research Center <b>County:</b> Labette <b>Town:</b> Mound Valley <b>Soil:</b> Parsons silty clay loam	<b>2002 FERTILIZATION:</b> 20-50-200 lb/a of N-P2O5-K2O on February 22.  <b>2002 PEST CONTROL:</b> Sprayed with Lorsban to control weevil on April 17. Poast applied on July 5 to control grass infestation.	<b>2002 CONDITIONS:</b> Dry conditions in July and August lowered yields from later cuttings and prevented a fifth cutting. Blister beetles invaded the plots in late July.
---	--	--

**Table 3. North Central Kansas, Belleville Alfalfa Performance Test, Seeded Sept. 2001.**

NAME	Forage Yield				2002 Total, 15% Moist.	2002 Total, % of Mean		
	tons/acre							
	Dry Matter		2002 Total					
	5-19	6-25	8-10					
<b>Released Cultivars</b>								
Pawnee	2.82	1.56	0.10	4.48	5.27	117		
645-II	2.63	1.45	0.08	4.16	4.89	108		
Abundance	2.63	1.45	0.08	4.16	4.89	108		
HybriForce-400	2.59	1.43	0.07	4.09	4.81	107		
Feast+EV	2.57	1.42	0.08	4.07	4.79	106		
Reliance	2.49	1.37	0.08	3.94	4.64	103		
Abilene+Z	2.50	1.38	0.06	3.94	4.64	103		
Enhancer	2.47	1.37	0.10	3.94	4.64	103		
6410	2.48	1.37	0.07	3.92	4.61	102		
Ameristand 403T	2.46	1.36	0.09	3.91	4.60	102		
US A4230	2.47	1.37	0.07	3.91	4.60	102		
GH 750	2.46	1.35	0.07	3.88	4.56	101		
DKA42-15	2.45	1.36	0.07	3.88	4.56	101		
Geneva	2.44	1.35	0.07	3.86	4.54	101		
Lightning II	2.44	1.34	0.08	3.86	4.54	101		
Dagger+EV	2.39	1.32	0.10	3.81	4.48	99		
Journey 204 Hybrid Alfalfa	2.41	1.33	0.07	3.81	4.48	99		
Pioneer 54Q53	2.39	1.32	0.08	3.79	4.46	99		
FK421	2.38	1.32	0.07	3.77	4.44	98		
Macon	2.37	1.31	0.06	3.74	4.40	97		
Kanza	2.01	1.11	0.05	3.17	3.73	83		
Perry	1.98	1.10	0.05	3.13	3.68	82		
<b>Experimental Strains</b>								
ZC9840A	2.62	1.45	0.09	4.16	4.89	108		
ZC9851A	2.57	1.42	0.08	4.07	4.79	106		
A 30-6	2.51	1.39	0.08	3.98	4.68	104		
ZG9840	2.50	1.38	0.07	3.95	4.65	103		
ZC9941A	2.45	1.35	0.08	3.88	4.56	101		
ZC9953A	2.44	1.35	0.06	3.85	4.53	100		
ZG9941	2.41	1.33	0.07	3.81	4.48	99		
ZC9940A	2.39	1.32	0.07	3.78	4.45	98		
4M72	2.37	1.31	0.06	3.74	4.40	97		
ZC9942A	2.37	1.30	0.06	3.73	4.39	97		
ZC9841A	2.34	1.29	0.07	3.70	4.35	96		
ZC9950A	2.31	1.28	0.06	3.65	4.29	95		
5M87	2.19	1.21	0.07	3.47	4.08	90		
ZC9842A	2.12	1.17	0.06	3.35	3.94	87		
<b>Summary Statistics</b>								
Average	2.43	1.34	0.07	3.84	4.52	100		
LSD(0.05)	0.26	0.14	0.03	0.41	0.48	11		
LSD(0.20)	0.17	0.09	0.01	0.27	0.32	7		
CV(%)	7.56	7.55	25.05	7.62	7.62	8		
MCV(%)	10.70	10.45	42.86	10.68	10.68	11		

<b>LOCATION:</b> North Central Kansas Site: North Central Kansas Exp. Field County: Republic Town: Belleville Soil: Crete silt loam	<b>2002 FERTILIZATION:</b> 36-92-0 applied April 15, 2002. <b>2002 PEST CONTROL:</b> None needed.	<b>2002 CONDITIONS:</b> Rainfall was well below normal for most of the summer, preventing a fourth cutting. Recorded precipitation was the lowest June, July, and August total since 1934.
<b>ESTABLISHMENT:</b> 8/22/01; RCBD, 4 reps Plots 5'x15'; 3'x15' harvested 18 lb. seed/acre		

**Table 4. South Central Kansas, Hutchinson Alfalfa Performance Test, Seeded Sept. 1999.**

NAME	Plant Height inches	Forage Yield										00-02 Total, 15% Moist. % of Mean	
		tons/acre					Dry Matter			2001 Total	2000 Total	00-02 Total	
		2002	5-29	6-25	7-17	8-28	Total	2001 Total	2000 Total				
NAME	8-28												
<b>RELEASED CULTIVARS</b>													
WL 327	13	2.17	1.69	0.98	1.68	6.52	4.43	5.89	16.84	19.81	109		
Magnum V	15	2.09	1.88	1.16	1.10	6.23	4.28	5.66	16.17	19.02	105		
Kanza	18	1.95	1.88	1.33	1.30	6.46	4.43	5.16	16.05	18.88	104		
Aspire	18	2.13	1.78	1.25	1.17	6.33	4.05	5.65	16.03	18.86	104		
Pioneer 54Q53	15	1.95	2.02	1.18	1.07	6.22	4.22	5.41	15.85	18.65	103		
Forecast 1001	16	2.01	1.98	1.10	1.23	6.32	4.21	5.28	15.81	18.60	103		
ABT350	13	2.02	1.99	0.92	1.34	6.27	4.60	4.90	15.77	18.55	103		
DK 142	14	2.14	2.09	0.98	1.02	6.23	4.38	5.15	15.76	18.54	102		
6420	16	1.99	1.84	1.10	1.08	6.01	4.57	5.12	15.70	18.47	102		
TMF 4464	14	2.13	2.09	1.04	0.98	6.24	4.44	4.94	15.62	18.38	102		
WL 232 HQ	13	1.95	2.04	1.05	1.18	6.22	4.45	4.94	15.61	18.36	101		
Abilene+Z	15	1.89	1.90	1.25	1.26	6.30	4.36	4.87	15.53	18.27	101		
Dagger+EV	15	2.07	1.89	0.98	0.96	5.90	4.35	5.01	15.26	17.95	99		
Cimarron SR	15	1.87	1.91	0.98	0.96	5.72	4.46	4.88	15.06	17.72	98		
DK 140	15	1.80	1.96	0.96	1.08	5.80	4.30	4.88	14.98	17.62	97		
Award	13	1.83	1.88	0.93	1.15	5.79	4.12	4.86	14.77	17.38	96		
Perry	13	1.99	1.55	0.94	0.95	5.43	4.48	4.77	14.68	17.27	95		
Cimarron 3i	13	1.85	1.81	0.93	0.97	5.56	4.15	4.96	14.67	17.26	95		
Macon	14	1.64	1.78	1.00	1.02	5.44	3.99	4.37	13.80	16.24	90		
<b>EXPERIMENTAL STRAINS</b>													
ZC9650	15	2.00	1.83	1.10	0.87	5.80	4.29	5.25	15.34	18.05	100		
ZC9850A	13	1.95	1.88	0.93	1.06	5.82	4.45	4.92	15.19	17.87	99		
<b>SUMMARY STATISTICS</b>													
Average	14	1.96	1.88	1.04	1.11	5.99	4.33	5.06	15.38	18.09	100		
LSD(0.05)	2	NS	NS	0.17	0.31	0.75	0.26	0.46	1.56	1.84	10		
LSD(0.20)	1	NS	0.19	0.10	0.21	0.48	0.17	0.23	1.00	1.18	7		
CV(%)	9	14.10	10.92	9.84	19.79	7.31	4.30	7.64	4.00	4.00	4		
MCV(%)	16	NS	NS	16.30	28.07	12.52	6.06	9.02	10.14	10.14	10		

<b>LOCATION:</b> South Central Kansas Site: South Central Experiment Field County: Reno Town: Hutchinson Soil: Ost silt loam <b>ESTABLISHMENT:</b> 9/14/99; RCBD, 4 reps Plots 5'x20', 3'x20' harvested 18 lb. seed/acre	<b>2002 FERTILIZATION:</b> 75-40-40 prior to planting <b>2002 PEST CONTROL:</b> Sinbar applied for weed control. Furadan applied to control alfalfa weevils.	<b>2002 CONDITIONS:</b> A June hailstorm lowered second-cutting yields. Dry conditions in July and August lowered third and fourth-cutting yields.
--	--	---

**Table 5. Northwest Kansas, Colby Irrigated Alfalfa Performance Test, Seeded August 2001.**

NAME	Forage Yield					2002 Total, 15% Moist.	2002 Total, % of Mean		
	tons/acre								
	Dry Matter				Total				
	2002	6-11	7-8	8-7	9-16				
<b>Released Cultivars</b>									
Lightning II	2.95	1.19	1.34	1.40	6.89	8.11	111		
Magnum V	2.73	1.29	1.38	1.36	6.76	7.95	109		
631	2.71	1.16	1.51	1.35	6.73	7.92	108		
Enhancer	2.63	1.24	1.30	1.48	6.65	7.82	107		
645-II	2.61	1.13	1.51	1.38	6.64	7.81	107		
Target II Plus	2.71	1.19	1.35	1.35	6.61	7.78	106		
Pioneer 53V08	2.56	1.20	1.34	1.40	6.51	7.66	105		
Pioneer 54Q53	2.36	1.14	1.40	1.39	6.29	7.40	101		
Reward	2.46	1.18	1.21	1.31	6.17	7.26	99		
Kanza	2.44	1.08	1.20	1.32	6.05	7.12	97		
4200	2.45	1.05	1.27	1.17	5.95	7.00	96		
Geneva	2.34	1.10	1.13	1.38	5.94	6.99	96		
Feast+EV	2.17	1.05	1.33	1.19	5.75	6.76	92		
Perry	2.05	0.77	1.14	1.20	5.18	6.09	83		
<b>Experimental Strains</b>									
A 30-6	2.68	1.12	1.41	1.55	6.77	7.96	109		
ZC9840A	2.51	1.09	1.48	1.41	6.50	7.65	105		
ZC9842A	2.53	1.14	1.25	1.50	6.43	7.56	103		
ZC9841A	2.44	1.03	1.37	1.58	6.43	7.56	103		
6M71	2.33	1.24	1.39	1.42	6.38	7.51	103		
CW 94008	2.40	1.24	1.46	1.27	6.37	7.49	102		
ZC9950A	2.42	1.24	1.34	1.35	6.35	7.47	102		
5M84	2.38	1.15	1.38	1.41	6.32	7.44	102		
ZC9851A	2.58	1.06	1.30	1.34	6.28	7.39	101		
5M85	2.39	1.23	1.29	1.32	6.24	7.34	100		
CW 75044	2.25	1.11	1.44	1.43	6.23	7.33	100		
ZC9854A	2.57	0.95	1.23	1.46	6.22	7.32	100		
CW 64004	2.33	1.18	1.27	1.41	6.19	7.28	100		
ZC9940A	2.22	1.16	1.15	1.46	6.00	7.06	96		
CW 64026	2.41	1.13	1.22	1.24	6.00	7.06	96		
CW 74040	2.19	1.03	1.21	1.41	5.84	6.87	94		
CW 64049	2.24	0.95	1.29	1.31	5.80	6.82	93		
CW 54033	2.21	0.99	1.12	1.25	5.57	6.55	90		
CW 73029	1.98	0.87	1.15	1.36	5.37	6.32	86		
<b>Summary Statistics</b>									
Average	2.43	1.11	1.31	1.37	6.22	7.32	100		
LSD(0.05)	0.35	0.19	0.19	0.19	0.47	0.55	8		
LSD(0.20)	0.23	0.13	0.13	0.13	0.31	0.36	5		
CV(%)	10.20	12.29	10.46	9.83	5.43	5.00	5		
MCV(%)	14.32	17.29	14.66	13.77	7.62	7.62	8		

<b>LOCATION:</b> Northwest Kansas Site: Northwest Research-Extension County: Thomas Town: Colby Soil: Keith silt loam <b>ESTABLISHMENT:</b> 8/29/01; RCBD, 4 reps Plots 3'x20'; 3'x17' harvested 18 lb. seed/acre	<b>2002 FERTILIZATION:</b> 17-60-0 prior to planting <b>2002 PEST CONTROL:</b> No insect or disease problems.	<b>2002 CONDITIONS:</b> Spring and early summer were very dry. A total of 35.3" of water was added in 10 irrigations.
---	--	---

**Table 6. Southwest Kansas, Garden City Irrigated Alfalfa Performance Test, Seeded Sept. 1999.**

NAME	Forage Yield										00-02 Total, % of Mean	
	tons/acre					Dry Matter						
	2002		2001			2000	00-02	Total,	15%	Moist.		
	5-20	7-1	8-1	9-5	Total	Total	Total	Total	Total	Total		
<b>RELEASED CULTIVARS</b>												
TMF 4464	3.27	4.12	2.31	2.05	11.75	9.61	12.16	33.52	39.44	104		
Pioneer 54Q53	3.19	3.99	2.42	2.10	11.70	9.71	12.08	33.49	39.40	104		
Magnum V	3.31	3.83	2.38	2.13	11.65	9.20	12.15	33.00	38.82	102		
Aspire	3.26	3.55	2.44	2.16	11.41	9.61	11.77	32.79	38.58	102		
Dagger+EV	3.25	3.82	2.34	2.06	11.47	9.55	11.75	32.77	38.55	102		
Pioneer 53V08	3.02	4.06	2.42	2.16	11.66	9.24	11.85	32.75	38.53	102		
WL 327	3.09	3.65	2.31	2.04	11.09	9.39	12.21	32.69	38.46	102		
Forecast 1001	3.07	3.74	2.41	2.14	11.36	9.41	11.87	32.64	38.40	101		
Emperor	3.12	3.71	2.30	2.07	11.20	9.34	11.97	32.51	38.25	101		
Abilene+Z	3.22	3.76	2.31	2.14	11.43	9.34	11.68	32.45	38.18	101		
Jade II	3.11	3.87	2.28	2.07	11.33	9.10	12.01	32.44	38.16	101		
Affinity+Z	3.32	3.36	2.32	2.10	11.10	9.47	11.84	32.41	38.13	101		
GH 750	3.12	3.93	2.23	1.97	11.25	9.17	11.77	32.19	37.87	100		
Cimarron 3i	3.21	3.52	2.20	2.04	10.97	9.25	11.97	32.19	37.87	100		
ABT 400SCL	3.23	3.48	2.23	2.07	11.01	9.28	11.87	32.16	37.84	100		
ABT350	3.15	3.95	2.29	2.05	11.44	9.27	11.34	32.05	37.71	100		
DK 140	3.05	3.97	2.29	2.01	11.32	9.13	11.47	31.92	37.55	99		
FQ315	2.96	3.79	2.30	2.05	11.10	9.11	11.62	31.83	37.45	99		
6420	3.13	3.59	2.34	2.01	11.07	9.15	11.60	31.82	37.44	99		
Cimarron SR	3.09	3.69	2.17	2.08	11.03	9.03	11.73	31.79	37.40	99		
Kanza	2.95	3.84	2.41	2.09	11.29	9.21	10.83	31.33	36.86	97		
Award	3.11	3.41	2.22	2.03	10.77	8.93	11.18	30.88	36.33	96		
Perry	3.23	3.46	2.08	1.98	10.75	8.76	11.30	30.81	36.25	96		
DK 142	3.04	3.40	1.98	1.97	10.39	8.95	11.43	30.77	36.20	96		
<b>EXPERIMENTAL STRAINS</b>												
ZC9850A	3.21	4.14	2.48	2.21	12.04	9.58	12.10	33.72	39.67	105		
DS983809	3.19	4.27	2.38	2.10	11.94	9.38	12.39	33.71	39.66	105		
HybriForce-400	3.20	4.01	2.35	2.11	11.67	9.67	11.98	33.32	39.20	103		
DS983810	3.23	3.78	2.36	2.15	11.52	9.20	12.37	33.09	38.93	103		
ZC9853A	3.48	3.77	2.29	2.11	11.65	9.55	11.85	33.05	38.88	103		
DS983811	3.26	4.06	2.33	2.02	11.67	9.36	11.99	33.02	38.85	103		
DS9707 HYB	3.23	4.08	2.39	2.12	11.82	9.21	11.92	32.95	38.76	102		

**Table 6. Southwest Kansas, Garden City Irrigated Alfalfa Performance Test, Seeded Sept. 1999.**

NAME	Forage Yield										00-02 Total, % of Mean	
	tons/acre					Dry Matter						
	2002					2001	2000	00-02	Total,	15% Moist.		
	5-20	7-1	8-1	9-5	Total	Total	Total	Total	Total	Total		
DS983808	3.13	3.80	2.38	2.13	11.44	9.14	12.36	32.94	38.75	102		
ZC9650	3.19	3.93	2.39	2.18	11.69	9.08	11.92	32.69	38.46	102		
ZC9851A	3.20	3.78	2.40	2.13	11.51	9.18	11.81	32.50	38.24	101		
DS983812	3.21	3.52	2.35	2.03	11.11	9.23	11.94	32.28	37.98	100		
CW 64025	3.03	3.74	2.31	2.07	11.15	9.31	11.67	32.13	37.80	100		
ZC9854A	3.10	3.84	2.29	2.05	11.28	9.15	11.64	32.07	37.73	100		
CW 74033	3.04	3.90	2.38	2.08	11.40	9.08	11.48	31.96	37.60	99		
DS983813	3.19	3.87	2.28	2.06	11.40	9.13	11.27	31.80	37.41	99		
DS9704 HYB	2.93	3.69	2.33	2.13	11.08	8.93	11.79	31.80	37.41	99		
CW 64018	2.95	3.80	2.30	2.03	11.08	9.03	11.40	31.51	37.07	98		
CW 84025	2.97	3.58	2.24	2.05	10.84	9.31	11.24	31.39	36.93	97		
CW 84024	2.94	3.50	2.26	2.11	10.81	8.94	11.64	31.39	36.93	97		
ZC9840A	3.13	3.91	2.09	1.93	11.06	8.90	11.40	31.36	36.89	97		
ZC9841A	3.18	3.89	2.16	2.06	11.29	8.78	11.13	31.20	36.71	97		
ZC9842A	3.24	3.36	2.25	2.02	10.87	9.21	11.07	31.15	36.65	97		
CW 74043	2.99	3.47	2.17	1.87	10.50	8.85	11.41	30.76	36.19	96		

**SUMMARY STATISTICS**

Average	3.14	3.77	2.30	2.07	11.28	9.21	11.71	32.20	37.88	100
LSD(0.05)	0.22	0.51	0.15	0.11	0.73	0.45	0.43	1.70	2.00	5
LSD(0.20)	0.14	0.33	0.10	0.08	0.48	0.29	0.22	1.11	1.31	3
CV(%)	4.91	9.70	4.68	3.86	4.66	3.48	3.17	2.24	2.64	3
MCV(%)	6.86	13.54	6.53	5.39	6.52	4.86	3.71	5.28	5.28	5

<b>LOCATION:</b> Southwest Kansas Site: Southwest Res.-Ext. Center County: Finney Town: Garden City Soil: Keith silt loam  <b>ESTABLISHMENT:</b> 8/24/99; RCBD, 4 reps Plots 3'x20'; 3'x20' harvested 32 lb. seed/acre	<b>2002 FERTILIZATION:</b> 22-104-0 applied at planting  <b>2002 PEST CONTROL:</b> Pursuit Plus applied in March to control grasses. Cygon 400 applied on August 8 to control cowpea aphids.	<b>2002 CONDITIONS:</b> Minimal disease and insect stress facilitated excellent yields. Hot, dry winds may have reduced yields to some extent.
---	--	---

**Appendix 1: Varieties in 2002 Kansas Alfalfa Performance Tests  
with unverified fall dormancy and disease and insect resistance ratings.**

COMPANY Name	S N										COMPANY Name	S N												
	F	B	V	F	A	R	A	P	A	S	P	K	K	P	F	B	V	F	A	R	A	P	S	N
D	W	W	N	R	A	A	A	N	H	N	N	L	D	W	W	N	R	A	A	A	N	H	N	L
<b>AgriPro</b>																								
Depend+EV	4	H	H	H	H	H	M	R	S	M	R	-	-	-										
Yielder	3	H	H	H	R	H	-	R	-	-	-	-	-	-										
<b>Allied</b>																								
350	3	-	-	-	-	-	-	-	-	-	-	-	-	-										
400SCL	4	-	-	-	-	-	-	-	-	-	-	-	-	-										
Macon	4	H	H	H	H	H	R	R	-	M	H	-	-	-										
Reliance	3	-	-	-	-	-	-	-	-	-	-	-	-	-										
<b>America's Alfalfa</b>																								
Abilene+Z	5	H	H	H	H	H	M	H	M	R	R	-	-	-										
Affinity+Z	4	H	H	H	H	H	-	R	-	R	R	-	-	-										
Amerigraze 401+Z	4	H	H	H	H	H	-	R	-	R	R	-	-	-										
Ameristand 403T	-	-	-	-	-	-	-	-	-	-	-	-	-	-										
Emperor	4	H	H	H	H	H	M	R	-	H	-	-	-	-										
<b>AV</b>																								
4200	-	-	-	-	-	-	-	-	-	-	-	-	-	-										
<b>Cargill</b>																								
FQ315	3	H	R	H	H	H	H	R	-	R	H	-	-	-										
<b>CroPlan Genetics</b>																								
5-Star	5	-	-	-	-	-	-	-	-	-	-	-	-	-										
Rebound 4.2	4	-	-	-	-	-	-	-	-	-	-	-	-	-										
<b>Dairyland</b>																								
Forecast 1001	4	H	R	R	R	H	-	R	-	R	R	-	H	H										
HybriForce-400	4	-	-	-	-	-	-	-	-	-	-	-	-	-										
Magnum V	4	H	R	H	R	H	R	R	M	R	M	-	M	M										
<b>Donley</b>																								
FK421	-	-	-	-	-	-	-	-	-	-	-	-	-	-										
<b>Garst</b>																								
631	4	H	R	H	R	H	R	H	M	R	M	-	-	-										
6410	4	H	H	H	H	H	H	R	-	M	H	-	-	-										
6420	4	H	R	H	-	H	R	R	-	R	R	-	H	H										
645-II	3	H	H	H	H	H	-	R	-	H	-	-	-	-										
<b>Garst/AgriPro</b>																								
Dagger+EV	5	H	H	H	H	H	M	H	M	R	H	-	L	L										
Feast+EV	3	H	H	H	R	H	-	M	-	H	-	-	-	-										
<b>Golden Harvest</b>																								
GH 750	4	H	H	H	H	H	R	R	R	M	H	-	-	-										
<b>Great Plains</b>																								
Cimarron 3i	4	H	R	H	H	R	R	R	-	R	M	R	-	-										
Cimarron SR	4	H	H	H	H	H	R	H	-	R	M	R	-	-										
<b>KS AES &amp; USDA</b>																								
Kanza	3	R	-	-	-	R	R	-	-	-	-	-	-	-										
<b>MBS</b>																								
Gold Plus	4	H	R	H	H	H	H	H	-	H	R	-	-	-										
ProGro	4	H	R	H	R	H	R	R	M	-	M	-	-	-										
Reward	4	H	R	H	R	H	R	H	M	M	M	-	-	-										

**Variety characterization codes:**

FD = Fall dormancy rating

BW = Bacterial wilt

VW = Verticillium wilt

FW = Fusarium wilt

AN = Anthracnose race 1

PRR = Phytophthora root rot

SAA = Spotted alfalfa aphid

PA = Pea aphid

BAA = Blue alfalfa aphid

SN = Stem nematode

APH = Aphanomyces root rot race 1

SRKN = Southern root knot nematode

NRKN = Northern root knot nematode

PL = Potato leafhopper

**Fall dormancy ratings:**

Check variety

Rating

Code

**Pest resistance ratings:**

Resistance class

% Resistant plants

Susceptible 0-5%

Low Resistance 6-14%

Moderate Resistance 15-30%

Resistance 31-50%

High Resistance >50%

Not adequately tested

Fall dormancy and disease and insect resistance ratings are from Alfalfa Varieties, a publication of the Certified Alfalfa Seed Council, NAAIC cultivar descriptions, or from developers of the varieties. Blank spaces indicate that the variety has not been tested adequately for that trait.

For those interested in accessing crop performance testing information electronically, visit our World Wide Web site. Most of the information contained in this publication is available for viewing or downloading.

The URL is <http://www.ksu.edu/kscpt>.

**Excerpts from the UNIVERSITY RESEARCH POLICY AGREEMENT  
WITH COOPERATING SEED COMPANIES**

Permission is hereby given to Kansas State University to test varieties and/or hybrids designated on the attached entry forms in the manner indicated in the test announcements. I certify that seed submitted for testing is a true sample of the seed being offered for sale.

I understand that all results from Kansas Crop Performance Tests belong to the University and the public and shall be controlled by the University so as to produce the greatest benefit to the public. Performance data may be used in the following ways: 1) Tables may be reproduced in their entirety provided the source is referenced and data are not manipulated or reinterpreted; 2) Advertising statements by an individual company about the performance of its entries may be made as long as they are accurate statements about the data as published, with no reference to other companies' names or cultivars. In both cases, the following must be included with the reprint or ad citing the appropriate publication number and title: "See the official Kansas State University Agricultural Experiment Station and Cooperative Extension Service Report of Progress 904 '2002 Kansas Performance Tests with Alfalfa Varieties', or the Kansas Crop Performance Test website, <http://www.ksu.edu/kscpt>, for details. Endorsement or recommendation by Kansas State University is not implied."

*These materials may be freely reproduced for educational purposes. All other rights reserved. In each case, give credit to the author(s), name of work, Kansas State University, and the date the work was published.*

## **CONTRIBUTORS**

### **MAIN STATION, MANHATTAN**

Kraig Roozeboom, Associate  
Agronomist (Senior Author)

### **RESEARCH CENTERS**

Pat Evans, Colby  
Joseph Moyer, Mound Valley  
Merle Witt, Garden City

### **EXPERIMENT FIELDS**

Barney Gordon, Belleville  
William Heer, Hutchinson  
Larry Maddux, Powhattan

NOTE: Trade names are used to identify products. No endorsement is intended, nor is any criticism implied of similar products not named.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service, Manhattan 66506  
SRP 904 December 2002

It is the policy of Kansas State University Agricultural Experiment Station and Cooperative Extension Service that all persons shall have equal opportunity and access to its educational programs, services, activities, and materials without regard to race, color, religion, national origin, sex, age, or disability. Kansas State University is an equal opportunity organization. These materials may be available in alternative formats.