



2004

KANSAS PERFORMANCE TESTS WITH

ALFALFA VARIETIES

REPORT OF PROGRESS 935

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service



● dryland ★ irrigated

TABLE OF CONTENTS

2004 Performance Tests

Objectives and Procedures.....	1
Variety Characterization.....	1
Northeast, Topeka, Shawnee County, Seeded 2003	Table 1..... 2
Southeast, Mound Valley, Labette County, Seeded 2001	Table 2..... 3
North Central, Belleville, Republic County, Seeded 2001	Table 3..... 4
South Central, Hutchinson, Reno County, Seeded 2002	Table 4..... 5
Northwest Irrigated, Colby, Thomas County, Seeded 2001	Table 5..... 6
Northwest Irrigated, Colby, Thomas County, Seeded 2003	Table 6..... 7
Southwest Irrigated, Garden City, Finney County, Seeded 2002	Table 7..... 8
Fall dormancy and pest resistance ratings of released varieties, Table 8.....	9
Electronic Access and University Research Policy.....	back cover

Entrants in 2004 Kansas Alfalfa Performance Tests.

AgVenture Seeds, Inc. (AV) Kentland, IN 888-999-0859 agventure.com	Donley Seed Co. (Donley) New Albany, IN 812-941-9822	Monsanto Seed (Monsanto) St. Louis, MO 800-833-5252 monsanto.com	Sharp Bros. Seed Company (Sharp) Healy, KS 800-462-8483 sharpseed.com
Allied Seed Cooperative (Allied) Tangent, OR 866-445-3107 alliedseed.com	Garst Seed Co. (Garst) Slater, IA 800-831-6630 garstseed.com	Mycogen Seeds (Mycogen) Indianapolis, IN 1-800-MYCOGEN mycogen.com	Star Seed, Inc. (Star) Beloit, KS 800-782-7611 gostarseed.com
America's Alfalfa (America's Alfalfa) Princeton, IL 800-873-2532 americasalfalfa.com	Great Plains Research Co. (Great Plains) Apex, NC 919-362-1583 greatplainsresearch.com	NC+ Hybrids (NC+) Lincoln, NE 800-279-7999 nc-plus.com	Syngenta Seeds, Inc. (NK) Golden Valley, MN 763-593-7324 nk-us.com
Bio-Plant Research (BioPlant) Camp Point, IL 800-593-7708	Hyttest Seeds (Hyttest) Shiremantown, PA 717-737-4529	PGI Alfalfa, Inc. (PGI) Oxnard, CA 866-744-5710	Taylor Seed Farms, Inc. (Taylor) White Cloud, KS 800-742-7473 taylorseedfarms.com
Channel Bio Corp. (Channel) Kentland, IN 800-369-8218 channelbio.com	J.C. Robinson Seed Co. (Golden Harvest) Waterloo, NE 800-228-9906 goldenharvestseeds.com	Pioneer, A DuPont Co. (Pioneer) Amarillo, TX 800-258-5604 pioneer.com	United Suppliers Inc. (US Seeds) Eldora, IA 877-714-4503 uniteds.com
CroPlan Genetics (CroPlan Genetics) Shoreview, MN 651-765-5713 croplangenetics.com	J.R. Simplot Company (Simplot) Boise, ID 208-672-2732 simplot.com	Power Seeds, Inc. (Power) Fraserville, Ontario Can 705-944-5600	W-L Research, Inc. (W-L Research) Madison, WI 608-240-0630
Dairyland Seed Co. (Dairyland) West Bend, WI 800-236-0163 dairylandseed.com	Midwest Seed Genetics (Midwest Seed) Carroll, IA 800-369-8218 midwestseed.com	Producers Hybrids (Producers) Battle Creek, NE 402-675-2975 producershybrids.com	

2004 PERFORMANCE TESTS

Objectives and Procedures

The Kansas Agricultural Experiment Station established an official alfalfa testing program in 1980 to provide Kansas growers with unbiased performance comparisons of alfalfa varieties marketed in the state. Each year, private companies are asked to enter varieties voluntarily at the locations slated for establishment that year. Announcements and entry forms are mailed to private companies in June for entry in fall-seeded tests. Companies enter varieties of their choice and pay entry fees to cover part of the costs of conducting the tests. Most tests are planted in mid-August or September, but the Southeast Kansas test usually is planted in the spring. Individual tests are conducted for a minimum of 3 years. New tests typically are established during the final production year of the previous test, or more frequently if there is enough interest.

Descriptive information is presented with the results for each test. This information, including soil type, establishment methods, fertilization, pest control, irrigation, harvest dates, and growing conditions unique to that location, can help explain test and/or variety performance.

Forage yields were estimated by harvesting four replications of each variety with a plot harvester. The amount of forage produced from a specific area (35-80 ft²) was weighed, and a subsample was taken to determine moisture content. This information was used to convert the plot weights to tons of dry matter per acre for each cutting, the season total, and the total for each previous season, as presented in Tables 1 through 7. The forage yield over the lifetime of a particular test is presented as the total tons of dry matter produced per acre, as the total tons of 15% moisture hay, and as a percentage of the test average.

Each table is separated into three sections. The first lists released cultivars that are generally available on the seed market or soon will be. The second section includes experimental cultivars that were entered in the test before being released for sale. These experimental lines often represent an earlier generation of seed than that used for the released cultivars. The third section includes summary statistics unique to that test.

At the bottom of each column, the Least Significant Difference (LSD) is listed at the 0.05 and 0.20 levels. These values indicate how large a difference is needed to be confident that one variety is superior to another. Differences between varieties that are equal to or greater than the 0.05 LSD have only a 1 in 20 chance of being due to chance or error. Differences equal to or greater than the 0.20 LSD have a 1 in 5 chance of being caused by chance or error.

The Coefficient of Variability (CV) provides an estimate of the consistency of the results of a particular test. In these tests, CVs less than 10% generally indicate reliable, uniform data, whereas CVs of 10 to 15% are not uncommon and generally indicate that the data are acceptable for rough comparisons. Tests with CVs greater than 15% may still be useful, but variety comparisons lack precision.

The Mean Coefficient of Variability (MCV) is similar to the CV in that it serves as an indicator of test precision. The MCV is calculated by dividing the 0.05 LSD by the test mean (average) and multiplying by 100. The MCV reveals the percentage difference required to detect differences between varieties with 95% confidence. Many alfalfa breeders and testers agree that tests with MCV values greater than 10% are of little benefit.

Variety Characterization

For variety selection, producers should consider the performance of a variety in each of the current tests in which it appears, its performance over time and locations relative to familiar or check varieties, and the disease and insect resistance characteristics that are potentially important in their situation.

Tables 1 through 7 contain updated yield data from individual tests currently in progress. First-season yields for a spring-planted test are often more variable than yields in subsequent years. Season totals are important, but yield distribution during the season may differ among varieties. Examine yields from individual cuttings to determine if differences in yield distribution exist. Yield totals over many years provide the best measure of variety performance over time.

Table 8 provides disease and insect-resistance ratings. These ratings were obtained primarily from the annual "Winter Survival, Fall Dormancy & Pest Resistance Ratings for Alfalfa Varieties" pamphlet published by the National Alfalfa Alliance. That report summarizes information submitted by developers of alfalfa varieties as part of the variety registration process. The Association of Official Seed Certifying Agencies (AOSCA) National Alfalfa Variety Review Board (NAVRB) reviewed the ratings before they were published. Companies submitting varieties for the tests provided ratings for some unregistered varieties.

Table 1. Northeast Kansas, Topeka Alfalfa Performance Test, Seeded September 9, 2003.

Kansas River Valley Experiment Field, Eudora silt loam
 18 lb. seed/acre
 Plots 3'x20'; 3'x20' harvested
 36-92-216 lb/a of N-P-K before planting

A light infestation of alfalfa weevils was present before the first cutting, but no control measures were required. Rainfall distribution was favorable throughout the growing season.

NAME	Forage Yield						Total, 15% Moist.	Total, % of Mean	
	tons/acre								
	Dry Matter								
	2004				2004	Total			
	5-6	6-9	7-14	9-2					
RELEASED CULTIVARS									
Power 4.2	2.01	1.89	1.64	1.67	7.21	7.21	8.48	105	
Notice II	2.09	1.79	1.60	1.70	7.18	7.18	8.45	105	
6400HT	1.98	1.83	1.65	1.64	7.10	7.10	8.35	104	
Perry	2.01	1.77	1.56	1.64	6.98	6.98	8.22	102	
4A421	1.82	1.80	1.58	1.70	6.90	6.90	8.12	101	
Phirst	1.88	1.77	1.52	1.69	6.86	6.86	8.07	100	
6530	1.78	1.79	1.57	1.63	6.76	6.76	7.96	99	
Reward II	1.77	1.73	1.51	1.75	6.75	6.75	7.95	99	
WL 357 HQ	1.90	1.74	1.52	1.54	6.71	6.71	7.89	98	
Journey 204 Hybrid Alfalfa	1.81	1.79	1.49	1.61	6.70	6.70	7.88	98	
HybriForce-420/wet	1.80	1.78	1.55	1.58	6.70	6.70	7.88	98	
Kanza	1.84	1.73	1.50	1.60	6.67	6.67	7.85	97	
SUMMARY STATISTICS									
Average	1.86	1.78	1.56	1.65	6.85	6.85	8.06	100	
LSD(0.05)	0.28	NS	NS	NS	NS	NS	NS	NS	
LSD(0.20)	0.18	0.10	0.09	0.10	0.47	0.47	0.55	7	
CV(%)	10.46	6.26	5.94	6.32	3.86	3.86	3.86	4	
MCV(%)	15.02	NS	NS	NS	NS	NS	NS	NS	

Table 2. Southeast Kansas, Mound Valley Alfalfa Performance Test, Seeded May 9, 2001.

Southeast Ag. Research Center, Parsons silty clay loam					Spring rains delayed the first cutting slightly. The third cutting was accelerated because of a blister beetle infestation. Dry weather inhibited regrowth after the fourth harvest.						
15 lb. seed/acre											
Plots 5'x30'; 3'x20' harvested											
20-50-200 lb/a of N-P-K on March 10, 2004											
NAME	Forage Yield									Total, 15% Moist.	Total, % of Mean
	tons/acre										
	Dry Matter										
	2004				2004	2003	2002	2001	Total		
5-10	6-8	7-12	8-8								
RELEASED CULTIVARS											
6420	2.32	1.13	1.32	1.04	5.81	5.23	5.36	1.21	17.61	20.72	107
HybriForce-400	2.15	1.12	1.40	1.05	5.72	5.08	5.42	1.34	17.56	20.65	107
Dagger+EV	1.97	1.07	1.21	0.91	5.16	5.19	5.08	1.29	16.72	19.67	101
Perry	2.29	1.02	1.37	0.99	5.67	4.83	5.03	1.07	16.60	19.53	101
5-Star	1.98	1.14	1.18	1.05	5.35	5.07	4.89	1.17	16.48	19.39	100
WL 327	2.07	1.12	1.24	0.98	5.40	5.26	4.70	1.12	16.48	19.38	100
Rebound 4.2	2.09	1.10	1.31	0.92	5.41	5.11	4.80	1.02	16.34	19.23	99
400SCL	2.05	0.98	1.23	0.90	5.15	5.12	4.94	1.04	16.25	19.12	99
Pioneer 54V54	2.04	1.04	1.17	0.98	5.23	4.96	5.04	1.01	16.24	19.11	99
WL 342	2.20	1.00	1.21	0.96	5.37	4.99	4.77	1.08	16.21	19.07	98
Pawnee	1.76	1.04	1.25	0.86	4.91	5.14	4.98	1.13	16.16	19.01	98
350	1.85	0.96	1.30	0.97	5.09	4.86	4.82	1.12	15.89	18.70	96
Kanza	1.75	1.06	1.26	1.02	5.10	4.83	4.70	1.25	15.88	18.68	96
SUMMARY STATISTICS											
Average	2.04	1.06	1.27	0.97	5.34	5.04	4.96	1.14	16.48	19.38	100
LSD(0.05)	0.26	NS	NS	NS	0.71	0.27	NS	0.15	1.66	1.96	10
LSD(0.20)	0.17	0.08	0.12	0.09	0.46	0.21	0.32	0.09	1.07	1.26	6
CV(%)	8.90	8.68	9.80	8.68	4.74	4.45	6.94	8.96	3.00	3.00	3
MCV(%)	12.77	NS	NS	NS	13.30	5.32	NS	12.85	10.09	10.09	10

Table 3. North Central Kansas, Belleville Alfalfa Performance Test, Seeded August 22, 2001.

North Central Kansas Exp. Field, Crete silt loam
 18 lb. seed/acre
 Plots 5'x15'; 3'x15' harvested
 11-50-0 lb/a of N-P-K applied in February and after first cutting

The first three cuttings benefited from normal to above-normal rains. August and September were extremely dry, generating little regrowth after the fourth cutting.

NAME	Forage Yield								Total, 15% Moist.	Total, % of Mean
	tons/acre									
	Dry Matter									
	2004				2004	2003	2002	Total		
5-20	6-14	7-14	8-11							
RELEASED CULTIVARS										
Abundance	2.50	1.64	1.51	0.37	6.02	5.53	4.16	15.71	18.48	107
Dagger+EV	2.51	1.60	1.71	0.40	6.22	5.35	3.81	15.38	18.09	105
645-II	2.41	1.48	1.35	0.38	5.61	5.58	4.16	15.35	18.06	105
GH 750	2.33	1.24	1.54	0.31	5.41	5.81	3.88	15.10	17.76	103
Feast+EV	2.54	1.26	1.49	0.33	5.62	5.32	4.07	15.01	17.66	103
DKA42-15	2.51	1.34	1.49	0.39	5.72	5.37	3.88	14.97	17.62	102
Pawnee	2.28	1.40	1.47	0.39	5.54	4.95	4.48	14.97	17.61	102
US A4230	2.41	1.32	1.54	0.34	5.61	5.44	3.91	14.96	17.60	102
A 30-06	2.44	1.25	1.44	0.38	5.51	5.45	3.98	14.94	17.58	102
Journey 204 Hybrid Alfalfa	2.53	1.40	1.58	0.39	5.91	5.21	3.81	14.93	17.56	102
Pioneer 54Q53	2.49	1.31	1.63	0.40	5.84	5.29	3.79	14.92	17.55	102
Macon	2.45	1.53	1.48	0.37	5.82	5.31	3.74	14.87	17.50	102
6410	2.41	1.29	1.44	0.34	5.48	5.35	3.92	14.75	17.36	101
Enhancer	2.27	1.22	1.43	0.38	5.31	5.38	3.94	14.63	17.22	100
Abilene+Z	2.54	1.25	1.51	0.35	5.66	5.02	3.94	14.62	17.19	100
Perry	2.62	1.37	1.39	0.41	5.79	5.68	3.13	14.60	17.18	100
FK421	2.53	1.23	1.45	0.27	5.48	5.35	3.77	14.60	17.17	100
Lightning II	2.30	1.22	1.42	0.41	5.34	5.35	3.86	14.55	17.12	100
Geneva	2.44	1.30	1.35	0.39	5.48	5.05	3.86	14.39	16.93	98
Ameristand 403T	2.42	1.36	1.28	0.25	5.31	5.04	3.91	14.26	16.77	97
HybriForce-400	2.30	1.27	1.30	0.40	5.26	4.85	4.09	14.20	16.71	97
Reliance	2.32	1.42	1.44	0.35	5.53	4.56	3.94	14.03	16.50	96
Kanza	2.50	1.61	1.46	0.35	5.92	4.40	3.17	13.49	15.87	92
EXPERIMENTAL STRAINS										
ZC9840A	2.62	1.55	1.55	0.36	6.07	6.02	4.16	16.25	19.12	111
ZG9941	2.62	1.18	1.55	0.33	5.68	5.31	3.81	14.80	17.41	101
ZC9851A	2.65	1.37	1.52	0.39	5.94	4.74	4.07	14.75	17.35	101
ZG9840	2.47	1.27	1.37	0.27	5.38	5.39	3.95	14.72	17.32	101
ZC9940A	2.63	1.25	1.40	0.37	5.65	4.98	3.78	14.41	16.96	99
ZC9950A	2.54	1.24	1.43	0.33	5.53	5.20	3.65	14.38	16.92	98
4M72	2.25	1.09	1.43	0.41	5.19	5.36	3.74	14.29	16.81	98
ZC9941A	2.52	1.19	1.32	0.42	5.44	4.92	3.88	14.24	16.76	97
ZC9841A	2.42	1.26	1.36	0.42	5.46	4.94	3.70	14.10	16.59	96
5M87	2.26	1.21	1.52	0.36	5.34	5.27	3.47	14.08	16.56	96
ZC9942A	2.47	1.26	1.32	0.27	5.31	4.96	3.73	14.00	16.48	96
ZC9842A	2.42	1.35	1.38	0.38	5.53	4.78	3.35	13.66	16.07	93
ZC9953A	2.20	1.15	1.43	0.31	5.08	4.51	3.85	13.44	15.82	92
SUMMARY STATISTICS										
Average	2.45	1.32	1.45	0.36	5.58	5.20	3.84	14.62	17.20	100
LSD(0.05)	0.27	0.17	0.21	NS	0.77	0.43	0.41	1.70	2.00	12
LSD(0.20)	0.17	0.11	0.14	0.07	0.50	0.34	0.27	1.11	1.30	8
CV(%)	7.76	9.42	10.44	22.61	5.10	7.08	7.62	3.75	3.75	4
MCV(%)	10.88	13.21	14.63	NS	13.75	8.31	10.68	11.62	11.62	12

Table 4. South Central Kansas, Hutchinson Alfalfa Performance Test, Seeded September 1, 2002.

South Central Experiment Field, Ost silt loam
 18 lb. seed/acre
 Plots 5'x20', 3'x20' harvested
 75-40-40 lb/a of N-P-K before planting

Above-normal rainfall in June and July, but late August and September were dry. Temperatures were generally below normal. Insect pests caused no problems.

NAME	Forage Yield								Total, 15% Moist.	Total, % of Mean
	tons/acre									
	Dry Matter									
	2004				2004	2003	Total	Total		
5-24	6-21	7-15	8-28							
RELEASED CULTIVARS										
350	2.23	0.77	1.26	1.32	5.58	4.27	9.85	11.59	108	
HybriForce-400	2.36	0.81	1.37	1.24	5.78	3.95	9.73	11.44	107	
Hyttest 410	2.34	0.86	1.17	1.21	5.59	3.99	9.58	11.27	105	
400SCL	2.33	0.80	1.13	1.07	5.33	4.19	9.52	11.20	105	
Journey 204 Hybrid Alfalfa	2.13	0.77	1.14	1.27	5.31	4.10	9.41	11.08	104	
Dagger+EV	2.25	0.79	1.38	1.21	5.63	3.78	9.41	11.08	104	
Perry	2.38	0.67	1.14	1.19	5.39	3.91	9.30	10.94	102	
Reliance	2.00	0.73	1.11	1.19	5.02	4.27	9.29	10.93	102	
Reward II	2.19	0.74	1.19	1.16	5.28	4.00	9.28	10.92	102	
WL 342	2.03	0.73	1.13	1.20	5.09	4.14	9.23	10.86	102	
Pawnee	2.03	0.85	1.19	1.37	5.44	3.78	9.22	10.85	102	
DKA42-15	2.15	0.75	1.30	1.21	5.41	3.80	9.21	10.84	101	
Aspire	2.36	0.84	1.19	1.32	5.70	3.38	9.08	10.69	100	
645-II	2.16	0.72	1.02	1.04	4.94	4.14	9.08	10.68	100	
Rebound 4.2	1.94	0.77	1.26	1.12	5.10	3.88	8.98	10.56	99	
5-Star	1.95	0.84	1.35	1.21	5.34	3.54	8.88	10.45	98	
Macon	1.99	0.74	1.16	1.05	4.93	3.88	8.81	10.36	97	
Key	2.04	0.73	0.95	1.22	4.94	3.61	8.55	10.06	94	
Lightning II	1.56	0.73	1.07	1.18	4.53	3.73	8.26	9.72	91	
Kanza	1.78	0.80	1.11	1.21	4.89	3.31	8.20	9.65	90	
EXPERIMENTAL STRAINS										
CW 83018	2.18	0.85	1.31	1.10	5.44	3.73	9.17	10.79	101	
CW 93018	2.02	0.76	1.19	1.22	5.19	3.78	8.97	10.56	99	
CW 94006	1.75	0.74	1.16	1.18	4.82	3.65	8.47	9.97	93	
CW 94022	1.90	0.72	1.14	1.11	4.87	3.54	8.41	9.90	93	
SUMMARY STATISTICS										
Average	2.08	0.77	1.18	1.19	5.23	3.85	9.08	10.69	100	
LSD(0.05)	0.44	NS	NS	NS	1.14	0.49	1.62	1.90	18	
LSD(0.20)	0.29	NS	NS	NS	0.74	0.32	1.05	1.23	12	
CV(%)	15.04	16.41	17.69	13.22	8.18	8.95	6.06	6.06	6	
MCV(%)	21.22	NS	NS	NS	21.76	12.62	17.81	17.81	18	

Table 5. Northwest Kansas, Colby Alfalfa Performance Test, Seeded August 29, 2001.

Northwest Res.-Ext. Center, Keith silt loam	Good growing conditions prevailed throughout the summer, with above-normal rainfall and below-normal temperatures. Insects and diseases caused no problems.									
18 lb. seed/acre										
Plots 3'x20'; 3'x17' harvested										
17-60-0 lb/a of N-P-K before planting										
	Forage Yield								Total, 15% Moist.	Total, % of Mean
	tons/acre									
	Dry Matter									
	2004				2004	2003	2002	Total		
NAME	6-1	7-12	8-6	9-15						
RELEASED CULTIVARS										
Pioneer 53V08	4.18	2.16	1.26	1.57	9.18	7.67	6.51	23.36	27.48	109
A 30-06	3.57	2.09	1.28	1.48	8.42	7.86	6.77	23.05	27.12	108
Enhancer	3.71	2.13	1.35	1.60	8.79	7.20	6.65	22.64	26.64	106
631	3.84	2.07	1.25	1.47	8.63	7.23	6.73	22.59	26.58	106
Lightning II	3.19	1.86	1.35	1.51	7.91	7.28	6.89	22.08	25.98	103
Kanza	3.63	2.06	1.41	1.66	8.76	6.91	6.05	21.72	25.55	102
Magnum V	3.38	2.05	1.29	1.40	8.12	6.80	6.76	21.68	25.51	102
Target II Plus	3.29	2.04	1.18	1.45	7.96	7.06	6.61	21.63	25.44	101
Reward	3.50	2.19	1.36	1.48	8.53	6.91	6.17	21.61	25.43	101
645-II	3.25	1.94	1.18	1.26	7.63	6.99	6.64	21.26	25.02	100
Geneva	3.27	2.20	1.34	1.47	8.28	6.99	5.94	21.21	24.95	99
Pioneer 54Q53	2.95	1.91	1.24	1.42	7.51	7.25	6.29	21.05	24.77	99
Feast+EV	3.19	1.96	1.19	1.39	7.73	7.15	5.75	20.63	24.27	97
4200	2.93	1.94	1.34	1.46	7.66	6.48	5.95	20.09	23.64	94
Perry	3.17	1.82	1.02	1.23	7.25	6.91	5.18	19.34	22.75	91
EXPERIMENTAL STRAINS										
ZC9842A	3.53	2.03	1.42	1.50	8.48	7.51	6.43	22.42	26.38	105
CW 94008	3.55	2.28	1.30	1.51	8.65	7.35	6.37	22.37	26.32	105
6M71	3.49	2.03	1.41	1.59	8.51	7.31	6.38	22.20	26.12	104
CW 64004	3.51	2.21	1.31	1.46	8.48	7.41	6.19	22.08	25.98	103
ZC9851A	3.63	2.22	1.35	1.47	8.67	7.05	6.28	22.00	25.88	103
5M85	3.38	2.16	1.27	1.55	8.37	7.15	6.24	21.76	25.59	102
ZC9840A	3.39	1.78	1.28	1.33	7.78	7.38	6.50	21.66	25.48	101
ZC9854A	3.52	2.00	1.39	1.44	8.36	6.94	6.22	21.52	25.32	101
5M84	2.89	2.19	1.39	1.51	7.98	7.22	6.32	21.52	25.32	101
ZC9950A	3.06	2.34	1.27	1.49	8.16	6.93	6.35	21.44	25.23	100
CW 75044	3.26	1.89	1.41	1.53	8.09	7.06	6.23	21.38	25.16	100
ZC9940A	3.22	2.04	1.23	1.38	7.87	7.35	6.00	21.22	24.97	99
ZC9841A	3.20	2.11	1.33	1.39	8.03	6.72	6.43	21.18	24.92	99
CW 74040	3.20	2.10	1.30	1.34	7.94	7.14	5.84	20.92	24.61	98
CW 64049	2.69	1.98	1.14	1.49	7.29	6.72	5.80	19.81	23.31	93
CW 64026	2.63	1.80	1.32	1.37	7.11	6.30	6.00	19.41	22.83	91
CW 73029	2.91	1.82	1.18	1.33	7.24	6.67	5.37	19.28	22.68	90
CW 54033	2.40	1.70	1.20	1.28	6.58	6.49	5.57	18.64	21.93	87
SUMMARY STATISTICS										
Average	3.29	2.03	1.29	1.45	8.06	7.07	6.22	21.35	25.12	100
LSD(0.05)	0.76	NS	0.17	0.21	1.54	0.65	0.47	2.68	3.15	13
LSD(0.20)	0.50	NS	0.11	0.14	1.00	0.43	0.31	1.74	2.05	8
CV(%)	16.53	14.08	9.13	10.55	7.99	6.60	5.43	4.04	4.04	4
MCV(%)	23.20	NS	12.82	14.80	19.16	9.26	7.62	12.53	12.53	13

Table 6. Northwest Kansas, Colby Alfalfa Performance Test, Seeded August 29, 2003.

Northwest Research-Extension Center, Keith silt loam 18 lb. seed/acre Plots 3'x20'; 3'x17' harvested 16-55-0 lb/a of N-P-K before planting	Good growing conditions prevailed throughout the summer, with above-normal rainfall and below-normal temperatures. Insects and diseases caused no problems.
---	---

NAME	Forage Yield						Total, 15% Moist.	Total, % of Mean
	tons/acre							
	Dry Matter							
	2004				2004	Total		
	6-2	7-13	8-9	9-16				
RELEASED CULTIVARS								
Phirst	3.62	1.96	1.50	1.14	8.22	8.22	9.66	105
Pioneer 54Q25	3.47	2.07	1.46	1.18	8.17	8.17	9.61	105
Pioneer 54V46	3.23	2.07	1.59	1.28	8.17	8.17	9.61	105
Arapaho	3.62	1.88	1.35	1.14	8.00	8.00	9.41	103
Notice II	3.16	2.11	1.39	1.29	7.96	7.96	9.36	102
WL 357 HQ	3.56	1.85	1.42	1.09	7.93	7.93	9.33	102
Expedition	3.27	2.03	1.47	1.10	7.88	7.88	9.27	101
Journey 204 Hybrid Alfalfa	3.34	1.84	1.52	1.13	7.83	7.83	9.21	101
Regal	3.29	1.94	1.33	1.25	7.82	7.82	9.20	100
Kanza	3.14	2.03	1.45	1.16	7.77	7.77	9.14	100
Jade III	3.29	1.85	1.44	1.11	7.69	7.69	9.05	99
6400HT	3.47	1.79	1.42	1.01	7.69	7.69	9.05	99
HybriForce-420/wet	3.46	1.75	1.31	1.12	7.65	7.65	9.00	98
Abundance	3.34	1.87	1.29	1.15	7.65	7.65	9.00	98
Maximizer	3.20	1.83	1.35	1.11	7.48	7.48	8.80	96
Evermore	2.84	2.08	1.46	1.07	7.44	7.44	8.76	96
Perry	3.27	1.84	1.33	1.01	7.44	7.44	8.76	96
631	2.83	2.02	1.42	1.16	7.43	7.43	8.74	95
EXPERIMENTAL STRAINS								
FSG351	3.38	1.89	1.50	1.19	7.96	7.96	9.37	102
FSG505	3.42	1.81	1.46	1.16	7.84	7.84	9.23	101
FSG406	3.23	1.95	1.45	1.14	7.77	7.77	9.14	100
CL2000	3.21	1.95	1.27	1.20	7.62	7.62	8.97	98
SUMMARY STATISTICS								
Average	3.30	1.93	1.42	1.14	7.79	7.79	9.17	100
LSD(0.05)	NS	NS	0.18	0.13	NS	NS	NS	NS
LSD(0.20)	NS	NS	0.12	0.08	0.84	0.84	0.99	11
CV(%)	14.12	12.04	9.20	7.78	6.99	6.99	6.99	7
MCV(%)	NS	NS	13.01	11.00	NS	NS	NS	NS

Table 7. Southwest Kansas, Garden City Alfalfa Performance Test, Seeded September 3, 2002.

NAME	Forage Yield								Total, 15% Moist.	Total, % of Mean
	tons/acre									
	Dry Matter									
	2004				2004	2003	Total	Total		
5-20	6-29	8-20	9-16							
RELEASED CULTIVARS										
WL 327	4.02	2.20	1.93	1.97	10.11	9.35	19.46	22.90	108	
Reward II	3.82	2.11	1.88	1.91	9.72	9.17	18.89	22.23	105	
GH 750	3.94	2.13	1.85	1.87	9.79	9.03	18.82	22.14	104	
WL 342	3.69	2.09	1.89	1.84	9.51	9.14	18.65	21.94	103	
Hyttest 410	3.74	2.04	1.90	1.92	9.60	9.04	18.64	21.93	103	
Abundance	3.66	2.10	1.87	1.95	9.58	9.00	18.58	21.86	103	
HybriForce-400	3.68	2.05	1.80	1.90	9.43	9.11	18.54	21.81	103	
Key	3.87	2.19	1.77	1.86	9.68	8.82	18.50	21.77	103	
4A421	3.63	2.13	1.92	1.82	9.49	9.00	18.49	21.75	103	
WL 319 HQ	3.65	2.10	1.84	1.77	9.35	9.11	18.46	21.72	102	
Journey 204 Hybrid Alfalfa	3.73	2.14	1.86	1.94	9.67	8.75	18.42	21.67	102	
Dagger+EV	3.71	2.09	1.84	1.89	9.52	8.70	18.22	21.43	101	
5-Star	3.61	2.13	1.89	1.79	9.42	8.73	18.15	21.36	101	
Hyttest 520	3.54	2.08	1.93	1.85	9.39	8.68	18.07	21.26	100	
Pioneer 54V54	3.65	2.08	1.82	1.80	9.36	8.61	17.97	21.14	100	
Masterpiece	3.54	2.07	1.78	1.85	9.24	8.69	17.93	21.09	99	
Feast+EV	3.67	2.04	1.72	1.80	9.24	8.56	17.80	20.94	99	
Magna 601	3.39	2.02	1.92	1.93	9.27	8.28	17.55	20.65	97	
Kanza	3.08	1.83	1.66	1.76	8.33	6.95	15.28	17.97	85	
EXPERIMENTAL STRAINS										
DS108HYB	4.02	2.15	1.87	1.92	9.96	8.79	18.75	22.06	104	
DS9809HYB	3.89	2.15	1.88	1.89	9.81	8.88	18.69	21.99	104	
ZC9953A	3.86	2.13	1.84	1.91	9.73	8.95	18.68	21.98	104	
DS218HYB	3.77	2.17	2.02	2.08	10.04	8.57	18.61	21.90	103	
DS107HYB	3.62	2.10	1.84	1.91	9.46	9.13	18.59	21.88	103	
DS201HYB	3.74	2.16	2.01	2.07	9.98	8.55	18.53	21.80	103	
DS106HYB	3.42	2.14	1.89	2.01	9.46	9.05	18.51	21.77	103	
CW 04022	3.51	2.15	1.90	1.90	9.45	8.80	18.25	21.47	101	
GPVL0144	3.77	2.08	1.78	1.88	9.52	8.30	17.82	20.96	99	
Pioneer 55V05	3.51	2.12	1.88	1.94	9.45	8.36	17.81	20.95	99	
CW 94025	3.44	2.01	1.80	1.80	9.06	8.65	17.71	20.83	98	
CW 14026	3.50	2.06	1.90	1.87	9.32	8.36	17.68	20.80	98	
CW 04027	3.49	2.02	1.83	1.85	9.20	8.46	17.66	20.77	98	
CW 04030	3.48	1.97	1.81	1.83	9.08	8.49	17.57	20.67	97	
CW 65086	3.32	2.00	1.86	1.94	9.12	8.43	17.55	20.65	97	
FG 40M159A	3.42	2.00	1.81	1.69	8.93	8.53	17.46	20.54	97	
Exp 80I	3.50	1.95	1.69	1.79	8.93	8.45	17.38	20.45	96	
CW 05009	3.27	2.04	1.91	1.98	9.20	8.12	17.32	20.37	96	
CW 94023	3.40	1.95	1.78	1.72	8.85	8.23	17.08	20.09	95	
CW 65085	3.28	1.96	1.70	1.75	8.69	8.12	16.81	19.78	93	
SUMMARY STATISTICS										
Average	3.60	2.06	1.84	1.87	9.37	8.66	18.03	21.21	100	
LSD(0.05)	0.54	0.16	0.14	0.20	1.04	0.30	1.40	1.65	8	
LSD(0.20)	0.35	0.10	0.09	0.13	0.68	0.23	0.91	1.07	5	
CV(%)	10.79	5.52	5.43	7.72	4.70	2.92	2.80	2.80	3	
MCV(%)	15.11	7.73	7.60	10.81	11.15	3.46	7.77	7.77	8	

Table 8. Varieties in 2004 Kansas Alfalfa Performance Tests, with disease and insect resistance ratings.*

Brand Name	P S B A R R											Brand Name	P S B A R R														
	B V F A R A P A S P K K P	W W W N R A A A N H N N L	Brand Name	B V F A R A P A S P K K P	W W W N R A A A N H N N L																						
Allied												Midwest Seed															
350	H	H	H	H	H	R	R	-	-	H	-	-	Pawnee	H	H	H	H	H	M	H	L	M	H	-	L	L	
400SCL	-	-	-	-	-	-	-	-	-	-	-	-	Monsanto														
Evermore	H	H	H	H	H	H	R	-	R	H	-	M	M	Aspire	M	R	H	H	H	H	H	R	H	-	-	-	
Macon	H	H	H	H	H	R	R	-	M	H	-	-	-	DKA42-15	H	H	H	H	H	R	H	-	R	H	-	-	
Reliance	H	H	H	H	H	R	-	-	M	R	-	-	-	Mycogen													
America's Alfalfa													4A421	H	H	H	H	H	H	H	-	-	H	-	M	M	
Abilene+Z	H	H	H	H	H	M	H	M	R	R	-	-	-	NC+													
Ameristand 403T	H	H	H	H	H	M	R	-	M	H	-	-	-	Jade III	H	R	H	H	H	R	R	R	R	R	-	H	H
AV													NK														
4200	-	-	-	-	-	-	-	-	-	-	-	-	Expedition	R	H	H	H	H	R	-	-	R	H	-	R	R	
BioPlant													Geneva	H	H	H	H	H	R	H	L	R	H	-	-	-	
Phirst	H	R	H	H	H	R	R	-	R	R	-	H	H	PGI													
Channel													A 30-06	H	H	H	H	H	-	R	-	-	H	-	-	-	
Notice II	-	-	-	-	-	-	-	-	-	-	-	-	Reward	H	R	H	R	H	R	H	M	M	M	-	-	-	
CroPlan Genetics													Reward II	H	R	H	R	H	R	R	R	R	R	-	H	H	
5-Star	R	R	H	R	R	R	R	R	R	R	-	-	-	Pioneer													
Rebound 4.2	H	H	H	H	H	R	R	-	M	H	-	-	-	53V08	H	H	H	H	H	R	H	M	H	L	-	H	H
Dairyland													54Q25	H	H	H	H	H	R	R	-	H	R	-	H	H	
Arapaho	H	R	H	R	H	-	M	-	R	R	-	H	H	54Q53	H	H	R	R	H	M	M	-	H	M	-	H	H
HybriForce-400	H	R	H	R	H	H	R	M	R	M	-	H	H	54V46	R	H	H	H	H	R	M	L	M	R	-	H	H
HybriForce-420/wet	H	R	H	R	H	R	R	-	H	R	-	H	H	54V54	H	H	H	H	H	R	-	-	L	M	-	-	-
Magna 601	R	M	H	R	H	H	R	-	R	M	R	R	R	Power													
Magnum V	H	R	H	R	H	R	R	M	R	M	-	M	M	Power 4.2	H	R	H	R	H	R	R	-	H	H	-	R	R
Donley													Producers														
FK421	H	H	H	H	H	-	R	-	-	H	-	-	-	Target II Plus	H	R	H	R	H	M	R	M	R	M	-	M	M
Garst													Sharp														
631	H	R	H	R	H	R	H	M	R	M	-	-	-	Abundance	H	R	H	R	H	R	R	M	R	R	-	H	H
6400HT	H	H	H	H	H	-	H	-	-	H	-	-	-	Enhancer	H	R	H	R	H	R	R	M	M	M	-	M	M
6410	H	H	H	H	H	R	R	-	-	H	-	-	-	Journey 204 Hybrid	H	R	H	H	H	R	R	-	R	R	-	H	H
6420	H	R	H	R	H	R	R	-	R	R	-	H	H	Simplot													
645-II	H	H	H	H	H	-	R	-	-	H	-	-	-	Masterpiece	H	R	H	H	H	R	-	R	H	R	-	R	R
6530	H	H	H	H	H	-	H	-	R	H	-	-	-	Star													
Dagger+EV	H	H	H	H	H	M	H	M	R	H	-	L	L	Lightning II	H	R	H	H	H	H	M	-	M	H	-	-	-
Feast+EV	H	H	H	R	H	-	M	-	-	H	-	-	-	Taylor													
Golden Harvest													Maximizer	H	H	H	H	H	R	-	-	R	H	-	-	-	
GH 750	H	H	H	H	H	R	R	-	M	H	-	-	-	US Seeds													
Great Plains													US A4230	H	H	H	H	H	R	R	-	R	H	-	-	-	
Key	H	H	H	H	H	H	H	M	M	M	M	-	-	W-L Research													
Regal	H	R	H	R	H	R	H	-	H	M	M	-	-	WL 319 HQ	H	H	H	H	H	R	H	-	M	H	-	-	-
Hyttest													WL 327	H	R	H	H	H	R	R	H	R	H	-	-	-	
Hyttest 410	H	H	H	H	H	H	M	-	R	H	-	-	-	WL 342	H	H	H	H	H	H	H	-	R	H	-	-	-
Hyttest 520	-	-	-	-	-	-	-	-	-	-	-	-	-	WL 357 HQ	H	H	H	H	H	-	H	-	-	H	-	-	-

*BW = Bacterial wilt
 VW = Verticillium wilt
 FW = Fusarium wilt
 AN = Anthracnose race 1
 PRR = Phytophthora root rot
 SAA = Spotted alfalfa aphid
 PA = Pea aphid

BAA = Blue alfalfa aphid
 SN = Stem nematode
 APH = Aphanomyces root rot race 1
 SRKN = Southern root knot nematode
 NRKN = Northern root knot nematode
 PL = Potato leafhopper

Pest resistance ratings:		
Code	Resistance class	% Resistant plants
S	Susceptible	0-5%
L	Low Resistance	6-14%
M	Moderate Resistance	15-30%
R	Resistance	31-50%
H	High Resistance	>50%
-	Not adequately tested	

Disease and insect resistance ratings are from the National Alfalfa Alliance, NAAIC descriptions, or from developers of the varieties.

For those interested in accessing crop performance testing information electronically, visit our World Wide Web site. Most of the information contained in this publication is available for viewing or downloading.

The URL is <http://www.ksu.edu/kscpt>.

Excerpts from the
UNIVERSITY RESEARCH POLICY AGREEMENT WITH COOPERATING SEED COMPANIES

Permission is hereby given to Kansas State University to test varieties and/or hybrids designated on the attached entry forms in the manner indicated in the test announcements. I certify that seed submitted for testing is a true sample of the seed being offered for sale.

I understand that all results from Kansas Crop Performance Tests belong to the University and the public and shall be controlled by the University so as to produce the greatest benefit to the public. Performance data may be used in the following ways: 1) Tables may be reproduced in their entirety provided the source is referenced and data are not manipulated or reinterpreted; 2) Advertising statements by an individual company about the performance of its entries may be made as long as they are accurate statements about the data as published, with no reference to other companies' names or cultivars. In both cases, the following must be included with the reprint or ad citing the appropriate publication number and title: "See the official Kansas State University Agricultural Experiment Station and Cooperative Extension Service Report of Progress 935 '2004 Kansas Performance Tests with Alfalfa Varieties', or the Kansas Crop Performance Test website, <http://www.ksu.edu/kscpt>, for details. Endorsement or recommendation by Kansas State University is not implied."

These materials may be freely reproduced for educational purposes. All other rights reserved. In each case, give credit to the author(s), name of work, Kansas State University, and the date the work was published.

CONTRIBUTORS

MAIN STATION, MANHATTAN

Kraig Roozeboom, Agronomist
(Senior Author)

RESEARCH CENTERS

Pat Evans, Colby
Joseph Moyer, Mound Valley
Merle Witt, Garden City

EXPERIMENT FIELDS

Barney Gordon, Belleville
William Heer, Hutchinson
Larry Maddux, Topeka

NOTE: Trade names are used to identify products. No endorsement is intended, nor is any criticism implied of similar products not named.