

SOME EFFECTS OF GIBBERELIC ACID
ON YEAR-OLD PECAN SEEDLINGS

by

LOY WILLIAM SHREVE

B. S. F., West Virginia University, 1951

A MASTER'S THESIS

submitted in partial fulfillment of the

requirements for the degree

MASTER OF SCIENCE

Department of Horticulture and Forestry

KANSAS STATE UNIVERSITY
Manhattan, Kansas

1967

Approved by


Major Professor

LD
2668
T4
1967
S518
c.2

TABLE OF CONTENTS

	PAGE
INTRODUCTION.....	1
Definition of Terms.....	2
REVIEW OF LITERATURE.....	2
MATERIALS AND METHODS.....	5
RESULTS.....	7
DISCUSSION.....	10
SUMMARY.....	14
ACKNOWLEDGEMENTS.....	16
LITERATURE CITED.....	17

TABLES

TABLE	PAGE
I. Some Effects on Growth of Year-old Pecan Seedling Trees From Aqueous Sprays of GA ₃ at 5,000 ppm in 1966. Treatment Groups Included Trees Sprayed in June or July and Those Sprayed Both Months.....	8
II. Comparison of Effects of Spray Applications of 5,000 ppm GA ₃ on Growth and Development of Year-old Pecan Seedlings.....	9

INTRODUCTION

The slow growth of the pecan seedling during its early years is a problem for nurserymen and others interested in pecan propagation. Under normal growing conditions, pecan seedlings may grow for three or four seasons before the diameter of the stem is large enough to be suitable for grafting or budding. Reducing the time required to grow pecan seedlings to a size convenient for propagation would enable nurserymen to produce planting stock at reduced costs. Also, pecan orchard owners could benefit from lower priced planting stock, and, if the owners do the propagations themselves, they could gain in the shorter time required for trees to begin bearing nuts.

Gibberellic acid causes increased plant height of year-old seedlings of yellow poplar, sweetgum, cherrybark oak, willow oak, and southern red oak (12). GA_3 also caused increased height growth in rooted cuttings of eastern cottonwood and American sycamore. Treatment with GA_3 caused increased stem elongation of newly germinated pecan seedlings grown under greenhouse conditions (9). After treatment with GA_3 , greenhouse grown seedlings of black walnut, willow oak, and loblolly pine had stems 40% taller and twice as thick as the controls (16).

Objectives of this study were to determine: 1, if field applications of gibberellic acid increased pecan seedling growth to a size suitable for budding and grafting earlier than untreated pecan seedlings; 2, if time of application affected response of pecan seedlings to GA_3 ; and, 3, how pecan seedlings responded to

repeated treatment with GA₃.

DEFINITION OF TERMS

Gibberellin. A general term for any of the plant stimulating substances produced by the Gibberella fungus.

Gibberellic Acid. The gibberellin also classified as GA₃, the type most abundantly secreted by the fungus (Gibberella) and easiest to extract.

REVIEW OF LITERATURE

Nelson (12) found that a one percent solution of gibberellic acid in lanolin applied to terminal buds of year-old seedlings of yellow-poplar, sweetgum, cherrybark oak, willow oak and southern red oak significantly increased their height. The same solution applied to terminal buds of cuttings of eastern cottonwood and American sycamore also caused significant increases in height growth of these species. Stems of newly germinated white oak seedlings showed less marked increases as did year-old seedlings of white pine from treatments with GA₃.

Kenworthy (6) reported that treatment in excess of 1,000 ppm GA₃ dissolved in methanol and introduced directly into the xylem of the tree trunk of young peach trees resulted in rosetting of twigs and reduced growth by a decrease in twig length, leaf area, and dry weight of twigs. One hundred ppm GA₃ treatment applied in the same way resulted in increased twig length, leaf area, and dry weight of leaves.

Bilan and Kemp (2,3) observed that year-old loblolly pine seedlings treated with one, two, or three per cent solutions of gibberellic acid applied by soaking shoots only, roots only, or soaking the entire plant increased the height of the tree with increases being proportional to concentrations used.

Martin and Wiggins (9) have found that soaking pecan seeds in gibberellic acid at concentrations up to 5,000 ppm resulted in earlier germination. Treated seed had better germination, and the resulting seedlings' height was greater than control. They found that GA₃ caused only a limited amount of growth over a 51 day period when sprayed on the foliage of newly germinated pecan seedlings grown under greenhouse conditions.

Agricultural Research Service scientists (16) have found that young greenhouse seedlings of black walnut and loblolly pine sprayed with 400 ppm GA₃ solution developed stems 40% taller and were twice the thickness of controls at the end of one growing season.

Kuskova (7) found that soaking of belladonna seeds, before sowing, with a 2% solution of gibberellin increases dry weight of the leaves and stems, increases the alkaloid content in the stems and leaves, and increases the alkaloid content of the entire plant while affecting only slightly the height of the plant. Supplementary spraying of belladonna plants with a 2% solution of GA₃ nearly doubled the height of the plant, did not affect the dry weight of the stems, reduced alkaloids in the leaves and roots, and increased the alkaloids in the stems.

Thomas et al (14) found that GA_3 applied in lanolin paste in a band on the last internode, foliage spray, and by a single drop to the apical bud of mango seedling stocks increased stem length and stem diameter of the plants over controls. Also, they found that each application of GA_3 , even when supplemental, was followed by immediate stem elongation within a month of treatment. Trees treated twice or three times had longer stems and larger stem diameters than trees treated once.

Nanda and Purohit (11) reported that enhanced extension growth in Salmalia malabarica caused by gibberellin treatment is probably a consequence of mobilization of stored food, and that it appears that gibberellin causes mobilization of starch into sugar which becomes readily available for extension growth of cells.

Ting and Lockhart (15), experimenting with applied gibberellic acid on pinto beans, found that applied GA_3 moves with carbohydrates within the plant.

Marth and Mitchell (8) increased the size of black walnut seedlings from an average of 1.5 feet (controls) to 8.5 feet in one season by three consecutive applications of gibberellic acid applied in lanolin bands at the base of terminal buds. GA_3 prevented summer dormancy from occurring. Continuous growth throughout the season caused the plants to produce the equivalent of several seasons growth.

Zych(22) found that gibberellic acid, applied as a spray, inhibited rooting of Purple Autumn raspberry. Inhibition was

greater for mature (basal cuttings) cuttings than for tips (terminal cuttings).

Mastalerz (10) reported that single applications of GA₃ applied as a spray increased the size and quality of Better Times roses. Repeated sprayings with the same concentration of GA₃ gave an increase in stem length but flowers were abnormal and of low quality.

Barker and Collins (1) found that GA₃ applied in lanolin or in water, as a fine spray, in concentrations as low as 20 ppm induces parthenocarpy in the low bush blueberry when applied to the flower immediately after anthesis. This concentration applied to parts of the plant other than the flowers did not induce parthenocarpy. None of the treatments at 20 ppm concentration caused an increase in stem growth.

MATERIALS AND METHODS

Stratified hardy northern pecan seed was planted in a sandy loam soil at the Ashland Horticulture Farm, Manhattan, Kansas, in April 1965. Seeds were planted one every six inches in three rows which were spaced four feet apart. Seedlings were weeded and irrigated as needed. Twenty-four plots each containing nine seedling pecan trees with ten unsprayed control seedlings separating sprayed plots were studied. The leaves of five seedlings in each of the twenty-four plots were sprayed June 20, 1966, with an aqueous solution of 5,000 ppm GA₃. Each sprayed tree in each treated plot was separated from the other treated

trees by an unsprayed tree.

Twelve previously untreated plots (60 trees) were sprayed July 20, as described above. Sixty trees sprayed June 20, received a second application of GA₃ at 5,000 ppm July 20.

Spray applications were made with a hand operated compressed air sprayer. Three drops of a surfacant, Tween 20, were added to each 135 milliliters of spray. The spray solution had a pH of 3.2. Plants were sprayed until run-off. Excessive spray drift was prevented by covering the seedling during spraying with a polyethelene bag and inserting the spray nozzle into the bag to make the application.

Fertilizer (20-10-5 NPK) was side-dressed at 350 pounds per acre August 20, 1966, to all seedlings. Fertilizer was applied uniformly in eight inch bands on each side of the row and then the plots were irrigated.

Twenty trees selected at random from groups treated each in June and in July and those sprayed twice were dug and prepared for sampling in early December as were 118 unsprayed control trees. Measurements recorded included stem and root length, collar and stem diameter four inches above the collar. All samples were cut into pieces approximately a half inch long and oven dried for ten days at 70 degrees Centigrade. Data collected following drying included dry weight of stems, roots and total plants. The ratio of dry weights of stem/root was calculated. Analysis of variance was calculated for each value obtained, and effects of treatments were compared by the Duncan Multiple Range

Test (4).

RESULTS

Table 1 shows response in growth of pecan seedlings to spray applications of gibberellic acid at 5,000 ppm. Comparisons can be made among trees sprayed only in June or July and those sprayed both months.

Seedlings sprayed in June had longer stems and greater dry weights of stems than unsprayed trees. However, dry weight of treated roots was less than for unsprayed seedlings. The ratio of stem dry weights to root weights was greater in treated than in control trees. Stem and collar diameters, total dry weights, and length of roots of treated seedlings were not significantly different from controls.

Seedlings sprayed with gibberellic acid in July had longer stems, greater stem diameters, heavier oven dry weights of stems than controls. However, average dry weights of roots and root length were less than those of controls. Ratio of dry weight of stems to roots was greater for sprayed trees than for controls. Average diameters of collars and total dry weights were not significantly different from the controls.

Table 11 shows effects of date of treatment and accumulative effects of two GA_3 sprays.

Trees receiving GA_3 sprays in both June and July had significantly longer stems, greater stem dry weights, and less root dry weights than trees sprayed only once in June or July. Also,

Table 1. Some effects on growth of year-old pecan seedling trees from aqueous sprays of GA₃ at 5,000 ppm in 1966. Treatment groups included trees sprayed in June or July and those sprayed both months.

Date treated	Mean measurements, inches				Mean weights, grams				Ratio dry wt. stem/ root
	Stem length	Root length	Collar diameter	Stem diam. 4 in. above collar	Stem dry wt.	Root dry wt.	Total dry wt.		
6/20/66	a.	17.930 **	18.952 ns	.388 ns	.333 ns	11.894 *	25.868 *	37.761 ns	.544 **
	b.	11.180	21.280	.441	.295	8.450	36.342	44.788	.238
7/20/66	a.	21.515 **	20.280 *	.437 ns	.388 **	14.927 **	15.245 **	30.172 ns	1.058 **
	b.	9.700	24.674	.415	.268	5.891	33.011	38.953	.182
6/20/66 and 7/20/66	a.	27.833 **	21.718 ns	.442 ns	.411 **	22.947 **	11.276 **	34.223 ns	1.983 **
	b.	11.180	21.280	.441	.295	8.450	36.342	44.788	.238

(a) treated seedlings
(b) control seedlings

* significant at 5%
** significant at 1%

ns non-significant at 5%

Table 11. Comparison of effects of spray applications of 5,000 ppm GA₃ on growth and development of year-old pecan seedlings.

Date treated	Mean measurements, inches				Mean weights, grams			
	Stem length	Root length	Collar diameter	Stem diam. 4 in. above collar	Stem dry wt.	Root dry wt.	Total dry wt.	Ratio dry wt. stem/root
6/20/66	17.930a	18.952a	.388a	.333a	11.894a	25.868a	37.761a	.544
7/20/66	21.515a	20.280a	.437a	.388ab	14.927a	15.245a	30.172a	1.058
6/20/66 and 7/20/66	27.833b	21.718a	.442a	.411b	22.947b	11.276b	34.223a	1.983

Values with the same letter have no significant difference
 Values with unlike or no letters differ significantly

trees sprayed twice had significantly larger stem diameters than June sprayed trees. Furthermore, twice treated trees had significantly greater stem-root ratios than seedlings sprayed only once in June or July.

The only significant difference between trees sprayed only in June and trees sprayed only in July was in stem-root ratio; ratio was larger for July sprayed trees.

DISCUSSION

Although seedlings sprayed in June only differed significantly from those sprayed in July only by just one factor, stem root ratio, other differences are large enough to indicate that time of treatment affects the response of pecan seedlings to gibberellic acid. Trees treated in July were 3.5 inches taller, roots 1.3 inches longer, and stem diameter .05 inches wider than comparative values for trees treated in June.

Observed differences in response by trees sprayed in June and trees sprayed in July are probably due to the amount of GA₃ that penetrated into the leaves of treated trees. Two of the factors that apparently affect the penetration of growth substances are temperature (20) and surface area treated (13). The average high temperature for five days before to five days after the June 20 treatment was just above 85 degrees Fahrenheit and the average low temperature for the same period was nearly 63 degrees Fahrenheit (17). Average high temperature five days before to five days after the July 20 treatment was nearly 94 de-

degrees F and the average low temperature for the same period was nearly 79 degrees Fahrenheit (18). According to Van Overbeek(20), natural permeation barriers of leaves have a higher permeability with increased temperature. Sargent (13) reports that it seems likely that a growth regulator will diffuse through surface layers at a net rate proportional to its concentration gradient across the layers. There was less leaf area to be covered on the June treated trees than on the July treated trees.

These same factors that influenced penetration of growth substances, temperature and surface area treated, probably account for the differences in response to GA₃ treatment between trees sprayed twice and those sprayed once in June or July.

Results of supplemental spraying observed in this study are similar to the findings of Nanda and Purohit (11) in their research involving supplemental GA₃ treatment of Samalia malbarica. They found that increasing the concentration of applied GA₃ caused increased growth of the stem with a corresponding increase in the dry stem-root ratio. Martin and Wiggins (9) found that increasing the soaking time of pecan seed treated with GA₃ from 12 to 192 hours reduced root dry weight and total dry weight of the plant. Xuskova (7) observed a similar response to GA₃ by belladonna plants. Supplemental spraying with GA₃ on belladonna resulted in doubled stem height and reduced root dry weight. However, stem dry weight was not affected.

Increased stem-root ratio from greater concentrations of applied gibberellic acid is probably the result of increased

stem growth at the expense of root development. Halevy et al(5) found that the effect of GA_3 on translocation of assimilates is to increase the mobilization of stored materials and to enhance distribution toward stem tips. Nanda and Purohit (11) report that gibberellic acid causes mobilization of starch into sugar which becomes readily available for extension growth of cells.

Pecan seedlings treated with 5,000 ppm gibberellic acid developed leaves that were smaller than controls and the leaves were pale and lacking the green color of leaves on controls. Trees sprayed in June had leaves that were a normal green color by July 20, 1966; although they were still smaller than leaves of control seedlings. Seedlings sprayed for the first time in July and those treated in June and again in July developed new small pale leaves on new stem growth. Fertilizer was applied August 20, 1966, and, within two weeks, leaves became a normal green. However, leaves remained small as did the leaves of the trees sprayed only in June. Results of fertilization are similar to those reported by ARS scientists (16) who reported that paleness in the green of developing leaves on plants treated with GA_3 was changed to a deeper green by the addition of fertilizer.

The growing points of trees treated with GA_3 in July only and the growing points of trees treated both in June and July were damaged by frost on October 16 (19). However, the rest of the stem did not appear to be damaged. Trees sprayed in June only and control seedlings were not injured other than leaf-kill by the frost.

If the normally extensive root system of pecan seedlings could be substantially reduced in overall development, lifting and transplanting of this species would be correspondingly easier; less labor would be required in lifting operations, packing, handling, and storage would require less space. Furthermore, it would be logical to assume that a smaller root system could result in lifting of seedlings with less injury to the taproot thereby resulting in more rapid early growth of transplanted seedlings than now occurs.

Root length was significantly reduced in seedlings sprayed with GA_3 in July only, and root length was less than controls, though not significantly so, in trees treated in June only. However, root length was slightly greater than controls for trees treated both in June and July. These findings indicate that additional research with gibberellic acid on pecan seedlings could discover the ideal combination of time of treatment, fertilizer, and concentration of applied GA_3 that will decrease seedling root size to a degree that will benefit growers. Also, other gibberellins (21) may hold promise in such research.

This study generally confirms the findings of Martin and Wiggins (9) with one exception. They found that gibberellic acid can cause only a limited amount of growth of pecan seedlings over a 51 day period of growth regardless of concentration used or number of spray applications. The observations reported in this study show that repeated application of GA_3 does cause increased growth of stems of year-old pecan seedlings. Further, they found

that increasing the soaking time of pecan seeds with GA_3 from 12 to 192 hours reduced the root dry weight and total dry weight of seedlings germinated. Increased dry stem-dry root ratio is in agreement with the findings reported in this paper.

The accomplishments of Marth and Mitchell (8) and other Agricultural Research Service scientists (16) in increasing the growth of black walnut seedlings with applications of gibberellic acid would indicate the possibility of inducing a greater growth in pecans than reported by Martin and Wiggins (9) or reported in this study. Additional research with the effects of gibberellic acid on pecan seedlings for stocks should attempt to determine the best time for GA_3 application, the best method of GA_3 application, and to find the GA_3 concentration that will yield the best results.

There are thirteen known gibberellins and three gibberellin-like substances that have been detected (21). Additional research to improve pecan stocks with gibberellic acid should also investigate the effects of other gibberellins on pecan seedling growth.

SUMMARY

Field application of 5,000 ppm gibberellic acid to year-old pecan seedlings significantly increased stem diameter on seedlings treated in July and those treated both in June and July. Effects of gibberellic acid on stem diameter were not as great on the seedlings sprayed in June. However, the diameter

increase was large enough in each case to improve the suitability of the seedlings for budding and grafting.

There was a difference in response to 5,000 ppm GA₃ spray between pecan seedlings treated in June and those treated in July. Although the only significant difference was in dry stem-dry root ratio, other differences were large enough to indicate that time of treatment is an important factor in response of pecan seedlings to gibberellic acid.

A second application of 5,000 ppm gibberellic acid in July to seedlings that had been previously treated in June gave significant increases over other treatments in stem length, dry stem-dry root ratio, and stem dry weight. Root dry weight was significantly less than root dry weight of seedlings sprayed only once in June or July. Stem diameter was significantly larger than the same measurement for trees treated only in June but not significantly different from stem diameters of trees treated only in July.

ACKNOWLEDGEMENTS

The author wishes to express appreciation to Dr. Ronald W. Campbell, Head of Department of Horticulture and Forestry, major advisor, for his help and counsel during the course of this study and in the preparation of this manuscript.

Thanks is also expressed to the other members of the supervisory committee for their suggestions and constructive criticism: Dr. Robert P. Ealy, Professor of Landscape Architecture; Dr. William J. Carpenter, Professor of Horticulture; and Dr. James K. Greig Jr., Associate Professor of Horticulture.

He also wishes to express his appreciation to Dr. Neil W. Miles, Assistant Professor of Horticulture, for his help in the collection and preparation of samples.

LITERATURE CITED

1. Barker, W.G. and W.B. Collins. 1965. Parthenocarpic Fruit set in the Lowbush blueberry. Proc. Amer. Soc. Hort. Sci. 87:229-233.
2. Bilan, M.V. and A.K. Kemp. 1960. Effect of gibberellin on height growth of one-year old seedlings of loblolly pine. Jour. Forestry 58:35-37.
3. Bilan, M.V. and A.K. Kemp. 1962. Effect of gibberellin on growth and development of loblolly pine seedlings. Jour. Forestry 60:391-394.
4. Duncan, David B. 1955. Multiple Range and Multiple F Tests. Biometrics. 11:1-42.
5. Halevy, A.H., S.P. Monselise and Z. Plaut. 1964. Effects of gibberellin on translocation and on dry matter and water content in several plant species. Physiol. Plantarum. 17: 49-62.
6. Kenworthy, Enoch Dawson. 1959. Some effects of gibberellic acid on fruit plants and seed. Master's thesis, Kansas State University, Manhattan, Kansas.
7. Kuskova, Z.R. 1965. The effect of gibberellin on belladonna. Soviet Plant Physiology. 12:548-553.
8. Marth, Paul C. and John W. Mitchell. 1961. Growth regulators, stimulants, and seeds. The Yearbook of Agriculture: 51-57.
9. Martin, L.W. and S.C. Wiggins. 1961. The effect of gibberellic acid on germination and seedling growth of pecans. Proc. Amer. Soc. Hort. Sci. 77:295-300.
10. Mastalerz, John W. 1965. The effect of gibberellic acid on the flowering shoots of Better Times roses. Proc. Amer. Soc. Hort. Sci. 87:525-530.
11. Nanda, K.K. and A.N. Purohit. 1965. The effect of gibberellin on mobilization of reserve food and its co-relation with extension growth. Planta. 66:131-125.
12. Nelson, T.C. 1957. Early responses of some southern tree species to gibberellic acid. Jour. Forestry. 55:518-520.
13. Sargent, J.A. 1965. The penetration of growth substances into leaves. Ann. Rev. Plant Physiol. 16:1-12.

14. Thomas, Paul, S. Krishnamurthi and V.N. Madhava Rao. 1963. Effect of gibberellic acid on nursery stocks of mango (*mangifera indica* linn.) and citrus spp. South Indian Horticultre. 11:18-30.
15. Ting, Yun Chin and James A. Lockhart. 1965. Translocation of applied gibberellin in bean seedlings. Am. Jour. of Botany. 52:828-833.
16. U.S.D.A. 1958. Gibberellic Acid An Unusual Plant-Regulating Chemical ARS 22-47.
17. U.S.D.C. 1966. Climatological Data. 80:No.6.
18. U.S.D.C. 1966. Climatological Data. 80:No.7.
19. U.S.D.C. 1966. Climatological Data. 80:No.10.
20. Van Overbeek, J. 1959. Auxins. The Bot. Rev. 25:271-350.
21. Van Overbeek, J. 1966. Plant harmones and regulators. Science. 152:721-731.
22. Zych, C.C. 1965. Rooting of leaf bud cuttings of *Rubus neglectus* inhibited by gibberellin. Proc. Amer. Soc. Hort. Sci. 87:238-239.

SOME EFFECTS OF GIBBERELIC ACID
ON YEAR-OLD PECAN SEEDLINGS

by

LOY WILLIAM SHREVE

B. S. F., West Virginia University, 1951

AN ABSTRACT OF A MASTER'S THESIS

submitted in partial fulfillment of the

requirements for the degree

MASTER OF SCIENCE

Department of Horticulture and Forestry

KANSAS STATE UNIVERSITY
Manhattan, Kansas

1967

The purpose of this study was to determine if field applications of gibberellic acid would increase pecan seedling growth to a size suitable for budding and grafting earlier than untreated pecan seedlings.

Stratified hardy northern pecan seed was planted in sandy loam soil at Ashland Horticulture Farm, Manhattan, Kansas, in April, 1965. Seeds were planted one every six inches in rows which were spaced four feet apart. Seedlings were weeded and irrigated as needed. Twenty-four plots were studied each containing nine seedling pecan trees with ten unsprayed control seedlings separating each plot. Five seedlings in each of the twenty-four plots were foliage sprayed June 20, 1966, with an aqueous solution of 5,000 ppm GA₃. Sprayed trees in each plot were separated from the other treated trees by an unsprayed guard tree.

Twelve previously untreated plots (sixty trees) were sprayed July 20, as described above. Sixty trees sprayed June 20, received a second application of GA₃ at 5,000 ppm July 20.

Spray applications were made with a hand operated compressed air sprayer. A few drops of a surfacant, Tween 20, were added to each spray treatment. The pH of the spray solution was 3.2. Plants were sprayed until run-off was observed.

Fertilizer (20-10-5 NPK) was side dressed at 350 pounds per acre, August 20, 1966, to all seedlings.

Twenty trees from each treated group as well as 118 unsprayed control trees were dug and prepared for sampling in early

December. Measurements recorded included stem and root length, collar and diameter of stem four inches above the collar. All samples were cut into pieces about $\frac{1}{2}$ inch long and oven-dried for ten days at 70 degrees Centigrade. Data collected following drying included dry weight of stems, roots, and total plant. Stem/root ratios were calculated. Analysis of variance was calculated for each value obtained, and effects of treatments were compared by the Duncan Multiple Range Test.

Seedlings sprayed in June had longer stems and greater dry weight of stems than unsprayed trees. However, dry weight of treated roots was less than for unsprayed seedlings. Stem/root ratios were greater in treated than in control trees.

Seedlings sprayed with GA₃ in July had longer stems, greater stem diameters, and heavier dry stem weights than controls. However, average dry weight of roots and root lengths were less than those of controls. Stem/root ratios were greater for treated trees than for controls.

Trees sprayed in both June and July had significantly longer stems, greater stem dry weight, less root dry weight, and a greater stem/root ratio than trees sprayed only once in June or July. Trees treated in both June and July had significantly larger stem diameters than trees treated only in June.

The only significant difference between trees sprayed in June and those sprayed in July was in stem/root ratios which were larger for July treated trees.

Differences in response to treatment were probably due to

temperature at the time of treatment and the size of the tree when GA₃ was applied; both are factors that may affect the amount of GA₃ utilized by the plant.