

STRUCTURAL COMPARATION OF CEREAL AND TUBER  
AMYLOPECTINS

by

BEATRIZ GÜTIERREZ P.

B.A. National University of Colombia, 1973.

---

A MASTER'S THESIS

submitted in partial fulfillment of the  
requirements for the degree

MASTER OF SCIENCE

Department of Grain Science and Industry

KANSAS STATE UNIVERSITY

Manhattan, Kansas

1985

Approved by



-----

Major Professor

LD  
7668  
.74  
1985  
G875  
c.2

111202 641023

i

TABLE OF CONTENTS

	page
INTRODUCTION.....	1
LITERATURE REVIEW.....	2
Amylytic enzymes.....	2
Amylose structure.....	4
Amylopectin structure.....	5
MATERIALS AND METHODS.....	10
STARCHES.....	10
Wheat.....	10
Pearl millet.....	10
Mandioc and Ocumo.....	11
Starch fractionation.....	12
Iodine affinity.....	13
Amylose content.....	13
Preparation of B-limit dextrin.....	14
Debranching with pullulanase.....	15
Debranching with isoamylase.....	16
Gel filtration.....	16
RESULTS AND DISCUSSION .....	18
Characterization of starches and their fractions.....	18
Characterizations of the isolated amylopectins.....	19
SUMMARY AND CONCLUSIONS.....	32
ACKNOWLEDGEMENTS.....	39
DEDICATIONS.....	40
BIBLIOGRAPHY.....	40

## LIST OF FIGURES

Figure	page
1. Bio-gel P-10. Elution profiles of millet amylopectin, pullulanase debranched millet amylopectin, isoamylase debranched millet amylopectin.....	21
2. Bio-gel P-10 Elution profiles of wheat amylopectin, pullulanase debranched wheat amylopectin, isoamylase debranched wheat amylopectin.....	22
3. Bio-gel P-10. Elution profiles of Venezuelan mandioc amylopectin, pullulanase debranched Venezuelan mandioc amylopectin, isoamylase debranched Venezuelan mandioc amylopectin.....	23
4. Bio-gel P-10 . Elution profiles of Colombian mandioc amylopectin, pullulanase debranched Colombian mandioc amylopectin, isoamylase debranched Colombian mandioc amylopectin.....	24
5. Bio-gel P-10 . Elution profiles of ocumo amylopectin, pullulanase debranched ocumo amylopectin, isoamylase debranched ocumo amylopectin.....	25

6. Percent B-amylolysis of amylopectin fractions from millet, wheat, Venezuelan mandioc, Colombian mandioc and ocumo..... 27
7. Bio-gel P-2 . Elution profiles of pullulanase debranched millet B-limit dextrin, wheat amylopectin B-limit dextrin..... 29
8. Bio-gel P-2. Elution profiles of pullulanase debranched Venezuelan mandioc B-limit dextrin, Colombian mandioc B-limit dextrin, ocumo B-limit dextrin..... 30

## LIST OF TABLES

Table		page
I	Proximate analysis of starch samples.....	34
II	Iodine affinity of starch samples and their fractions.....	35
III	Amylose content of starch samples and their fractions.....	36
IV	Characteristic of the isolated amylopectins.....	37
V	A:B chain ratio of isolated amylopectins.....	38
APPENDIX I	.....	47
APPENDIX II	.....	48
APPENDIX III	.....	49

## INTRODUCTION

Starch constitutes the major food reserve of higher plants. It constitutes up to 70% of the cereal kernel (22) and up to 24% of fresh tubers (38). Starch granules vary in size, shape molecular structure and properties depending upon the botanical source (38). Starch performs two main roles in food products, it is a dietary carbohydrate source (i.e nutritive) and it provides the characteristic viscosity, texture, mouth-feel and consistency of many food products, both cereal based and non-cereal based. Starch also is frequently used as a processing aid to facilitate the manufacture of food products. As a consequence of its importance, starch, its structural and functional properties has been the subject of a great deal of research over the last 60 years.

In order to understand the role of the starch, as it relates to the quality of the final food products, it is necessary to know the differences of the intrinsic structure of the molecular components of starch as they exist from different starch sources. This study was undertaken to compare the fine structure of amylopectins from millet (Pennisetum americanum) wheat (Triticum aestivum) mandioc (Manihot esculenta crantz) and ocumo (Xanthosoma sacittifolium).

## LITERATURE REVIEW

Starch is an organized polymer composed of repeating D-glucose units. In nature it exists as discrete collections of molecules or granules. Chemical and biochemical evidence indicate that two kinds of carbohydrate polymers are present in the starch granule, amylose and amylopectin. They exist in different proportions according to the source of starch. The amylopectin fraction predominates in tubers and most of cereal starches. Significant differences in molecular structure and physical properties exist between these two types of polymers. The application of enzymatic methods and chromatographic techniques to the structural analysis of amylaceous compounds has been useful in elucidating the molecular structure of these polymers (2,8,9,15,21).

## AMYLITIC ENZYMES.

The enzymes which attack and degrade starch can be divided into two main groups; debranching enzymes, those that specifically hydrolyze the alpha 1-6 interchain linkages of amylopectin, and the alpha 1-4 glucosidases, those that split the alpha 1-4 glycosidic linkages. The latter can be further subclassified into enzymes that make random or internal cleavages (endo acting enzymes) and those that act sequentially from chain ends (exo acting enzymes) (27).

Among the enzymes which attach starch ALPHA-AMYLASE is specific for alpha 1-4 glycosidic linkages.(26) Its action pattern is essentially random and directed to form primarily alpha- limit dextrans and some D-glucose and maltose .The former are branched oligosaccharides of dp 4 or more, containing alpha 1-6 D-glucosidic linkages (15,26).

B-AMYLASE (E.C.3.2.1.2.), on the other hand, is an exo-enzyme starting at the non-reducing end of the polymer, it splits alternate alpha 1-4 bonds in amylaceous polysaccharides yielding maltose. It can not split alpha 1-6 glycosidic linkages of amylopectin neither can it pass alpha 1-6 branch points.The A or outer chains of amylopectin are degraded by B-amylase to a predictable length of two or three glucose units depending on whether the chain contained and odd or even numbers of glucose residues before digestion (19,35).Thus,an odd number will result in two residues remaining while an even number in three residues remaining. When amylopectin is treated with B-amylase approximately 50% of the carbohydrates are converted into maltose (26,27).The remaining B-amylase resistant polymer is referred to as a B-limit dextrin.

Enzymes that specifically hydrolyse alpha 1-6 glycosidic bonds have great utility in the structural analysis of starch. PULLULANASE (E.C.3.2.1.41) first isolated in 1961 (35), acts on

amylopectin to affect nearly complete removal of its branches (1,28) by its action on alpha 1-6 bonds. The endo enzyme also debranches B-limit dextrans, releasing maltose and maltotriose from the original A chains and higher oligosaccharides (presumably from the B chains) (26). Pullulanase debranching actin on amylopectin, glycogen, and their B limit dextrans, followed by chromatographic fractionation of the resulting carbohydrate polymers has been used in the determination of the average of chain length ( $\bar{CL}$ ) of these polysaccharides and their constituent branches (2,8,9,10,19,24,25).

Harada et al (20) and Whelan et al (18) isolated and characterized a second enzyme, ISOAMYLASE, which can be used in structural determinations. The specificity of isoamylase is such that it completely debranches glycogen and amylopectin as well as the phosphorylase limit dextrans of these polysaccharides. (26) Its action on the B-limit dextrans of glycogen and amylopectin is incomplete (10,18). Specifically, isoamylase is able to cleave only alpha 1-6 linkages present at branch points in oligo and polysaccharides, and requires an outer or A chain of three or more glucose units before debranching can occur (27).

#### AMYLOSE STRUCTURE.

Enzymatic and physical studies have indicated that the

amylose component of starch is not a simple linear alpha 1-4 glucoside. (8,32) .Pure B-amylase preparations result in only approximately 70% conversion of amylose to maltose This has led to the conclusion that amylose consists of essentially linear polysaccharides and some what larger polysaccharides containing barriers to enzyme action (4,5) , probably some low levels of branching. Of course,an alternate explanation for the incomplete conversion of amylose to maltose by B-amylase is the presence of contaminating amylopectin or artefactual barriers to enzyme created during the polysaccharide solution procedure (26) .


#### AMYLOPECTIN STRUCTURE.

In the period before it became possible to explore the fine structure of amylopectin enzymatically, several structures for amylopectin were proposed that equally satisfied the known facts about the molecule. Early chemical methods showed that, although glucose units in amylopectin were linked primarily through alpha 1-4 glycosic bonds branching occurred through alpha 1-6 glycosidic linkages. his fact, the presence of one alpha 1-6 linkage in about twenty-four glucose units results in an infinitely large number of possible arrangements.

Peat and coworkers (38) divided the carbohydrate chains in the amylopectin molecule into three types,A,B,and C chains.

A-chains are those being no substituent chains or branch


points. Thus, they are linked to the rest of molecule only through their reducing ends.



- terminal non reducing group
- reducing group



B-chains are those that are linked to the molecule through their reducing ends but, in addition are branched at the C-6 position in one or more of their B-glucopyranosyl residues. The C-chain is the only carbohydrate chain in the molecule having a free reducing group.

The structures proposed by Haworth et al (38), Standinger et al (35) and Meyer (35) differ primarily in the proportions of A and B chains. In the Haworth structure, in which the chains carry only one substituent chain, there is only a single



LAMINATE STRUCTURE:  
of Harword et al.

HERRINGBONE MODEL:  
of Staudinger et al.



RANDOMLY BRANCHED MODEL:  
of Meyer et al.

A chain and a single C chain, the rest of the molecule is composed of B chains. Standinger's structure, on the other hand, has all its chains with the exception of the C chain, unsubstituted. A chains branch out exclusively from the C chain with the result that there are no B-chains at all. Meyer's model has a randomly branched structure with equal numbers of A and B chains.

To distinguish between these three structures, enzymatic digestion followed by gel filtration has been used to directly measure the numbers of chains of each type following debranching, the average chain length ( $Cl^*$ ) of the released polymers can be determined (19,26). These values vary according to the amylopectin source. A  $Cl$  range of 17 to 20 has been reported for wheat amylopectin (24) and values of 25 for mandioc amylopectin (21).

An important parameter used to characterize the degree of multiple branching in the amylopectin molecule is the ratio of A to B chains (A:B ratio). By using isoamylase or pullulanase on B-limit dextrin that ratio can be calculated (25). Incubation with isoamylase alone results in incomplete debranching of amylopectin as a consequence of the inability of this enzyme to hydrolyse A chains that are two glucose units or less. This constitutes roughly half of the total numbers of chains. Amylopectin typically contains approximately twice as many A chains as B chains. A:B chain ratios are markedly higher in

amylopectin (1.5-2.6:1) than they are in glycogen (0.6-1.2:1) (27). Lii et al. (24) reported A:B chain ratios for different types of wheat amylopectin ranging from 1.21 to 2.11.

More recently gel-filtration chromatography of the low molecular weight debranched products of amylopectin, used to obtain a chain size profile has shown the existence of a bimodal distribution of carbohydrate polymers in amylopectin (17,21,24,27,29,33). This leads to the conclusion that amylopectin structure can be correctly described by the revised Meyer model proposed by Gunja-Smith et al. and further by the cluster models of French 1972 (15) and Robin et al 1974 (33). Whether the bimodal distribution results from the same polysaccharide molecules or not has been studied by different investigators (2,9,26). Marshall et al (27) suggested that the chain populations come from two different populations of polysaccharides. Atwell et al (2), by studying the B- limit dextrans of wheat amylopectin found that the overall amylopectin population was uniform and concluded that the bimodal distribution of chain sizes came from two chain populations or classes that existed within the same macromolecule.

## MATERIALS AND METHODS

## A.- MATERIALS.

## 1.-STARCHES.

## WHEAT:

Hard wheat flour was obtained from the pilot scale mill at Kansas State University. Wheat starch was isolated by the dough kneading procedure (39). Dough balls were kneaded under running water over 20xx bolting cloth and the aqueous slurry obtained centrifugated at 2000 xg for 30 minutes. The tailings layer was removed by scraping with spatula and the prime starch resuspended in water and centrifuged until no tailing layers was obtained. The recovered prime starch was lyophilized.

## PEARL MILLET:

To obtain millet starch, pearl millet grain was soaked at 4° C for 24 hours in distilled water containing 0.01% sodium azide. It was then washed several times with distilled water. The wet millet grain was ground in a Waring blender for 3 minutes, and screened through a 112 xx bolting cloth. The material retained on the screen was ground again for one minute and passed again through the bolting cloth. The slurry

obtained was allowed to settle and the supernatant was removed. the sediment was reslurried and passed through a 20 xx screen .The aqueous slurry obtained was centrifuged at 2000xg for 30 minutes and the tailings layer was removed by scraping it with a spatula.The resulting prime starch was reslurried in water and centrifuged until no tailing layer was obtained.The recovered prime starch was lyophilized.

#### MANDIIOC AND OCUMO :

Starches from mandioc (Manihot esculenta crantz) and ocumo (Xanthosoma-sacittifolium) were the generous gift of Dr. E.Schultz of the Central University of Venezuela, Maracay,Venezuela. These tuber starches were obtained by wet milling and dried in a forced air oven at 45 °C. In addition, mandioc starch obtained from Colombia,was produced by my mother, Rosa Gutierrez. It was made by a household procedure. Fresh mandioc was peeled and scraped. It was suspended in water and screened through a bolting cloth and the aqueous slurry was allowed settle. After decanting the supernatant the tailings layer was removed by scraping. The procedure was repeated several times until no more tailings layer was present.The prime starch was dried in the open air for three days.

## B.- METHODS.

## STARCH FRACTIONATION:

Prime starches were fractionated into amylose and amylopectin fractions by the extraction- sedimentation procedure of Montgomery (30). Two liters of 2% (w/v) starch in phosphate buffer, (pH 6.0-6.3) was heated at 95-98 °C for 11 minutes before autoclaving (20 psi) for two hours. The solution was saturated with n- pentanol (500 ml) and allowed to cool overnight with gentle stirring. The amylose - pentanol complex was collected by centrifugation (3000xg for 20 minutes). "Pure" amylose was obtained by adding 1.5 l. of hot water that had been saturated with n- butanol (100 mls) to the amylose and stirring overnight. The precipitated amylose complex was recovered by centrifugation (3000 xg for 20 minutes). The above procedure was repeated three times for mandioc and ocumo samples and five times for wheat and millet samples. Recovered purified amylose was mixed with fresh n- pentanol in a Waring blender and filtered thru a coarse porosity, scintered glass funnel before being lyophilized.

Amylopectin was recovered from the original supernatant by adding an equal volume of methanol to the aqueous solution. The amylopectin was allowed to precipitate overnight under refrigeration. The

amylopectin, then, was mixed with methanol in a waring blendor (15 sec) and filtered thru a coarse porosity scintered glass funnel before being lypholyzed.

#### IODINE AFFINITY.

The iodine affinity of starches, and its fractions amylose, and amylopectin were determined by potentiometric titration (34). Starches, amylose, and amylopectin fractions from millet and wheat were defatted by soxhlet extraction for 24 hours with 95% ethanol. The samples were then lyophilized and pulverized before the determination. A Corning pH-meter (Model 140) with a calomel electrode as reference and a bright platinum electrode was used for analysis. Two minutes was allowed for equilibration between the addition of iodine solution and obtaining a reading. Since deionized water was found to be essential for reproducible results, it was used for all determinations. Results are report as percent og iodine bound.

#### AMYLOSE CONTENT:

The amylose content of starches, amylose, and amylopectin fractions was determined by the colorimetric procedure of Bienvenido (7) using pure amylose and pure amylopectin from potatoe as standards. Spectrophotometric readings at 620 mm were takeng with a Beckman spectronic 21

(250 Harbor Blvd., Fullerton, C.A. US. 92634). All determinations were run at least four times. Results are report as percent of amylose.

#### PREPARATION OF B-LIMIT DEXTRIN.

Amylopectin (1g) was dissolved in 0.1 M NaOH (30 ml) in a 50 ml volumetric flask. The pH was adjusted to approximately 6 with 1M HCl. Five ml of 0.2 M acetate buffer (pH 4.8) was added and the sample diluted to volume. An aliquot (1 ml) diluted 1:100 was used to determine total carbohydrate in the samples by the phenol - sulfuric acid procedure (14). The determination was run 5 times for each sample. B-amylase (4500 units, type 1-B lot 122F-8120, Sigma Chemical Co, St Louis, Mo.) was added to the flask and the total solution dialyzed against repeated changes of 0.02M acetate buffer (pH 4.8) to remove carbohydrate, generated by the enzyme action. Several drops of toluene were added to the dialysis bag to inhibit microbial growth. After 24 hours an additional 1000 units of B- amylase were added and the dialysis continued until no carbohydrate was detected in the dialyzate. The contents of the dialysis bag were concentrated, transferred to a 50 ml volumetric flask and diluted to volume. An aliquot (1 ml) diluted 1:100 was used to determine total carbohydrate (see above) and the results compared with the original

solution to obtain the B-amylolysis limit. The remainder of the solution was heated for 15 minutes in a boiling water bath, filtered through Whatman # 42 filter paper and lyophilised.(24,25,36).

#### DEBRANCHING WITH PULLULANSE:

Pullulanse from Eterobacter aerogenes (lot 122F-8120 and 14F 02031) was obtained from Sigma Chemical Co.St Louis,MO. as a suspension in 3.2 M  $(\text{NH}_4)_2\text{SO}_4$ .It was used as received unless the Nelson's colorimetric copper procedure was to be subsequently performed (31).Since ammonium sulfate interferes with the reduction of copper in the assay it was removed by dialysis against water before any analysis of reducing power was done. The enzyme did not show any alpha (1-4) glucosidase activity ( see appendix II.)

Amylopectin (100 mg) or B-limit dextrin (50 mg) was dissolved in 0.1M NAOH (5 mls) and neutralized with 1M acetic acid before dilution to 10 mls with acetate buffer (0.1 M,pH 5.5). Total debranching was obtained by adding 16 units of pullulanase to the solution and incubating at 37°C for 36 hours.The digests were subsequently heated in boiling water for 20 minutes to inactivate the enzyme.Insoluble material was removal by filtration.(2,9,24).

## DEBRANCHING WITH ISOAMYLASE:

Isoamylase from Pseudomonas amyloclavata (lot 73F-03581B) was obtained from Sigma Chemical Co. St Louis Mo. Since it did not show contaminating alpha 1-4 glucosidase activity (see appendix III) it was used as received.

Amylopectin (50 mgs) was dissolved in 0.1 M NaOH (5mls) and neutralized with 1M acetic acid before dilution to 10 mls with acetate buffer (0.1 M, pH 5.5). Total debranching was performed using 0.27 units of isoamylase and incubating at 37° C for 24 hours. The digest was heated in boiling water for 5 minutes to inactivate the enzyme and insoluble material was removed by filtration (24).

## GEL FILTRATION :

Solutions containing approximately 5 mgs of debranched amylopectin were chromatographed on Pharmacia K 26/70 columns (2.5x75 cms) of Bio-gel P-10 or Bio-gel P-2. The columns were operated in the upflow mode. Carbohydrate was eluted with water containing 0.02% NaN<sub>3</sub> (2,9,24). The flow rate was 10 to 15 ml per hour. Fractions of approximately 4 mls were automatically collected with a Golden Retriever, Fraction collector, model 1200 pup

(Instrumentation Specialties Company, Inc. Lincoln, Nebr. 68504, U.S.A.) using the drop counting mode. Carbohydrate in the effluent fractions was quantitated by assaying an aliquot of each fraction using the phenol-sulfuric acid procedure (14). Each column was calibrated using 80 mgrs of pullulanase debranched waxy corn starch. The DP of the resulting fractions was obtained by calculating the total amount of carbohydrate in each fraction over the amount of reducing power in that fraction. Occasionally it was necessary to combine three or more fractions and concentrate to obtain sufficient material for the determination.

## RESULTS AND DISCUSSION

## CHARACTERIZATION OF STARCHES AND THEIR FRACTIONS:

The proximate analyses of starches from millet (Pennisetum americanum), wheat (Triticum aestivum), Venezuelan mandioc and Colombian mandioc (Manihot esculenta crantz) and ocumo (Xanthosoma sacittifolium), are shown in table I. Their isolated amylose and amylopectin fractions are shown in the table II and III respectively. As measured by potentiometric titration (34) the iodine affinity values (table II) for whole starches, ranged from 3.01 for Venezuelan mandioc starch to 4.89 for millet starch. These values agree well with those reported for wheat (24) millet (6) Mandioc (37). The iodine affinity of ocumo starch has not been reported previously. The value obtained for this starch (4.06) is higher than those previously reported for tuber starches (37). The amylose content for these starches is reported in table II. Values ranged from 15% for Colombian mandioc starch to 23% for wheat starch. These values agree well with those reported for wheat (37) millet (3,6) and mandioc (11). The amylose content of ocumo starch has not been reported. The results here (22%) suggest that ocumo starch contains a greater amount of amylose than do other tuber starches.

The isolated amylose, obtained by Montgomery's procedure, (30)

had iodine affinities ranging from 17.91% for Colombian mandioc to 83% for millet. Iodine affinity values for mandioc amylose samples agree well with those reported before (11). Defatted wheat and millet amyloses, on the other hand, showed iodine affinities some what lower than those reported previously (2,6,24,26,), although they were recrystallized five times. Similar results were obtained by measuring amylose content in these starch samples samples by a second method (7) (tableIII). The tuber amylose samples (mandioc and ocumo) were purer (84-90%) than the amylose samples isolate from cereals (wheat and millet) (75%-79%). This results showed that probably the structure of these cereal amylose fractions are different from that of tuber sample used as standard. These structural differences may have resulted in the purity of these cereal amylopectin fractions to be underestimated.

#### CHARACTERIZATION OF THE ISOLATED AMYLOPECTINS.

The isolated amylopectins from all five sources were quite pure as judged by their iodine affinity values and amylose contents (tables II and III), since the values agreed well with those reported by previous workers (2,3,4,6,11,24,26,). The samples were pure enough to use for subsequent debranching and chromatographic studies.

Incubation of amylopectin samples with pullulanase

(E.C.3.2.1.41) or isoamylase (E.C. 3.2.68) followed by gel filtration, on a Bio-gel P-10 column, yielded a bimodal distribution of carbohydrate chains from all the samples ( Fig 1-5). Elution profiles of all the amylopectin samples appeared to be similar with slight differences in the DP of their carbohydrate populations (Table IV). The leading peak, with largest Mw material, showed several differences between samples. It contained material with an average degree of polymerization (DP) of 50-58 for debranched wheat and ocumo amylopectins. The DP of the material in this fraction was larger than 58 for debranched Venezuelan and Colombian mandioc amylopectins as well as for debranched millet amylopectin. The second, later, elution peak contained material with an average degree of polymerization  $\overline{DP}$  of 10-16 for debranched wheat and ocumo amylopectins (table IV). The DP of the same peak for debranched amylopectin from millet and mandioc was larger (18-22). These results agree with previous studies on mandioc (21) wheat (2,24). The presence of these two carbohydrate populations leads to the conclusion that the amylopectin fractions for the samples studied here can be best described by the symmetric model proposed by Gunja-Smith et al (31) as well as the cluster model proposed by French (29).

The third, minor, peak observed in the chromatographs at the void volume can be explained as the presence of amylose contamination in the amylopectin (2,9,24). Incubation with the

Fig. 1.- Bio- gel P-10 .Elution profiles of millet amylopectin (——). Pullulanase debranched millet amylopectin (- - -) . Isoamylase debranched millet amylopectin (-·-·-·).





Fig.2.- Bio-gel P-10. Elution profile of wheat amylopectin (—) Pullulanase debranched wheat amylopectin (----) Isoamylase debranched wheat amylopectin (-.-.-).








Fig. 4.- Bio-gel P-10 . Elution profiles of  
Colombian mandioc amylopectin(——).  
Pullulanase debranched Colombian mandioc  
amylopectin ( - - - ) Isoamylase debranched  
Colombia mandioc amylopectin (-.-.-)





Fig. 5.- Bio-gel P-10 . Elution profiles of ocumo amylopectin (——). Pullulanase debranched ocumo amylopectin (----). isoamylase debranched ocumo amylopectin (-.-.-).




enzymes was done until no changes in that peak as detected.

The average of the chain length of the debranched amylopectins, obtained from the total amount of the polysaccharide debranched by pullulanase divided by the amount of reducing power liberated during debranching, (19) (table IV) revealed that Venezuelan mandioc amylopectin has the largest  $\bar{Cl}$ , 24, while ocumo amylopectin has the smallest  $\bar{Cl}$ , 16. The average chain length of wheat and mandioc amylopectins are comparable to those reported by previous workers (21,23,24).  $Cl$  of millet and ocumo amylopectins have not been previously reported. Our data suggests that the average chain length for millet amylopectin ( $Cl$  17) was similar to that obtained for wheat amylopectin. The  $\bar{Cl}$  of ocumo, (16), on the other hand, was lower than that of mandioc amylopectins, (24-22). Thus ocumo amylopectin has both more and shorter chains than does mandioc amylopectin (25).

Incubation of amylopectin samples with B- amylase produced B- amylolysis values ranging from 54% for Venezuelan and Colombian mandioc to 61% for millet (Fig.6) . B-amylolysis limits for wheat and millet amylopectins are higher than that for Venezuelan and Colombian mandioc and ocumo amylopectins. This may well reflect structural differences in the molecule. Thus, wheat and millet amylopectin molecules may be more branched than are mandioc and ocumo amylopectins .

After removal of maltose and by dialysis the amylopectin

Fig. 6.- Per cent B- amylolysis of amylopectins  
from: Millet ( ≡ ). Wheat ( / / / ).  
Venezuelan mandioc ( / / / ). Colombian  
mandioc (    ). Ocumo ( / / / ).



B-limit dextrans were debranched by pullulanase, chromatographed on Bio-gel P-2. Their elution profiles showed a similar distribution of carbohydrates in all the samples with only slight differences in average chain lengths. Three main populations can be distinguished (fig 7-8). The highest Mw material consisted of long linear chains of  $\overline{DP}$  37-50. The intermediate Mw material contained linear chains with  $\overline{DP}$  of 5-37. Those populations correspond to the B-chains. The smallest Mw population contained maltose and maltotriose as the result of the release of fragments of debranched A chains. These patterns corroborated the profiles obtained from pullulanase debranched starch-limit dextrin of waxy maize (23). The A:B chain ratio can be calculated by measuring the amount of reducing groups liberated from A chains after debranching (8) as a fraction of the reducing groups liberated from B chains during debranching (the rest of population). When this was done (table V) it shown that millet amylopectin has the largest A:B chain ratio ( 1.79 ) ( table V) while Venezuelan mandioc amylopectin, the shortest (1.42). Because A chains are unbranched and B chains are those which have another chain attached, the larger A:B ratio for millet amylopectin indicates that its amylopectin has a greater extent of multiple branching than the do other samples. This is particularly true relative to the Venezuela mandioc amylopectin. Colombian mandioc amylopectin apparently has is less multiple branched than are







Fig. 8.- Bio-gel P-2. Elution profile of pullulanase debranchin of Venezuelan mandioc B-limit dextrin (——) Colombian mandioc B- limit dextrin amylopectin (----) Ocumo B- limit dextrin (-.-.).



wheat or ocumo amylopectins.

For the amylopectin fractions studied the average chain length ; the B- amylolysis limits ;and the A:B chain ratio were internally consistent and supported these conclusion . Millet, for example , has the lowest average chain length (17) ,the highest B- amylolysis limit (61%) and thehighest A:B chain ratio (1.79) as expected for a more highly branched molecule. Manioc, on the other hand, has the highest average chain length (22-24), the lowest B- amylolysis limit (54%), and the lowest A:B chain ratio (1.42) again consistent and supporting a less highly branched molecule. These results led to the conclusion that millet, wheat, and ocumo amylopectin fractions possess more and shorter branches than do mandioc amylopectin fractions.

Thus, the fine structure of these amylopectins apparently have definite but minor diferences in their structure.

## SUMMARY AND CONCLUSIONS

Starch isolated from millet ( Pennisetum americanum) wheat (Triticum aestivium) Venezuelan and Colombian mandioc (Manihot esculenta crantz) and ocumo (Xanthosoma sacittifolium) were fractionated into their amylose and amylopectin components. The fine structures of the isolated amylopectins and its B-Limit dextrans were investigated using hydrolytic enzymes in conjunction with gel-filtration chromatography . The isolated amylopectin fractions treated with B-amylase appear to have B-amylolysis limits ranging from 54% for Venezuelan and Colombian mandioc amylopectin fractions to 61% for millet amylopectin fraction.

When the amylopectin fractions were debranched with pullulanase or isoamylase, the debranched fractions had unit chain profiles very similar to each other with two chains populations DP of approximately 10-16 and 50-58 respectively in wheat and ocumo amylopectins and DPs of 18-22 and > 58 respectively in millet, and Venezuelan and Colombian mandioc amylopectins.

The average chain length of the amylopectin ranged from 16 for ocumo amylopectin to 24 for Venezuelan mandioc amylopectin. Ocumo amylopectin appears to be composed of more and shorter chains than for other tubers .

The ratio of A:B chains in each amylopectin fraction, determined enzymatically ranged from 1.42 for Venezuelan mandioc amylopectin to 1.78 for millet amylopectin.

The data obtained for B-amylolysis, Cl\* debranched amylopectin and A:B ratio chain length were internally consistent and leads to the conclusion that millet, wheat, and ocumo amylopectin fractions has more and shorter branches that dose mandioc amylopectin fractin .

In conclusion the amylopectins investigated here appear to have definite, but minor differences in structure.

TABLE I

Proximate analysis of starch samples.

Starch-Samples	% Moisture	% Protein (Nx5.7) (dry basis)	%.Ash (dry basis)
Millet	5.0	0.68	0.15
Wheat	7.4	0.43	0.17
Venezuelan mandioc	11.6	0.17	0.18
Colombian mandioc	14.7	0.35	0.10
Ocumo	9.5	0.22	0.08

TABLE II

Iodine affinity of starch samples and their fractions.

Source	% of Iodine bound		
	Starches	Amyloses	Amylopectins
Millet	4.89 $\pm$ 0.23	18.83 $\pm$ 0.5	0.93 $\pm$ 0.03
Wheat	4.56 $\pm$ 0.35	17.95 $\pm$ 0.32	1.08 $\pm$ 0.09
Venezuelan mandioc	3.01 $\pm$ 0.31	18.86 $\pm$ 0.30	0.75 $\pm$ 0.17
Colombian mandioc	3.08 $\pm$ 0.44	17.91 $\pm$ 0.15	0.35 $\pm$ 0.05
Ocumo	4.06 $\pm$ 0.20	18.06 $\pm$ 0.14	1.35 $\pm$ 0.13

TABLE III

Amylose content of starch samples and their fractions.

Source	amylose content %		
	Starches	Amyloses	Amylopectins
Millet	22 $\pm$ 0.69	79 $\pm$ 0.95	2.0 $\pm$ 0.4
Wheat	23 $\pm$ 1.2	75 $\pm$ 1.41	1.7 $\pm$ 0.2
Venezuelan mandioc	16 $\pm$ 1.7	84 $\pm$ 1.82	6.0 $\pm$ 0.6
Colombian manioc	15 $\pm$ 1.8	85 $\pm$ 0.96	4.0 $\pm$ 0.2
Ocumo	22 $\pm$ 1.1	90 $\pm$ 0.5	6.0 $\pm$ 0.4

TABLE IV

Characteristic of the isolated amylopectins.

Source	$\overline{CL}^*$	$\overline{DP}^{**}$	
		I	II
Millet	17 $\pm$ 1.5	>58	18-22
Wheat	17 $\pm$ 1.5	50-58	10-16
Venezuelan mandioc	24 $\pm$ 1.8	>58	18-22
Colombian mandioc	22 $\pm$ 2.0	>58	18-22
Ocumo	16 $\pm$ 1.5	50-58	10-16

\* Average chain length of pullulanase debranched amylopectins.

\*\* Average of degree of polymerization of chain population resulting from the chromatography of pullulanase debranched amylopectins.

TABLE V

A:B chain ratio of isolated amylopectins.

Source	A:B ratio
Millet	1.79 <u>±</u> 0.08
Wheat	1.54 <u>±</u> 0.06
Venezuelan mandioc	1.42 <u>±</u> 0.05
Colombian mandioc	1.50 <u>±</u> 0.06
Ocumo	1.64 <u>±</u> 0.05

## ACKNOWLEDGEMENTS

The author express her gratitude to; Dr. J. Faubion, and Dr. R.C. Hoseney for their guidance and patience during the course of her work.

Also, she wants extended her thanks to; Dr. P. Nordin for serving on her graduate committee, Dr. D. Fung who made it possible for her to achieve a goal, and the entire staff of the Department of Grain Science and Industry who in one way or other have made possible the development of this work.

## DEDICATION

To my family especially my mother, Rosa, and my son, Edgar Alejandro, I dedicate my work and my life for. Without the moral support and encouragement of them, who I love, accomplishment of my goals would not been possible.

## BIBLIOGRAPHY

1. Abubllah, M. Catley, B. J . Lee, E.Y. (1966). The mechanism of carbohydrase action. XI. Pullulanase, an enzyme specific for the hydrolysis of alpha 1-6 bonds in amylaceous oligo - and polysaccharides. Cereal Chem. 43, 111-118.
2. Atwell,W.A. (1980). Debranching of Wheat Amylopectin. Cereal. Chem. 57, (1), 12-16.
3. Badi, S.M. Hoseney, R.C. Finney, P.L.(1976).Pearl millet II. Partial characterization of starch and use of millet flour in breadmaking. Cereal Chem.53 (5) 718-724.
4. Banks, W. Greenwood, C.T. (1966). The fine structure of amylose. The action of pullulanase as evidence of branching. Arch. Biochem. Biophys. 117, 674-675.
5. Banks, W.Greenwood, C.T. (1975). The fine structure of amylose in: Starch and its components. ed. by John wiley and Sons,Inc,N.Y. pg 15.
6. Belia,A. (1980). Characterization of starch and alpha-amylase from pearl millet. Master thesis.Kansas State University. Manhattan,Ks.

7. Bienvenido, J. (1974) . Manual of cereal chemistry procedures. ed. by International Rice Research institute. Los Banos,Laguna.Philippines.
8. Biliaderis,C.G. Grant, D.R. and Vose, J.R. (1979). Molecular weight distribution of legume starches by gel chromatography Cereal Chem.56 (5) 475-480.
9. Biliaderis, C.G. Grant, D.R. and Vose, J.R. (1981) Structural characterization of legume starches I.Studies on amylose, amylopectin and B-limit dextrin. Cereal Chem. 58 (6) 496-502.
10. Casola, L. Brumby, P.E. and Massey, V.(1966). A method for investigation of the fine structure of amylopectin. J.Biol.Chem.241 (4977) 1028-1029.
11. Ciacco, F.C. (1977). Characterization of starch from various tubers and their use in bread -baking .Cereal Chem.54 (5) 1096-1107.
12. Ciacco, C. F. D'Appolonia, B.L. (1978).Baking studies with cassava and yam II. Rheological and baking studies of tuber-wheat flour blends. Cereal Chem. 55 (4) 423-435.
13. Ciacco,C.F. D'Appolo, B.L.(8).Biochemical composition of cassava and yam flour.Cereal Chem.55 (3) 402-411.
14. Dubois, M. Gilles, K. Hamilton, J.K. Rebers, P.A.and Smith,

- F. (1956). Colorimetric method for determination of sugar and related substances. *Am.Chem.* 28 (3) 350-560.
15. French, D. (1972). Fine structure of starch and its relationship to the organization of the granule. *J. Japan. Soc. Starch. Sci.* 19, 8-12.
16. Greenwood, C. T. (1964). Viscosity - molecular weight relation. in: *Methods in carbohydrate chemistry* .ed. by Whistler, R.L. Ac.Press, N.Y. Vol 4, pg 179.
17. Gunja-Smith, Z. Marshall, J.J. Mercier, C. Smith, E. and Whelan, W. (1970). A Revision of the Meyer-Bernfeld model of amylopectin and glycogen. *FEBS. letters.* 12, (2) 101-104.
18. Gunja-Smith, Z. Marshall, J.J. Smith, E.E. and Whelan, W.J. (1970). A glycogen -debranching enzyme from cytophaga. *FEBS. Letters.* 12, (2) 96-100.
19. Gunja-Smith, Z. Marshall, J.J. and Smith, E.E. (1971). Enzymatic determination of the unit chain length of glycogen and related polysaccharides. *FEBS. letters.* 13, (5) 309-311.
20. Harada, T. Misaki, A. Akai, H. Yokoboyashi, K. and Kaname, S. (1972). Characterization of pseudomonas isoamylase by its actions on amylopectin and glycogen

- comparison with aerobacter pullulanase. Bichem. Biophys. Acta. 286, 497-498.
21. Hood, L.F. and Mercier, C. (1978). Molecular structure of unmodified and chemical modified mandioc starch. Carbohyd. Res. 61, 53-66.
  22. Kent, N.L. (1983). Technology of cereals .Ed. by Pergamon Press.
  23. Lii , E.Y.C. and Whelan, W.J. (1966). Enzymic methods for the micro-determination of glycogen and amylopectin; and their units chain lengths. Arch. Biochem. Biophys. 116, 162-167.
  24. Lii , C.Y. and Liweback, D.R. (1973). Characterization and comparison of cereal starches. Cereals. Chem. 54 (1) 138-149.
  25. Marshall, J.J. and Whelan, W.J. (1974). Multiple branching in glycogen and amylopectin .Arch. Biochem. Biophys. 161, 234-238.
  26. Marshall, J.J.(1974). Application of enzymatic methods to The structural analysis of polysaccharides. Adv. Carbohydr. Chem. Biochem. (30) 257-370.
  27. Marshall, J.J.(1975). Starch\_degrading enzymes,old and new. Die Staerke. 27, (11) 377-383.

28. Mercier, C. Frantz, B.M. Whelan, W.J. (1972). An improved purification of cell bound pullulanase from aerobacter aerogenes. Eur. J. Biochem. 26, 1-9.
29. Mercier, C. and Whelan, W.J. (1970). The Fine structure of glycogen from type IV. Glycogen-storage disease. Eur.J. Biochem. (16) 579- 583
30. Montgomery, E. Senti, F.R. (1958).Separation of amylose from amylopectin starch by an extraction sedimentation procedure. J .of Polymer Sci. (18) 1-9.
31. Norton, Nelson. (1944). A photometric Adaptation of smogy method for the determination of glucose. J. Bio Chem. 153:375-380.
32. Peat, S.Whelan, W.J. and Pirts,J. (1949). The amyolytic enzymes of soya bean. Natura. 164, 499-500.
33. Robin, J.D. Mercier, C. Charbonniere. R. and Guilbot A. Lintnerized starches. Gel filtration and enzymatic studies of insoluble residues from prolonged acid treatment of potatoe starch. Cereal. Chem.(51), May-jun 389-405.
34. Schoch, T. (1964). Iodimetric determination of amylose . In: Methods in carbohyadrates. chem.Vol 4, 157-175.
35. Whelan, W.J. (1971). Enzymatic explorations of the

structure of Starch and Glycogen. Biochem. J. 122,609-622.

36. Whelan, W.J. (1964). Hydrolysis with B-amylase and preparation of the B- amylase limit dextrin of amylopectin. In: Methods in carbohydrates chem. Vol 4, (57) 261-267.
37. Whistler, R.L. (1965). Starch:Chemistry and technology . Ed. Academic Press N.Y. and London. Vol 1.
38. Whistler, R.L. (1984). Starch:Chemistry and technology.Ed. by Academia Presss,Inc, N.Y.
- 39.- Wolff M.S. (1964). Wheat starch isolation in: Methods carbohydres chem. Vol4, (2) 6-9.

## APPENDIX I

B-amylolysis of each amylopectin fraction was determined by comparison of the total amount of carbohydrate present in the amylopectin solution before and after digestion with B-amylase .

$$\frac{\text{Total carbohydrate before} - \text{total carbohydrate after}}{\text{total carbohydrate before}}$$

Average of total carbohydrate determination before and after digestion with B-amylase was.

<u>Samples</u>	<u>Total carbohydrate (gluc.mg/ml).</u>	
	Before digestion	After digestion
	$\bar{X}_b$	$\bar{X}_a$
Millet	4.98 $\pm$ 0.5	1.94 $\pm$ 0.05
Wheat	4.75 $\pm$ 0.4	1.97 $\pm$ 0.03
Venezuelan mandioc	5.10 $\pm$ 0.2	2.34 $\pm$ 0.03
Colombian mandioc	5.10 $\pm$ 0.2	2.33 $\pm$ 0.02
Ocumo	4.98 $\pm$ 0.2	2.18 $\pm$ 0.02

## APPENDIX II

To test for the existence of alpha (1-4) glucosidase activity in pullulanase a maltotriose solution was treated with the enzyme at 37<sub>o</sub> C for 24 hours. The reducing power of the solution was determined before and after the treatment. by Nelson's copper procedure .

Reducing power

gluc. mgs / ml.

Before

After

2.78

2.78

2.74

2.79

2.78

2.75

2.76

2.74

$$\bar{X} = 2.76 \pm 0.02$$

$$\bar{X} = 2.76 \pm 0.02$$

## APPENDIX III

To evaluate the alpha (1-4) glucosidase contamination of isoamylase a maltoheptaose solution was treated with the enzyme at 37 °C for 24 hours. The reducing power of the solution was evaluated by Nelson's procedure before and after the treatment, each determination was done four times.

Reducing power  
gluc.mgs /ml.

Before enzyme treatment	After enzyme treatment
3.54	3.54
3.55	3.55
3.58	3.59
3.54	4.00
$\bar{X} = 3.55 \pm 0.02$	$\bar{X} = 3.67 \pm 0.22$

STRUCTURAL COMPARISON OF CEREAL AND TUBER  
AMYLOPECTINS

BY

BEATRIZ GUTIERREZ P.

B.A., National University of Colombia, 1973

---

AN ABSTRACT OF A MASTER THESIS

submitted in partial fulfillment of the

requirements for the degree

MASTER OF SCIENCE

Department of Grain Science and Industry

KANSAS STATE UNIVERSITY

Manhattan ,Kansas.

1985

The fine structure of amylopectin fractions from millet, (Penneasetum americanum) wheat ( Triticum aestiviun ) mandioc (Manihot sculenta crazt) and ocumo (Xanthosoma sacittifolium) were investigated using hydrolytic enzymes ( B- amylase, pullulanase, and isoamylase) in conjunction with gel filtration chromatography ( bio- gel P-10 and P-2). The unit chain profiles of debranched amylopectin fractions were characterized by two populations with average degrees of polimerization of 10-16 and 50-58 respectively in wheat and ocumo and 18-22 and >58 in millet and mandioc. The average chain length for these amylopectin fractions (16-24) ,the B-amyloysis limits (54-61%) and the A:B chain ratios (1.42-1.79) were internally consistent and led to the conclusion that millet, wheat and ocumo amylopectins possess more and shorter branches that dose mandioc amylopectin. In conclusion the amylopectin fractions study here appear to have definite but minor differences in structure.