

STUDIES ON ASCARIDIA COLUMBAE: EXPERIMENTAL LIFE CYCLE
IN PARENTERALLY INFECTED PIGEONS, AND FACTORS
AFFECTING THE OVIPOSITION IN VITRO

by

ROY DANIEL MELENDEZ,
DVM., Universidad Centro Occidental, 1970
Barquisimeto, Venezuela

A MASTER'S THESIS

submitted in partial fulfillment of the
requirements for the degree

MASTER OF SCIENCES

Department of Infectious Diseases

KANSAS STATE UNIVERSITY
Manhattan, Kansas
1977

Approved by:


Major Professor

Document
LD
2668
T4
1977
M44
C.2

TABLE OF CONTENTS

	<u>Page</u>
TABLE OF CONTENTS	i
LIST OF TABLES, GRAPHS, AND FIGURES	iv
SECTION I. INTRODUCTION AND LITERATURE REVIEW	1
SECTION II. MATERIALS AND METHODS	3
1. Pigeons	3
2. Source and Culture of Eggs	4
3. Experiment No. 1	5
a) Counting and administration of embryonated <u>Ascaridia columbae</u> eggs	5
b) Length of infection	6
c) Recovery of larvae and adult worms	6
d) Histopathologic techniques	7
4. Experiment No. 2.. . . .	7
a) Source of embryonated eggs	7
b) Hatching technique	7
c) Counting and administration of infective L2 <u>A. columbae</u> .	9
d) Length of infection	9
e) India ink injections	11
f) Recovery of larvae and adult worms	11
g) Histopathologic techniques	12

	<u>Page</u>
5. Experiment No. 3.	12
a) Origin of nematodes	12
b) Preparation of preconditioned saline media	13
c) Females <u>A. columbae</u> in normal and preconditioned saline media.	13
d) Technique for counting <u>A. columbae</u> eggs laid <u>in vitro</u>	15
e) pH of normal and preconditioned saline media.	17
6. Statistical Analysis	17
SECTION III. RESULTS	18
1. Experiment No. 1	18
a) Population of <u>A. columbae</u> recovered from orally infected pigeons	18
b) Number of <u>A. columbae</u> embryonated eggs required to produce one worm	19
c) Body length of <u>A. columbae</u> in 50 and 55 day infections. . .	19
d) Macroscopic and microscopic observations	20
2. Experiment No. 2.	23
a) Efficacy of hatching techniques	23
b) Distribution and stage of development of <u>A. columbae</u> in intravenously infected pigeons	23
c) Length of adult <u>A. columbae</u> and size of infection in intravenously infected pigeons.	31
d) India ink injections	31

	<u>Page</u>
e) Post-mortem examination. Macroscopical and microscopic observations	33
3. Experiment No. 3	34
a) Partial and total egg outputs of females <u>A. columbae</u> kept in natural and preconditioned saline media.	34
b) pH readings	36
SECTION IV. DISCUSSION	38
SECTION V. CONCLUSIONS	50
SECTION VI. ABSTRACT	51
SECTION VII. ACKNOWLEDGEMENTS	52
SECTION VIII. REFERENCES	53

LISTS OF TABLES, GRAFTS, AND PLATES

<u>TABLES</u>	<u>Page</u>
Table 1. Total number of <u>Ascaridia columbae</u> infective eggs necessary to produce 1 worm in orally infected pigeons	13
Table 2. Solutions and elements present in natural and pre- conditioned saline media	16
Table 3. Length of <u>Ascaridia columbae</u> from orally infected pigeons	21
Table 4. Efficacy of hatching techniques tested with <u>A. columbae</u> embryonated eggs	24
Table 5. Location and number of worms recovered from tissues and organs of pigeons intravenously infected with 600 \pm 50 L2 stage <u>Ascaridia columbae</u>	25
Table 6. pH values recorded for the different media at several intervals of time.	37

GRAPHS

Graph 1. Frequency distribution of the length of <u>Ascaridia columbae</u> kept in natural and preconditioned saline media.	32
Graph 2. Partial and total egg outputs of female <u>Ascaridia columbae</u> kept in natural and preconditioned saline media.	35

PLATES

Plate 1. Figure 1. Second stage or infective larva (L2) of <u>Ascaridia columbae</u> . Penetrating Knob (arrow); i, intestine. X350.	10
--	----

	<u>Page</u>
Figure 2. Posterior extremity of L2 <u>A. columbae</u> . Anus (arrow); t, tapering tail. X350	10
Plate 2. Figure 3. Liver section from orally infected pigeon showing granulomatous lesions. Note groups of giant cells (Gc) clustered around fragment of <u>A. columbae</u> larva (L.) Lymphocytes (arrow) are also involved in the inflammatory reaction. X350.	22
Figure 4. Liver section from orally infected pigeon showing multiple granulomas. Giant cells (Gc) of foreign-body type and mononuclear cells (arrow) surround fragments of <u>A. columbae</u> larvae (L). X350.	22
Figure 5. Liver section from orally infected pigeon showing germinal centre-like lesion limited by a layer of connective tissue (arrows). A group of giant cells (Gc) enclosing larval fragments (L) is also observed. X350	22
Plate 3. Figure 6. L2 <u>A. columbae</u> retrieved from lungs of pigeon 72 which had a pulmonary fungal disease. Some polymorphonuclear leucocytes (arrow) are adhered to the larva cuticle. This also shows wrinkles (w). X350.	26

	<u>Page</u>
Figures 7 and 8. Another L2 <u>A. columbae</u> obtained from the lungs of pigeon 72. Some inflammatory cells look like eosinophils (arrows). Fig. 7, X350.	
Fig. 8, X700..	26
Plate 4. Figure 9. Third stage larva (L3) of <u>Ascaridia columbae</u> collected from lungs of intravenously infected pigeons. Lips (arrow) are distinct but weakly developed. X350..	27
Figure 10. L3 <u>A. columbae</u> collected from lungs of intravenously infected pigeon at higher magnification. Rudimentary lips (arrow) are helpful structures identifying this larval stage..	27
Plate 5. Figure 11. Anterior extremity of fourth-stage larva of <u>A. columbae</u> obtained from the gut wall of intravenously infected pigeon. Note the unshed larval sheath (S) which shows the impression left by the lips (arrow) and the cervical ala (longest arrow). E, esophagus; i, intestine. X350	29
Figures 12 and 13. Anterior and posterior ends of a same L4 <u>A. columbae</u> collected from the gut wall of an intravenously infected pigeon. Two sheaths are seen on the same larva (arrows).. . . .	29