



Corn Response as Affected by Planting Distance from the Center of Strip-Till Fertilized Rows

Eric Adee*, Fernando D. Hansel, Dorivar A. Ruiz Diaz and Keith Janssen

Department of Agronomy, Kansas State University, Manhattan, KS, USA

OPEN ACCESS

Edited by:

David Bryla,
U.S. Department of Agriculture, USA

Reviewed by:

Hans-Peter Kaul,
University of Natural Resources and
Life Sciences, Austria

Oscar Urrutia,
University of Navarra, Spain

*Correspondence:

Eric Adee
eadee@ksu.edu

Specialty section:

This article was submitted to
Crop Science and Horticulture,
a section of the journal
Frontiers in Plant Science

Received: 27 April 2016

Accepted: 03 August 2016

Published: 18 August 2016

Citation:

Adee E, Hansel FD, Ruiz Diaz DA and
Janssen K (2016) Corn Response as
Affected by Planting Distance from the
Center of Strip-Till Fertilized Rows.
Front. Plant Sci. 7:1232.
doi: 10.3389/fpls.2016.01232

Strip-till has been used at a large scale in east central Kansas as an alternative to earlier planting dates under a no-till system. To determine the effects of planting corn (*Zea mays*) under previously established strip-tilled fertilized rows, experiments were conducted on an Osage silty clay loam soil in 2006 and 2008 and on a Woodson silt loam soil in 2009, 2010, and 2011 using three different planting distances from the strip-tilled fertilized rows (0, 10, 20, and 38 cm) with a strip-till operation performed between 1 and 73 days before planting. The depth of the strip-till fertilizer application was 13–15 cm below the soil surface. Corn that was planted 10 cm from the fertilized row showed greater early season growth, higher plant population, and grain yield. Planting 20 and 38 cm from the center of the fertilized rows showed none of the benefits that are typically associated with strip-tillage system. Enough time should be allowed between the strip-till operation and planting to reach satisfactory soil conditions (e.g., moist and firm seedbed). Our results suggest that the best location for planting strip-tilled fertilized corn vary depending on soil and climatic conditions as well as the time between fertilizer application with the strip-till operation and planting. With fewer number of days, planting directly on the center of fertilized strip-till resulted in decreased plant population and lower grain yield. However, the greatest yield benefit across different planting conditions was attained when planting within 10 cm of the strip.

Keywords: strip-till, corn, *Zea mays*, nutrient uptake, stand establishment, grain yield

INTRODUCTION

The ability to strip-till for corn production in east central Kansas has allowed farmers to address several challenges they have faced. With the clay pan soils that are susceptible to erosion because of limited water infiltration, no-till has offered protection against soil erosion (Unger and Vigil, 1998; Lamm et al., 2009). While the layer of crop residue has offered protection against erosion, it has increased soil moisture and reduced soil temperatures, thus limiting the opportunities to plant corn early (Vetsch and Randall, 2002; Perez-Bidegain et al., 2007).

Earlier planting dates for corn have been shown to improve yields by getting the corn farther along in the growing season, especially to reach pollination, before the hot/dry weather of July (Roozeboom et al., 2009). Strip-till allows for residue to remain over much of the field to protect against erosion (Vetsch and Randall, 2002), while allowing the soil to warm and dry in the planting zone. The strip-till method also allows for banding of fertilizers to save a trip across the field and to place fertilizers closer to the plants for more efficient utilization (Fernandez and White, 2012).

TABLE 4 | Significance of F-values for the fixed effects of planting distance from the center of strip-till row on corn growth, nutrient concentration, nutrient uptake, population, and grain yield.

Variable	Years					Across years
	2006	2008	2009	2010	2011	
	P > F					
Plant dry weight V2–V3	0.030	0.001	0.018	–†	<0.001	<0.001
N concentration V2–V3	0.028	0.012	0.033	–	0.029	<0.001
P concentration V2–V3	0.304	0.027	0.005	–	0.243	0.001
K concentration V2–V3	0.790	<0.001	0.014	–	0.014	<0.001
N Uptake V2–V3	0.019	0.001	0.025	–	0.072	<0.001
P Uptake V2–V3	0.100	0.007	0.115	–	0.124	0.001
K Uptake V2–V3	0.031	0.001	0.018	–	0.039	<0.001
Plant dry weight V7–V8	0.651	0.001	0.008	0.002	0.001	<0.001
N concentration V7–V8	0.211	0.014	0.487	0.002	0.070	0.013
P concentration V7–V8	0.329	0.348	0.872	0.021	0.660	0.220
K concentration V7–V8	0.663	0.720	0.335	0.712	0.203	0.147
N Uptake V7–V8	0.909	0.001	0.016	0.005	0.217	<0.001
P Uptake V7–V8	0.867	0.001	0.014	0.009	0.179	<0.001
K Uptake V7–V8	0.995	0.002	0.011	0.172	0.101	<0.001
Plant population	0.046	0.054	0.834	<0.001	0.001	<0.001
Grain yield	0.047	0.008	0.338	0.127	0.365	0.002

†Not available.

TABLE 5 | Plant dry weight, nutrient concentration, and nutrient uptake at the V2–V3 growth stage†.

Distance‡ cm	Plant dry weight g plant ⁻¹	Nutrient concentration			Nutrient uptake		
		Nitrogen	Phosphorus g kg ⁻¹	Potassium	Nitrogen	Phosphorus mg plant ⁻¹	Potassium
2006							
0	0.33b§	3.9a	0.26	2.1	13.0cb	0.88	7.0b
10	0.49a	3.9a	0.30	2.2	19.1a	1.47	10.7a
20	0.40ab	3.6ab	0.31	2.2	14.8ab	1.30	8.7ab
38	0.30b	3.0b	0.28	2.3	9.2c	0.87	6.8b
2008							
0	0.44ab	3.5a	0.33b	2.0a	15.4a	1.47ab	9.0a
10	0.47a	3.4a	0.33b	1.9b	15.8a	1.56a	9.0a
20	0.38b	3.3a	0.33b	1.6c	12.7b	1.27b	6.0b
38	0.23c	2.9b	0.42a	1.4d	6.6c	0.95c	3.1c
2009							
0	0.80a	4.0a	0.4b	2.3a	31.7a	3.17	18.9a
10	0.71a	4.0a	0.4b	2.2a	28.4a	2.85	15.7a
20	0.62ab	4.0a	0.44a	2.0a	25.5ab	2.73	12.9ab
38	0.46b	3.7b	0.46a	1.4b	17.0b	2.09	6.5b
2011							
0	0.56a	4.0a	0.41	2.3a	30.9a	3.17	18.4a
10	0.52b	4.0a	0.40	2.2a	28.4a	2.85	15.7a
20	0.37c	4.0a	0.44	2.0a	25.5ab	2.73	12.9ab
38	0.35c	3.7b	0.44	1.4b	17.8b	2.10	7.0b

†Early growth data (V2–V3 growth stage) for 2010 was not collected.

‡Distance from the center of strip till row.

§Numbers followed by the same letter in a column within each year are not statistically significant at the 0.05 probability level.

TABLE 6 | Plant dry weight, nutrient concentration, and nutrient uptake at the V7–V8 growth stage.

Distance [†] cm	Plant dry weight g plant ⁻¹	Nutrient concentration			Nutrient uptake			Population Plants ha ⁻¹	Grain yield Mg ha ⁻¹
		Nitrogen	Phosphorus	Potassium	Nitrogen	Phosphorus	Potassium		
		g kg ⁻¹			mg plant ⁻¹				
2006									
0	6.5	4.3	0.48	1.9	279	30.9	124	58924b [‡]	8.2b
10	6.4	4.3	0.47	1.9	274	30.3	119	63015a	8.9a
20	6.6	4.2	0.46	1.8	281	30.2	121	63407a	8.5ab
38	5.7	4.6	0.50	2.1	262	28.2	119	59858ab	8.7ab
2008									
0	6.9a	2.8a	0.33	1.8	195a	22.7a	125ab	56128b	6.0cb
10	6.9a	2.8a	0.36	2.0	195a	24.5a	138a	57956a	6.5a
20	5.5a	2.8a	0.36	1.8	154b	19.9a	99b	55876b	6.3ab
38	2.4b	2.5b	0.33	1.9	59c	7.5b	44c	55805b	5.7c
2009									
0	19.5a	3.0	0.41	2.6	586a	79.0a	501a	56998	5.0
10	18.8a	3.3	0.40	2.1	609a	75.8a	392ab	57833	5.2
20	16.1a	3.4	0.42	2.1	536a	67.0a	341b	56998	4.7
38	11.8b	3.2	0.41	2.4	375b	48.4b	278b	56486	4.5
2010									
0	12.7a	3.7c	0.49b	3.3	471a	63.0a	437	45609a	3.7
10	9.7b	4.1b	0.54a	3.5	392ab	53.3ab	392	48118a	3.9
20	7.0c	4.3ab	0.53a	3.4	297cb	37.2cb	249	40407b	3.8
38	5.3c	4.5a	0.55a	3.9	236c	29.4c	221	30067c	3.3
2011									
0	8.6a	2.8b	0.40	2.7	536	74.1	481	57213a	1.8
10	8.0a	3.3ab	0.40	2.1	609	75.8	392	57884a	1.8
20	5.9b	3.4a	0.42	2.1	536	67.0	341	52317b	1.8
38	5.0b	3.3a	0.42	2.3	425	53.4	298	49978b	1.5

[†]Distance from the center of strip till row.

[‡]Numbers followed by the same letter in a column within each year are not statistically significant at the 0.05 probability level.

(warmer soil, cleared residue, and rapid fertilizer-root contact). The objective of this study was to evaluate the effect of planting distance from the center of strip-tilled fertilized rows for corn production.

MATERIALS AND METHODS

Field experiments were conducted on an Osage silty clay loam soil (Fine, smectitic, thermic Typic Epiaquerts, on 0–1% slope, land capability class IIw) near Lane, Kansas (38°25'N 95°07'W), in 2006 and 2008, and on a Woodson silt loam soil (Fine, smectitic, thermic Abruptic Argiaquolls, on 0–1% slope, land capability class IIs) at the East Central Kansas Experiment Field near Ottawa, Kansas (38°32'N 95°14'W), in 2009, 2010, and 2011. Both locations receive an average annual precipitation of ~1017 mm (Kansas State Weather Data Library, 2012). Soil samples were collected at the 0- to 15-cm depth from the fields where the studies were located. Samples were analyzed

for soil test P (STP) using the Mehlich-3 extraction, and for soil test K (STK) using the ammonium acetate extraction (Warncke and Brown, 1998). Soil pH was measured in a 1:1 water suspension, and total organic matter was determined by loss on ignition (Combs and Nathan, 1998). Soil texture was estimated by the hydrometer method (Model 6026Q20, Thomas Scientific; Bouyoucos, 1962). Cation exchange capacity (CEC) was estimated by summing the exchangeable acidity and the exchangeable bases (Warncke and Brown, 1998, **Table 1**).

The combination of strip-till and fertilizer application was performed 2–73 days before planting (**Table 2**). Fertilizer was applied at a standard rate (134-34-11 kg ha⁻¹ of N, P, and K, respectively). The fertilizer source used for this study include urea, diammonium phosphate, and potassium chloride. The depth of the fertilizer application with the strip-till was 13–15 cm. The applicator was a Yetter (Yetter Mfg., Colchester, IL) pull caddy with Maverick Generation 2 openers and residue managers (model 2984) and a 5 cm mole knife, equipped with a Gandy

Orbit Air model 623016 box and metering system (Gandy Co, Owatanna, MN).

Planting was with a White 6100 planter (AGCO, Duluth, GA). The planting distances evaluated were directly on top of the strip-tilled fertilized rows and ~10, 20, and 38 cm off the center of the rows. The experiment was designed as a randomized complete block with three to four replications. Individual plot size ranged from 3.04 m (4 rows × 76 cm) wide and 48 m long to 6.08 (8 rows × 76 cm) and 240 m long, depending on the field location and year.


The planting treatments were evaluated for effects on early season corn growth, nutrient uptake, plant population at harvest, and grain yield. Early growth assessments were made by collecting the above ground portion of 12 plants at growth stage V2 to V3 (Abendroth et al., 2011), and the above ground portion of six plants at the V7 to V8 growth stage (Abendroth et al., 2011) from each plot. Plant dry weight, and nutrient concentration (N, P, and K) were measured, and nutrient uptake calculated based on these two parameters. Plant samples were dried at 65°C in a forced-air oven, weighed, and ground to pass through a 2-mm sieve. Nutrient analysis was completed on digested plant materials using the sulfuric acid-hydrogen peroxide (H₂SO₄-H₂O₂) method. Nitrogen and phosphorus in the extract was determined with the Technicon AAI autoanalyzer, and potassium by Flame Atomic Absorption. Plant populations at harvest were measured by counting plants in 48–131 m of row, depending on the length of row each year. The center two rows of each plot were harvested with a plot combine, grain weight, and moisture were measured, and yields adjusted to 155 g kg⁻¹ moisture.

Statistical analysis was completed with SAS 9.4 (SAS Institute, 2014) using the GLIMMIX procedure. Significance level was set at the $P < 0.05$ for mean separation and using the LINES procedure of GLIMMIX. Anova analysis was completed by site-year using blocks as random in the model; and across site-years and using blocks and site-years as random effect in the model. Our statistical analysis used site-year as random in the model given that this factor can represent a larger population with a probability distribution in addition to the exchangeability of its effect. Our discussion and conclusions put emphasis on the main treatment effects across 5 site-years and the potential applicability for producers in future years.

RESULTS AND DISCUSSION

Total rainfall from January to September was lower than average for 2006 and 2011 (Table 3). The amount of rainfall after the strip till operation and planting was also lower for years 2006 (zero) and 2011 (39 mm; Table 2). Dry soil conditions can be associated with higher potential for seedling damage from the fertilizer application as well as poor soil condition for early plant establishment (Perez-Bidegain et al., 2007).

Across site-year analysis of all the parameters evaluated in the study showed significant effect by planting distance from the center of fertilized strip till except for P and K concentration at



the V7–V8 growth stage (Table 4). Above-ground corn biomass early in the season (V2–V3, and V7–V8) was affected by planting distance treatments for each year during the study, other parameters were statistically significant only for some years or across years (Table 4).

Corn early growth at the V2–V3 and V7–V8 growth stages were generally higher for corn planted directly on top of the strip-tilled fertilized rows or within 10 cm (Tables 5, 6, Figure 1). Previous studies showed similar results where the beneficial effects of in-row tillage and residue removal in the seed-row zone with strip-till led to greater early growth of corn (Vetsch et al., 2007). Planting corn 20 cm from the center of the strip-tilled fertilized rows reduced early season growth of corn at the V7–V8 growth stage by 24% on average, and planting 38 cm away reduced early season growth by 40%. Cool conditions slow the germination of corn seed and slow shoot elongation before emergence influencing early season corn growth (Miedema and Sinnaeve, 1980).

There was no consistent pattern for nutrient concentration in the plants at different planting distance from the row at the V2–V3 and V7–V8 growth stages (Tables 5, 6, Figure 2). This could be due the variability in the size of plants at different distances from the row at these growth stages. For example, a




FIGURE 2 | Nitrogen, phosphorus, and potassium concentration as affected by distance from the center of strip-till fertilized row across site-years. Samples collected at to growth stage, V2–V3 and V7–V8.

plant farther from the center of the row could have a higher concentration of nutrients but lower biomass due to cooler soil temperatures, and as a result have less total nutrient uptake. Plants may lose the potential for growth even though they have accumulated “adequate” concentrations of the limiting element (Hiatt and Massey, 1958); on the other hand when the growth-limiting element is supplied (e.g., higher soil temperatures) the relative rate of dry matter accumulation increases more rapidly

than the rate of nutrient accumulation (Jarrell and Beverly, 1981).

Uptake of plant nutrients (i.e., nitrogen, phosphorus, and potassium) followed a pattern similar to that for plant growth (Tables 5, 6, Figure 3). Nutrient uptake tended to be greater in plants growing closer to the band of fertilizer (Tables 5, 6, Figure 3). Nitrogen, P and K uptake was less when corn was planted 38 cm from the strip-till row (Tables 5, 6). The trend





FIGURE 3 | Nitrogen, phosphorus, and potassium uptake as affected by distance from the center of strip-till fertilized row across site-years. Samples collected at to growth stage, V2-V3 and V7-V8.

for decreasing uptake of N, P, and K by the corn plants the farther they are from the row suggest that the plants lost the beneficial effect of tillage and fertilizer placement from strip-till (Figure 3). Plants within 10 cm of the row well as those over the row showed higher plant biomass and nutrient uptake likely from the combined effect of localized tillage (strip-till) and fertilizer placement. However, this increase in early plant growth and utilization of the fertilizer applied with the strip-till did not always result in higher grain yields (Table 6). Other yield limiting factors during the growing season, such as moisture and temperature had significant role in the resulting yield and the potential benefit of planting distance from the strip-till and

fertilizer band. This was particularly the case for 2006 with below average rainfall (Tables 3, 6).

Average corn plant populations were higher when planted ~10 cm off the center of the strip-tilled fertilized rows compared with planting directly on top of the rows for every year and across years (Table 6, Figure 4). These results were statistically significant for 2 years with relatively different planting and weather conditions (2006 and 2008; Tables 2, 3). In 2006 the strip-till fertilization operation was completed 2 days before planting likely contributing to loose soils with air pockets under the row. However, in 2008 plating was completed 2 weeks after the strip-tillage operation and plant population was



still higher when planted 10 cm off the center of the strip-tilled fertilized rows. Very small differences in plant populations occurred in 2009, when planting was completed 73 days after the strip-till operation. These results are similar to those of Bordoli and Mallarino (1998). They found that deep and shallow fertilizer banding did not affect plant population with any tillage treatment when applied 3–5 weeks before planting with stabilized soil with firm seedbed. In 2010, high rainfall around planting generated waterlogged and cold conditions (Tables 2, 3). This year showed the overall lowest plant population of any year of the study with significantly higher values when planting within 10 cm of the center of the row (Table 6). Low soil temperatures, a consequence of untilled soil may cause slow plant growth, and in some cases injuries and chlorosis in corn, and therefore affecting plant population (Miedema, 1982; Perez-Bidegain et al., 2007). Plant stand reduction can also occur when fungal pathogens rot seed or kill seedlings in corn when cold and wet conditions are present after planting (Nyvall, 1999). Planting 20 cm off the center of the strip-tilled fertilized rows reduced plant population 3122 plants ha⁻¹, and 38 cm off reduced population 7308 plants ha⁻¹ compared to planting 10 cm from the center. Early growth was reduced 38 and 53%, respectively (Tables 5, 6). Effects for 2011 were similar to 2010. Averaged across all years, the highest plant population was 10 cm off the row (Figure 4).

Across all years, the grain yield was greatest when corn was planted 10 cm off the center of the row (Figure 4). Each year had different confounded factors that contributed to the overall greatest yield with that treatment. In 2006, when corn was planted 10 cm off the center of the rows grain yield was ~8% higher than corn planted directly on top of the strip-tilled fertilized rows (Table 6). This was likely in part due to a significant reduction in plant population (1656 fewer plants ha⁻¹). Results were similar in 2008 with higher corn population for treatments planted 10 cm off the center of the strip-tilled fertilized rows, this also resulted in average higher grain yield. In the 2009 the strip-till operation was completed 73 days before planting, and under these conditions there were no statically significant differences in plant population or grain yield between planting distance treatments. However, average higher grain yield was observed for treatments with planting within 10 cm of the center of the strip-till row. Yield response to planting distance from the center of strip-till was not significant in 2010 (Table 6). It is likely that overall lower plant population and excessive rainfall contributed to plant stress including nitrogen loss. In 2011, under extreme heat, drought (Campos et al., 2006), and near crop failure conditions, all treatments yielded below 1.9 Mg ha⁻¹, and planting distances from the row had no significant effect on yield.

These results suggest that the best location for planting strip-tilled fertilized corn will vary depending on the growing season, condition of the strip-tilled fertilized zone, and the amount of time between planting and when the strip-tilled fertilizer operation was performed.

In a year when a strip-till operation is just prior to planting, planting 10 cm off the strip would be advisable. If the strip-till operation is well in advance of planting, and the soil has had time to settle, then planting directly over the strip or within 10 cm showed similar results. However, our results show no benefit of planting at more than 10 cm from the center of the strip-till row where the benefits of the localized tillage and fertilizer application disappeared. Strip-tilled fertilized corn should be planted in a moist, firm seedbed to obtain best stands and within 10 cm of strip-tilled fertilized rows to ensure quick contact between corn roots and fertilizer without causing potential damage to the seedlings.

Additional studies are needed to determine if these results might be different when planting strip-tilled fertilized corn on coarse-textured soils and when higher rates of fertilizer and other sources of nitrogen, such as anhydrous ammonia, are used. Phytotoxicity due to nutrients being too close to the seedling could be affected by the timing and distance of planting and more relevant for some fertilizer sources such as anhydrous ammonia.

CONCLUSIONS

There are many benefits to planting corn into strips made in no-till fields ahead of time, but there can be disadvantages under some circumstances. Results from this study suggest that all the benefits can be obtained if planting is done within 10 cm of

the center of the strip, which also reduces many of the risks associated with planting directly over the strip, such as poor or uneven stand establishment. The benefits of planting in the strips with fertilizer applied include a warmer, drier seed bed which allows for earlier planting, improved stand establishment, and early season growth than could be realized in the same field in no-till, but still maintains many of the advantages of the residue cover that no-till offers. Facilitating early season growth and development of corn in dryland production to avoid heat and drought stress at critical development stages later in the season can be very important in avoiding yield loss.

REFERENCES

- Abendroth, L. J., Elmore, R. W., Boyer, M. J., and Marlay, S. K. (2011). *Corn Growth and Development. PMR 1009*. Ames, IA: Iowa State University.
- Bordoli, J. M., and Mallarino, A. P. (1998). Deep and shallow banding of phosphorus and potassium as alternatives to broadcast fertilization for no-till corn. *Agron. J.* 90, 27–33.
- Bouyoucos, G. J. (1962). Hydrometer method improved for making particle size analyses of soils. *Agron. J.* 54, 464–465.
- Campos, H., Cooper, M., Edmeades, G. O., Löffler, C., Schussler, J. R., and Ibanez, M. (2006). Changes in drought tolerance in maize associated with fifty years of breeding for yield in the US Corn Belt. *Maydica* 51, 369–381.
- Combs, S. M., and Nathan, M. V. (1998). "Soil organic matter," in *Recommended Chemical Soil Test Procedures for the North Central Region*, North Central Regional Research Publication No. 221 (revised), ed J. R. Brown (Columbia, MO: Missouri Agricultural Experiment Station), 53–58.
- Cresswell, H. P., Painter, D. J., and Cameron, K. C. (1993). Tillage and water content effects on surface soil hydraulic properties and shortwave albedo. *Soil Sci. Soc. Am. J.* 57, 816–824.
- Fernandez, F. G., and Schaefer, D. (2011). Assessment of soil phosphorus and potassium following real time kinematic-guided broadcast and deep-band placement in strip-till and no-till. *Soil Sci. Soc. Am. J.* 76, 1090–1099. doi: 10.2136/sssaj2011.0352
- Fernandez, F. G., and White, C. (2012). No-till and strip-till corn production with broadcast and subsurface-band phosphorus and potassium. *Agron. J.* 104, 996–1005. doi: 10.2134/agronj2012.0064
- Hakansson, I., Myrbeck, A., and Etana, A. (2002). A review of research on seedbed preparation for small grains in Sweden. *Soil Tillage Res.* 64, 23–40. doi: 10.1016/S0167-1987(01)00255-0
- Hallauer, A. R., and Colvin, T. S. (1985). Corn hybrids response to four methods of tillage. *Agron. J.* 77, 547–550. doi: 10.2134/agronj1985.00021962007700040010x
- Hiatt, A., and Massey, H. (1958). Zinc levels in relation to zinc content and growth of corn. *Agron. J.* 50, 22–24. doi: 10.2134/agronj1958.00021962005000010007x
- Jarrell, W. M., and Beverly, R. B. (1981). "The dilution effect in plant nutrition studies," in *Advances in Agronomy*, ed N. C. Brady (New York, NY: Academic Press), 197–224.
- Kansas State Weather Data Library (2012). Available online at: <https://www.ksre.k-state.edu/wdl/>

AUTHOR CONTRIBUTIONS

Conceived, designed by KJ. Performed the experiment under field conditions: KJ and EA. Performed statistical analysis: EA, DD, FH, JK. All authors wrote and revised the paper.

FUNDING

This research was partially supported by funds from Department of Agronomy and Kansas State Research and Extension. This is contribution no. 16-292-J from the Kansas Agricultural Experiment Station.

- Lamm, F. R., Aiken, R. M., and Abou Kheira, A. A. (2009). Corn yield and water use characteristics as affected by tillage, plant density, and irrigation. *Trans. ASABE* 52, 133–143. doi: 10.13031/2013.25954
- Miedema, P. (1982). "The effects of low temperature on *Zea mays*," in *Advances in Agronomy*, Vol. 35, ed N. C. Brady (San Diego, CA: Academic Press), 93–125.
- Miedema, P., and Sinnaeve, J. (1980). Photosynthesis and respiration of maize seedlings at suboptimal temperatures. *J. Exp. Bot.* 31, 813–819. doi: 10.1093/jxb/31.3.813
- Nyvall, R. F. (1999). *Field Crop Diseases*. Ames, IA: Iowa State University Press.
- Perez-Bidegain, M., Cruse, R. M., and Ciha, A. (2007). Tillage system by planting date interaction effects on corn and soybean yield. *Agron. J.* 99, 630–636. doi: 10.2134/agronj2006.0058
- Roozeboom, L., Claassen, M., and Maddux, L. (2009). *Pushing the Early Corn Planting Envelope in No-till On High-Clay Soils*. Agronomy Abstracts. ASA Madison, WI.
- SAS Institute (2014). *The SAS System for Windows. Release 9.3*. Cary, NC: SAS Institute.
- Unger, P., and Vigil, M. F. (1998). Cover crop effects on soil/water relationships. *J. Soil Water Cons.* 53, 200–207.
- Vetsch, J. A., and Randall, G. W. (2002). Corn production as affected by tillage system and starter fertilizer. *Agron. J.* 94, 532–540. doi: 10.2134/agronj2002.5320
- Vetsch, J. A., Randall, G. W., and Lamb, J. A. (2007). Corn and soybean production as affected by tillage systems. *Agron. J.* 99, 952–959. doi: 10.2134/agronj2006.0149
- Warncke, D., and Brown, J. R. (1998). "Potassium and other cations," in *Recommended Chemical Soil Test Procedures for the North Central Region*, North Central Regional Research Publication, 221 (Rev.), ed J. R. Brown (Columbia, MO: Missouri Agric. Exp. Stn. SB 1001), 31–33.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Adee, Hansel, Ruiz Diaz and Janssen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.