

Spectral Properties of Schrödinger-type Operators and Large-time Behavior of the Solutions to the Corresponding Wave Equation

A.G. Ramm *

Department of Mathematics, Kansas State University, Manhattan, KS 66506-2602, USA

Abstract. Let L be a linear, closed, densely defined in a Hilbert space operator, not necessarily selfadjoint. Consider the corresponding wave equations

$$(1) \quad \ddot{w} + Lw = 0, \quad w(0) = 0, \quad \dot{w}(0) = f, \quad \dot{w} = \frac{dw}{dt}, \quad f \in H.$$
$$(2) \quad \ddot{u} + Lu = fe^{-ikt}, \quad u(0) = 0, \quad \dot{u}(0) = 0,$$

where $k > 0$ is a constant. Necessary and sufficient conditions are given for the operator L not to have eigenvalues in the half-plane $\operatorname{Re} z < 0$ and not to have a positive eigenvalue at a given point $k_d^2 > 0$. These conditions are given in terms of the large-time behavior of the solutions to problem (1) for generic f .

Sufficient conditions are given for the validity of a version of the limiting amplitude principle for the operator L .

A relation between the limiting amplitude principle and the limiting absorption principle is established.

Keywords and phrases: elliptic operators, wave equation, limiting amplitude principle, limiting absorption principle

Mathematics Subject Classification: 35P25, 35L90, 43A32

1. Introduction

Let L be a linear, densely defined, closed operator in a Hilbert space H . Our results and techniques are valid in a Banach space also, but we wish to think about L as of a Schrödinger-type operator in a Hilbert space and, at times, think that L is selfadjoint. For a Schrödinger operator $L = -\nabla^2 + q(x)$ the resolvent $(L - k^2)^{-1}$, $\operatorname{Im} k > 0$, is an integral operator with a kernel $G(x, y, k)$, its resolvent kernel. If q is a real-valued function, sufficiently rapidly decaying then L is selfadjoint, $G(x, y, k)$ is analytic with respect to k in the half-plane $\operatorname{Im} k > 0$, except, possibly, for a finitely many simple poles ik_j , $k_j > 0$, the

*Corresponding author. E-mail: ramm@math.ksu.edu

semiaxis $k \geq 0$ is filled with the points of absolutely continuous spectrum of L , and there exists a limit

$$\lim_{\epsilon \rightarrow 0} G(x, y, k + i\epsilon) = G(x, y, k)$$

for all $k > 0$.

Sufficient conditions for $k^2 = 0$ not to be an eigenvalue of L are found in papers [5], [6]. Spectral analysis of the Schrödinger operators is presented in many books (see, for example, [2] and [11]). In papers [3], [4], such an analysis was given in a class of domains with infinite boundaries apparently for the first time, see also [8]. In [7] an eigenfunctions expansion theorem was proved for non-selfadjoint Schrödinger operators with exponentially decaying complex-valued potential q . The operator L in this paper is not necessarily assumed to be selfadjoint.

In [1] the validity of the limiting amplitude principle for some class of selfadjoint operators L has been established.

This principle says that, as $t \rightarrow \infty$, the solution to problem

$$\ddot{u} + Lu = fe^{-ikt}, \quad u(0) = 0, \quad \dot{u}(0) = 0, \quad \dot{u} = \frac{du}{dt}, \quad (1.1)$$

has the following asymptotics

$$u = e^{-ikt}v + o(1), \quad t \rightarrow \infty, \quad (1.2)$$

where k is a real number and $v \in H$ solves the equation

$$Lv - k^2v = f. \quad (1.3)$$

The v is called *the limiting amplitude*. It turns out that a more natural definition of the limiting amplitude is:

$$v = \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t u(s)e^{iks} ds, \quad (1.4)$$

if this limit exists and solves equation (1.3).

Why is this definition more natural than (1.2)? There are good reasons for this. One of the reasons is: if (1.2) and (1.3) hold, then the limit (1.4) exists and solves equation (1.3). The other reason is: the limit (1.4) may exist and solve equation (1.3) although the limit (1.2) does not exist.

Example. If $u = e^{ikt}v + e^{ik_1t}v_1$, then the limit (1.2) does not exist, while the limit (1.4) does exist and is equal to v .

To describe our assumptions and results, some preparation is needed.

Consider the problem

$$\ddot{w} + Lw = 0, \quad w(0) = 0, \quad \dot{w}(0) = f. \quad (1.5)$$

Assuming that $\|u(t)\| \leq ce^{at}$, where $c > 0$ stands throughout the paper for various generic constants, and $a \geq 0$ is a constant, one can define the Laplace transform of $u(t)$,

$$\mathcal{U} := \mathcal{U}(p) := \int_0^\infty e^{-pt}u(t)dt, \quad \sigma > a,$$

where $p = \sigma + i\tau$, $\text{Re} p = \sigma$.

Let us take the Laplace transform of (1.1) and of (1.5) to get

$$L\mathcal{U} + p^2\mathcal{U} = \frac{f}{p + ik}, \quad (1.6)$$

and

$$L\mathcal{W} + p^2\mathcal{W} = f, \quad (1.7)$$

where

$$\mathcal{W} = \mathcal{W}(p) = \int_0^\infty w(t)e^{-pt} dt.$$

We also denote $\mathcal{W}(p) := \bar{w}(t)$.

The complex plane p is related to the complex plane k by the formula

$$p = -ik, \quad k = k_1 + ik_2, \quad k_2 \geq 0, \quad \sigma = k_2 \geq 0. \tag{1.8}$$

We assume throughout that f is generic in the following sense:

If I is the identity operator and a point p is a pole of the kernel of the operator $(L + p^2I)^{-1}$, then it is a pole of the same order of the element $(L + p^2I)^{-1}f = \mathcal{W}$.

If k^2 is an eigenvalue of L and $\operatorname{Re} k^2 < 0$, then $\operatorname{Im} k > 0$, where $k = |k|e^{\frac{i \arg k^2}{2}}$, $p = -ik$, so $\sigma = \operatorname{Re} p > 0$. Let $k > 0$ and assume that $-k^2 < 0$ is an eigenvalue of L . Then ik is a pole of the resolvent kernel $G(x, y, k)$, and $p = -i(ik) = k$ is a pole of the kernel of the operator $(L + p^2I)^{-1}$. If $k^2 > 0$ is an eigenvalue of L , then $p = -ik$ is a pole of the operator $(L + p^2I)^{-1}$.

The following known facts from the theory of Laplace transform will be used.

Proposition 1.1. *An analytic in the half-plane $\sigma > \sigma_0 \geq 0$ function $F(p)$ is the Laplace transform of a function $f(t)$, such that $f(t) = 0$ for $t < 0$ and*

$$\int_0^\infty |f(t)|^2 e^{-2\sigma_0 t} dt < \infty \tag{1.9}$$

if and only if

$$\sup_{\sigma > \sigma_0} \int_{-\infty}^\infty |F(\sigma + i\tau)|^2 d\tau < \infty. \tag{1.10}$$

Proposition 1.2. *If $F(p) = \overline{f(t)}$, then*

$$\frac{F(p)}{p} = \overline{\int_0^t f(s) ds}. \tag{1.11}$$

Let us now formulate the main Assumptions A and B standing throughout this paper.

Assumption A. For a generic f the $\mathcal{W}(p) = (L + p^2)^{-1}f$ is analytic in the half-plane $\sigma > 0$, except, possibly, at a finitely many simple poles at the points $-ik_j$, $1 \leq j \leq J$, k_j are real numbers, and at the points κ_m , $\operatorname{Re} \kappa_m > 0$,

$$\mathcal{W}(p) = \sum_{j=1}^J \frac{v_j}{p + ik_j} + \mathcal{W}_1(p) + \sum_{m=1}^M \frac{b_m}{p - \kappa_m}, \tag{1.12}$$

where v_j and b_m are some elements of H , $\mathcal{W}_1(p)$ is an analytic function in the half-plane $\operatorname{Re} p = \sigma > 0$, continuous up to the imaginary axis $\sigma = 0$, and satisfying the following estimate

$$\|\mathcal{W}_1(p)\| \leq \frac{c}{1 + |p|^\gamma}, \quad \gamma > \frac{1}{2}. \tag{1.13}$$

Assumption B. There exists the limit

$$\lim_{\sigma \rightarrow 0} \|\mathcal{W}_1(\sigma - ik) - \mathcal{W}_1(-ik)\| = 0 \tag{1.14}$$

for all real numbers k .

Theorem 1.3. *Let the Assumption A hold. Then a necessary and sufficient condition for the operator L to have no eigenvalues in the half-plane $\operatorname{Re} k^2 < 0$ is the validity of the estimate*

$$\left\| \int_0^t w(s) ds \right\| = O(e^{\epsilon t}), \quad t \rightarrow \infty, \quad (1.15)$$

for an arbitrary small $\epsilon > 0$.

A necessary and sufficient condition for the operator L not to have any positive eigenvalues $k^2 > 0$ is the validity of the estimate

$$\left\| \frac{1}{t} \int_0^t e^{iks} w(s) ds \right\| = o(1), \quad t \rightarrow \infty, \quad \forall k \in \mathbb{R}. \quad (1.16)$$

A point $ik_0 > 0, k_0 > 0$, is not a pole of the resolvent kernel of the operator $(L - k^2 - i0)^{-1}$ if and only if estimate (1.16) holds with $k = k_0 > 0$.

Remark. If condition (1.16) holds for $k = 0$, then $\left\| \int_0^t w(s) ds \right\| = o(t)$, so condition (1.15) holds, and the operator L has no eigenvalues in the half-plane $\operatorname{Re} k^2 < 0$.

Theorem 1.4. *Let the Assumptions A and B hold. Suppose that estimates (1.14) and (1.15) hold. Then the limiting amplitude principle (1.4) holds for every $k \in \mathbb{R}, k \neq k_j, 1 \leq j \leq J$.*

In section 2, proofs are given.

2. Proofs

2.1. Proof of Theorem 1.3

From the Assumption A and Proposition 1.1, it follows that $\mathcal{W}(p)$ is a Laplace transform of a function $w(t)$ such that

$$w(t) = \sum_{j=1}^J v_j e^{-ik_j t} + \sum_{m=1}^M b_m e^{\kappa_m t} + w_1(t), \quad (2.1)$$

where

$$w_1(t) = \frac{1}{2\pi i} \int_{\sigma_0 - i\infty}^{\sigma_0 + i\infty} e^{pt} \mathcal{W}_1(p) dp, \quad (2.2)$$

and the integral in (2.2) converges in L^2 -sense due to the assumption (1.13). It is clear from formula (2.1) that all $b_m = 0$ if and only if estimate (1.15) holds with $0 < \epsilon < \min_{1 \leq m \leq M} \operatorname{Re} \kappa_m$. This proves the first conclusion of Theorem 1.3.

Let us calculate the expression on the left side of formula (1.16) and show that this expression is $o(1)$ unless $k = k_j$ for some $1 \leq j \leq J$. In this calculation it is assumed that L does not have any eigenvalues in the half-plane $\operatorname{Re} k^2 < 0$, in other words, that all $b_m = 0$. Otherwise the expression on the left of formula (1.16) tends to infinity as $t \rightarrow \infty$ at an exponential rate.

If all $b_m = 0$ in (2.1), then

$$\sum_{j=1}^J v_j \frac{1}{t} \int_0^t e^{i(k-k_j)t} dt + \frac{1}{t} \int_0^t w_1(t) e^{ikt} dt := I_1 + I_2. \quad (2.3)$$

If k and k_j are real numbers, then

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t e^{i(k-k_j)t} dt = \begin{cases} 1, & k = k_j, \\ 0, & k \neq k_j. \end{cases} \quad (2.4)$$

Thus, $I_1 = 0$ if and only if k does not coincide with any of k_j , $1 \leq j \leq J$.

Let us prove that

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t w_1(t) e^{ikt} dt = 0. \quad (2.5)$$

By proposition (1.2) and the Mellin inversion formula, one has

$$I := \frac{1}{t} \int_0^t w_1(t) e^{ikt} dt = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \mathcal{W}_1(p-ik) \frac{e^{pt}}{pt} dp, \quad (2.6)$$

where $\text{Re } p = \sigma > 0$ can be chosen arbitrarily small.

Let $pt = q$, take $\sigma = \frac{1}{t}$, write $q = 1 + is$, and write the integral on the right side of (2.6) as:

$$I = \frac{1}{2\pi i} \int_{1-i\infty}^{1+i\infty} \mathcal{W}_1\left(\frac{q}{t} - ik\right) \frac{q e^q}{t q^2} dq. \quad (2.7)$$

If one uses estimate (1.13) and formula $|q| = (1 + s^2)^{1/2}$, then one obtains the following inequality

$$\|I\| \leq \frac{1}{2\pi t} \int_{-\infty}^{\infty} \frac{1}{(1 + s^2)^{1/2}} \frac{c ds}{[1 + |\frac{1+is}{t} - ik|^\gamma]} = \frac{c}{2\pi t^{1-\gamma}} \int_{-\infty}^{\infty} \frac{1}{(1 + s^2)^{1/2}} \frac{ds}{(t^\gamma + [1 + (s - kt)^2]^{\gamma/2})}. \quad (2.8)$$

Let $s = ty$. Then the integral on the right side of (2.8) can be written as

$$\begin{aligned} & \frac{ct}{2\pi t^{1-\gamma}} \int_{-\infty}^{\infty} \frac{dy}{(1 + t^2 y^2)^{1/2}} \frac{1}{(t^\gamma + [1 + t^2(y - k)^2]^{\gamma/2})} \\ &= \frac{c}{2\pi} \int_{-\infty}^{\infty} \frac{dy}{(1 + t^2 y^2)^{1/2}} \frac{1}{(1 + [t^{-2} + (y - k)^2]^{\gamma/2})} \\ &\leq \frac{c}{2\pi} \int_{-\infty}^{\infty} \frac{dy}{(1 + t^2 y^2)^{1/2}} \frac{1}{[1 + (y - k)^\gamma]} \rightarrow 0, \text{ as } t \rightarrow \infty, \end{aligned} \quad (2.9)$$

and the convergence of the last integral to zero is uniform with respect to $k \in \mathbb{R}$.

Thus

$$\lim_{t \rightarrow \infty} \|I\| = 0. \quad (2.10)$$

From (2.3)-(2.5) the last two conclusions of Theorem 1.3 follow. Theorem 1.3 is proved. \square

2.2. Proof of Theorem 1.4

Using Proposition 1.2, equation (1.6), and the Mellin formula, one gets

$$\frac{1}{t} \int_0^t u(t) e^{ikt} dt = \frac{1}{t} \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \frac{\mathcal{U}(p-ik)}{p} e^{pt} dp, \quad (2.11)$$

where, according to (1.6),

$$\mathcal{U}(p-ik) = \frac{\mathcal{W}(p-ik)}{p}. \quad (2.12)$$

Let $\sigma = \frac{1}{t}$ and $pt = q$. Then

$$\frac{1}{t} \int_0^t u(t) e^{ikt} dt = \frac{1}{2\pi i} \int_{1-i\infty}^{1+i\infty} \mathcal{W}\left(\frac{q}{t} - ik\right) \frac{e^q}{q^2} dq. \quad (2.13)$$

Estimate (1.15) and Theorem 1.3 imply that all $b_m = 0$ in formula (2.1). Therefore, using formula (2.1) with $b_m = 0$, one gets

$$\mathcal{W} = \sum_{j=1}^J v_j \frac{1}{p + ik_j} + \mathcal{W}_1,$$

and

$$\mathcal{W}\left(\frac{q}{t} - ik\right) = \mathcal{W}_1\left(\frac{q}{t} - ik\right) + \sum_{j=1}^J v_j \frac{1}{\frac{q}{t} - i(k - k_j)}. \quad (2.14)$$

One has $\bar{t}^n = \frac{n!}{p^{n+1}}$. Therefore

$$\frac{1}{2\pi i} \int_{1-i\infty}^{1+i\infty} \frac{e^q}{q^2} dq = 1,$$

and

$$\lim_{t \rightarrow \infty} \frac{v_j}{2\pi i} \int_{1-i\infty}^{1+i\infty} \frac{1}{\frac{q}{t} - i(k - k_j)} \frac{e^q}{q^2} dq = \begin{cases} \frac{iv_j}{k - k_j}, & k \neq k_j, \\ \infty, & k = k_j. \end{cases} \quad (2.15)$$

Furthermore,

$$\lim_{t \rightarrow \infty} \frac{1}{2\pi i} \int_{1-i\infty}^{1+i\infty} \mathcal{W}_1\left(\frac{q}{t} - ik\right) \frac{e^q}{q^2} dq = \mathcal{W}_1(-ik), \quad (2.16)$$

as follows from assumption (1.14) and the Lebesgue's dominated convergence theorem if one passes to the limit $t \rightarrow \infty$ under the sign of the integral (2.16). Let us check that this v solves equation (1.3). This would conclude the proof of Theorem 1.4. We need a lemma.

Lemma 2.1. *If $h \in L^1_{loc}(0, \infty)$ and the limit $\lim_{t \rightarrow \infty} t^{-1} \int_0^t h(s) ds$ exists, then the limit $\lim_{p \rightarrow 0} p \int_0^\infty e^{-pt} h(t) dt$ exists, and*

$$\lim_{t \rightarrow \infty} t^{-1} \int_0^t h(s) ds = \lim_{p \rightarrow 0} p \int_0^\infty e^{-pt} h(t) dt. \quad (2.17)$$

Proof of Lemma 1. One has

$$p \int_0^\infty e^{-pt} h(t) dt = pe^{-pt} \int_0^t h(s) ds|_0^\infty + p^2 \int_0^\infty te^{-pt} t^{-1} \int_0^t h(s) ds dt.$$

For any $p > 0$ one has

$$pe^{-pt} \int_0^t h(s) ds|_0^\infty = 0.$$

Let $q = pt$ and denote $H(t) := t^{-1} \int_0^t h(s) ds$, $J := \lim_{t \rightarrow \infty} H(t)$. Then

$$\lim_{p \rightarrow 0} p^2 \int_0^\infty te^{-pt} t^{-1} \int_0^t h(s) ds dt = \lim_{p \rightarrow 0} \int_0^\infty qe^{-q} H(qp^{-1}) dq.$$

Passing in the last integral to the limit $p \rightarrow 0$ one obtains (2.17). Lemma 1 is proved. \square

Using equation (2.17), one writes $v = \lim_{p \rightarrow 0} p\mathcal{U}(p - ik)$, where \mathcal{U} solves equation (1.6). Thus,

$$L\mathcal{U}(p - ik) + (p - ik)^2 \mathcal{U}(p - ik) = p^{-1} f.$$

Multiplying both sides of this equation by p and passing to the limit $p \rightarrow 0$, one obtains equation (1.3). In the passage to the limit under the sign of the unbounded operator L the assumption that L is closed was used.

Thus, the conclusion of Theorem 1.4 follows. \square

If the limit (1.14) exists at a point $p = i\tau$ then one says that the limiting absorption principle holds for the operator L at the point $k = ip = i(-ik) = k$, $k > 0$.

Thus, Assumption B means that the limiting absorption principle holds for L at the point $k > 0$, that is, $\lim_{\epsilon \rightarrow 0} (L - k^2 - i\epsilon)^{-1} f$ exists.

3. Applications

Let $L = -\nabla^2 + q(x)$, where $q(x)$ is a real-valued function, $|q(x)| \leq c(1 + |x|)^{-2-\epsilon}$, $\epsilon > 0$, $x \in \mathbb{R}^3$. Then L is selfadjoint on the domain $H^2(\mathbb{R}^3)$. Its resolvent $(L - k^2 - i0)^{-1}$ satisfies Assumptions A and B if one keeps in mind the following.

Let $G(x, y, k)$ be the resolvent kernel of L , that is, the kernel of the operator $(L - k^2 - i0)^{-1}$,

$$LG(x, y, k) = -\delta(x - y) \quad \text{in } \mathbb{R}^3,$$

$G \in L^2(\mathbb{R}^3)$ for $\text{Im } k > 0$. If $f \in L^2(\mathbb{R}^3)$ is compactly supported, then for $k > 0$ the function

$$v(x) := (L - k^2 - i0)^{-1}f = \int_{\mathbb{R}^3} G(x, y, k)f(y)dy$$

does not necessarily belong to $L^2(\mathbb{R}^3)$.

For example, if $q(x) = 0$, then $G(x, y, k) = \frac{e^{ik|x-y|}}{4\pi|x-y|}$, and the function

$$v(x, k) = \int_{|y| \leq 1} g(x, y, k)dy = O\left(\frac{1}{|x|}\right) \quad (3.1)$$

does not belong to $L^2(\mathbb{R}^3)$ (except for those $k > 0$ for which $x(x, k) = 0$ in the region $|y| \geq 1$. These numbers $k > 0$ are the zeros of the Fourier transform of the characteristic function of the ball $|y| \leq 1$, see [10], Chapter 11.

By this reason the abstract results of theorem (1.3) and (1.4) can be used in applications if one defines some subspace of H , for example, a subspace of functions with compact support, denote by \mathcal{P} , a projection operator on this subspace, and replaces \mathcal{W} and \mathcal{W}_1 by $\mathcal{P}\mathcal{W}$ and $\mathcal{P}\mathcal{W}_1$ in equations (1.12) and (1.14). For example, the function (3.1) one replaces by $\eta(x)v(x, k)$, where $\eta(x)$ is a characteristic function of a compact subset of \mathbb{R}^3 .

The analytic properties of $\eta(x)v(x, k)$ and of $v(x, k)$ as functions of k are the same. A similar suggestion is used in [1].

With the above in mind, one knows (for example, from [2] or [11]) that Assumptions A and B hold for $L = -\nabla^2 + q(x)$.

Consequently, the conclusions of Theorems 1.3 and 1.4 hold.

In addition, the assumptions

$$|q(x)| \leq c(1 + |x|)^{-2-\epsilon}, \quad \epsilon > 0, \quad \text{Im } q = 0,$$

imply that L does not have positive eigenvalues, so all $v_j = 0$, and zero is not an eigenvalue of $L \geq 0$ if $\epsilon > 0$ (see [5], [6]).

A new method for estimating of large time behavior of solutions to abstract evolution problems is developed in [9], where some applications of this method are given.

References

- [1] D. Eidus. *The limiting amplitude principle*. Russ. Math. Surveys, 24, (1969), 51–94
- [2] D. Pearson. *Quantum scattering and spectral theory*. Acad. Press, London, 1988.
- [3] A.G. Ramm. *Spectral properties of the Schrödinger operator in some domains with infinite boundaries*. Doklady Acad of Sci. USSR, 152, (1963) 282-285.
- [4] A.G. Ramm. *Spectral properties of the Schrödinger operator in some infinite domains*. Mat. Sborn., 66, (1965), 321-343.
- [5] A.G. Ramm. *Sufficient conditions for zero not to be an eigenvalue of the Schrödinger operator*. J. Math. Phys., 28, (1987), 1341–1343.
- [6] A.G. Ramm. *Conditions for zero not to be an eigenvalue of the Schrödinger operator*. J. Math. Phys. 29, (1988), 1431–1432.
- [7] A.G. Ramm. *Eigenfunction expansion for nonselfadjoint Schrödinger operator*. Doklady Acad. Sci. USSR. 191, (1970), 50-53.

- [8] A.G. Ramm. *Scattering by obstacles*. D.Reidel, Dordrecht, 1986.
- [9] A.G. Ramm. *Stability of solutions to some evolution problems*. Chaotic Modeling and Simulation (CMSIM), 1, (2011), 17-27.
- [10] A.G. Ramm. *Inverse problems*. Springer, New York, 2005.
- [11] M. Schechter. *Operator methods in quantum mechanics*. North Holland, New York, 1981.